

2-Chloro-N-(4-fluorophenyl)acetamide

Si-shun Kang, Hai-su Zeng, Hai-lin Li and Hai-bo Wang*

College of Science, Nanjing University of Technology, Xinmofan Road No. 5
Nanjing, Nanjing 210009, People's Republic of China
Correspondence e-mail: wanghaibo@njut.edu.cn


Received 27 May 2008; accepted 28 May 2008

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$;
 R factor = 0.047; wR factor = 0.126; data-to-parameter ratio = 8.4.

In the title compound, $\text{C}_8\text{H}_7\text{ClFNO}$, an intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond forms a six-membered ring. In the crystal structure, molecules are linked by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming infinite chains along the c axis.

Related literature

For related compounds, see: Wen *et al.* (2006); Zhang *et al.* (2006). For reference structural data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_8\text{H}_7\text{ClFNO}$	$V = 842.7(3)\text{ \AA}^3$
$M_r = 187.60$	$Z = 4$
Monoclinic, Cc	Mo $K\alpha$ radiation
$a = 4.7410(9)\text{ \AA}$	$\mu = 0.42\text{ mm}^{-1}$
$b = 20.062(4)\text{ \AA}$	$T = 293(2)\text{ K}$
$c = 8.9860(18)\text{ \AA}$	$0.30 \times 0.20 \times 0.05\text{ mm}$
$\beta = 99.60(3)^\circ$	

Data collection

Enraf–Nonius CAD-4	861 independent reflections
diffractometer	610 reflections with $I > 2\sigma(I)$
Absorption correction: ψ scan	$R_{\text{int}} = 0.015$
(North <i>et al.</i> , 1968)	3 standard reflections
$T_{\text{min}} = 0.885$, $T_{\text{max}} = 0.980$	every 200 reflections
974 measured reflections	intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	$\Delta\rho_{\text{max}} = 0.16\text{ e \AA}^{-3}$
$wR(F^2) = 0.126$	$\Delta\rho_{\text{min}} = -0.20\text{ e \AA}^{-3}$
$S = 1.00$	Absolute structure: Flack (1983),
861 reflections	92 Friedel pairs
103 parameters	Flack parameter: 0.18 (17)
	H-atom parameters constrained

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}3-\text{H}3\text{A}\cdots\text{O}$	0.93	2.36	2.925 (8)	119
$\text{N}-\text{H}1\cdots\text{O}^i$	0.86	2.02	2.853 (6)	164

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2738).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Wen, Y. H., Zhang, S. S., Yu, B. H., Li, X. M. & Liu, Q. (2006). *Asian J. Chem.* **18**, 1032–1038.
Zhang, S.-S., Wen, H.-L., Li, X.-M., Xu, L.-L. & Wen, Y.-H. (2006). *Acta Cryst. E* **62**, o3412–o3413.

supplementary materials

Acta Cryst. (2008). E64, o1194 [doi:10.1107/S1600536808016152]

2-Chloro-N-(4-fluorophenyl)acetamide

S. Kang, H. Zeng, H. Li and H. Wang

Comment

N-(substituted phenyl)-2-chloroacetamides are important intermediates in organic synthesis. They can be used in the synthesis of many derivatives such as (quinolin-8-yloxy) acetamide (Zhang *et al.*, 2006) and 2,5-piperazinedione (Wen *et al.*, 2006). In our studies in this area, the title compound, (I), was synthesized and structurally characterised.

The bond lengths and angles in (I) are within normal ranges (Allen *et al.*, 1987). An intramolecular C—H···O interaction occurs (Fig. 1) and an intermolecular N—H···O hydrogen bond helps to establish the packing (Table 1).

Experimental

Chloroacetyl chloride (0.05 mol) was added to a solution of 4-nitrophenylamine (0.05 mol) and triethylamine (0.05 mol) in toluene (50 ml) over a period of 30 min, with cooling in an ice bath, and then the mixture was stirred at room temperature for 4 h. After separation of the triethylamine hydrochloride by filtration, the organic phase was washed three times with water. The toluene layer was removed and evaporated. Pink blocks of (I) were obtained by slow evaporation of a chloroform solution over a period of 7 d.

Refinement

The H atoms were positioned geometrically (N—H = 0.86 Å, C—H = 0.93–0.97 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{carrier})$.

Figures



Fig. 1. The molecular structure of (I) with displacement ellipsoids for the non-hydrogen atoms drawn at the 50% probability level. The intramolecular hydrogen bond is shown as a dashed line.

2-Chloro-N-(4-fluorophenyl)acetamide

Crystal data

C ₈ H ₇ ClFNO	$F_{000} = 384$
$M_r = 187.60$	$D_x = 1.479 \text{ Mg m}^{-3}$
Monoclinic, <i>Cc</i>	Mo $K\alpha$ radiation
	$\lambda = 0.71073 \text{ \AA}$
Hall symbol: C -2yc	Cell parameters from 25 reflections
$a = 4.7410(9) \text{ \AA}$	$\theta = 8\text{--}12^\circ$

supplementary materials

$b = 20.062 (4) \text{ \AA}$	$\mu = 0.42 \text{ mm}^{-1}$
$c = 8.9860 (18) \text{ \AA}$	$T = 293 (2) \text{ K}$
$\beta = 99.60 (3)^\circ$	Block, pink
$V = 842.7 (3) \text{ \AA}^3$	$0.30 \times 0.20 \times 0.05 \text{ mm}$
$Z = 4$	

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.016$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.2^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.0^\circ$
$T = 293(2) \text{ K}$	$h = 0 \rightarrow 5$
$\omega/2\theta$ scans	$k = 0 \rightarrow 24$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -10 \rightarrow 10$
$T_{\text{min}} = 0.885, T_{\text{max}} = 0.980$	3 standard reflections
974 measured reflections	every 200 reflections
861 independent reflections	intensity decay: none
610 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.046$	$w = 1/[\sigma^2(F_o^2) + (0.P)^2 + 0.5P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.126$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.16 \text{ e \AA}^{-3}$
861 reflections	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
103 parameters	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 92 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.18 (17)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl	-0.4033 (4)	0.15199 (10)	0.5685 (2)	0.1096 (7)
N	0.1497 (11)	0.2970 (2)	0.6731 (5)	0.0738 (14)
H1	0.2033	0.2848	0.7653	0.089*
O	-0.1268 (10)	0.2652 (2)	0.4535 (5)	0.084
F	0.6949 (14)	0.5279 (2)	0.5602 (6)	0.147 (2)
C1	0.5590 (19)	0.4687 (3)	0.5826 (8)	0.095 (2)
C2	0.3302 (19)	0.4493 (4)	0.4775 (8)	0.097 (2)
H2A	0.2684	0.4749	0.3920	0.116*
C3	0.1935 (15)	0.3901 (3)	0.5030 (6)	0.0817 (18)
H3A	0.0450	0.3743	0.4308	0.098*
C4	0.2759 (13)	0.3546 (3)	0.6344 (6)	0.0707 (15)
C5	0.5091 (15)	0.3787 (3)	0.7356 (7)	0.0798 (17)
H5A	0.5715	0.3539	0.8224	0.096*
C6	0.6503 (19)	0.4357 (4)	0.7157 (8)	0.099 (2)
H6A	0.7995	0.4516	0.7874	0.119*
C7	-0.0366 (13)	0.2576 (3)	0.5950 (5)	0.0710 (16)
C8	-0.1284 (15)	0.1998 (3)	0.6748 (6)	0.089 (2)
H8A	-0.1937	0.2153	0.7654	0.107*
H8B	0.0358	0.1712	0.7058	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl	0.1293 (15)	0.1275 (15)	0.0742 (9)	-0.0277 (13)	0.0237 (9)	-0.0087 (10)
N	0.089 (3)	0.082 (3)	0.051 (2)	0.010 (3)	0.013 (2)	0.004 (2)
O	0.084	0.084	0.084	0.000	0.014	0.000
F	0.213 (7)	0.120 (3)	0.121 (3)	-0.062 (4)	0.067 (4)	0.001 (3)
C1	0.117 (6)	0.091 (5)	0.086 (5)	-0.031 (5)	0.043 (5)	0.001 (4)
C2	0.121 (6)	0.103 (5)	0.074 (4)	0.000 (5)	0.039 (4)	0.017 (4)
C3	0.090 (4)	0.096 (5)	0.063 (3)	-0.006 (4)	0.025 (3)	0.002 (3)
C4	0.078 (4)	0.080 (4)	0.058 (3)	0.008 (3)	0.020 (3)	0.008 (3)
C5	0.095 (4)	0.083 (4)	0.067 (3)	-0.016 (4)	0.029 (3)	0.000 (3)
C6	0.115 (6)	0.116 (5)	0.074 (4)	-0.007 (5)	0.038 (4)	0.010 (4)
C7	0.072 (3)	0.102 (4)	0.039 (2)	-0.003 (3)	0.009 (2)	-0.011 (3)
C8	0.111 (5)	0.110 (5)	0.044 (3)	-0.017 (4)	0.005 (3)	0.013 (3)

Geometric parameters (\AA , $^\circ$)

Cl—C8	1.765 (7)	C3—C4	1.379 (8)
N—C7	1.300 (7)	C3—H3A	0.9300
N—C4	1.373 (8)	C4—C5	1.396 (9)
N—H1	0.8600	C5—C6	1.353 (10)
O—C7	1.281 (6)	C5—H5A	0.9300
F—C1	1.381 (7)	C6—H6A	0.9300
C1—C2	1.371 (10)	C7—C8	1.467 (8)

supplementary materials

C1—C6	1.372 (10)	C8—H8A	0.9700
C2—C3	1.390 (9)	C8—H8B	0.9700
C2—H2A	0.9300		
C7—N—C4	131.3 (5)	C6—C5—C4	124.2 (6)
C7—N—H1	114.3	C6—C5—H5A	117.9
C4—N—H1	114.3	C4—C5—H5A	117.9
C2—C1—C6	124.2 (7)	C5—C6—C1	115.6 (7)
C2—C1—F	118.6 (7)	C5—C6—H6A	122.2
C6—C1—F	117.0 (7)	C1—C6—H6A	122.2
C1—C2—C3	117.7 (6)	O—C7—N	123.2 (6)
C1—C2—H2A	121.1	O—C7—C8	120.1 (5)
C3—C2—H2A	121.1	N—C7—C8	116.5 (4)
C4—C3—C2	120.7 (6)	C7—C8—Cl	114.7 (4)
C4—C3—H3A	119.7	C7—C8—H8A	108.6
C2—C3—H3A	119.7	Cl—C8—H8A	108.6
N—C4—C3	125.4 (6)	C7—C8—H8B	108.6
N—C4—C5	117.3 (5)	Cl—C8—H8B	108.6
C3—C4—C5	117.3 (6)	H8A—C8—H8B	107.6
C6—C1—C2—C3	-4.3 (12)	C3—C4—C5—C6	2.8 (10)
F—C1—C2—C3	-179.1 (7)	C4—C5—C6—C1	-3.0 (11)
C1—C2—C3—C4	3.9 (11)	C2—C1—C6—C5	3.8 (12)
C7—N—C4—C3	11.2 (11)	F—C1—C6—C5	178.7 (7)
C7—N—C4—C5	-168.0 (7)	C4—N—C7—O	4.7 (11)
C2—C3—C4—N	177.7 (6)	C4—N—C7—C8	-179.3 (6)
C2—C3—C4—C5	-3.1 (10)	O—C7—C8—Cl	-9.0 (9)
N—C4—C5—C6	-177.9 (7)	N—C7—C8—Cl	174.8 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C3—H3A···O	0.93	2.36	2.925 (8)	119
N—H1···O ⁱ	0.86	2.02	2.853 (6)	164

Symmetry codes: (i) $x+1/2, -y+1/2, z+1/2$.

Fig. 1

