

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**(±)-1-(1*H*-Benzimidazol-2-yl)ethanol**

Rong Xia and Hai-Jun Xu\*

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China  
Correspondence e-mail: xuhj@seu.edu.cn


Received 18 April 2008; accepted 9 May 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.059;  $wR$  factor = 0.124; data-to-parameter ratio = 10.0.

The asymmetric unit of the title molecule,  $\text{C}_9\text{H}_{10}\text{N}_2\text{O}$ , contains two molecules. The fused benzene and imidazole rings are nearly coplanar, the largest deviations from the mean plane being 0.025 (3) Å at the non-bridgehead imidazole C atom of one molecule and 0.018 (3) Å at one of the bridgehead C atoms in the other molecule. Intermolecular  $\text{O}-\text{H}\cdots\text{N}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds result in the formation of a sheet parallel to the (010) plane.

## Related literature

For related literature, see: Allen *et al.* (1987); Chen & Ruan (2007); Garuti *et al.* (1999); Matsuno *et al.* (2000); Tlahuext *et al.* (2007).



## Experimental

## Crystal data

 $\text{C}_9\text{H}_{10}\text{N}_2\text{O}$  $M_r = 162.19$ Orthorhombic,  $P2_12_12$  $a = 13.734$  (3) Å $b = 15.376$  (3) Å $c = 7.9163$  (16) Å $V = 1671.7$  (6) Å<sup>3</sup> $Z = 8$ Mo  $K\alpha$  radiation $\mu = 0.09$  mm<sup>-1</sup> $T = 293$  (2) K

0.20 × 0.18 × 0.05 mm

## Data collection

Rigaku Mercury2 diffractometer  
Absorption correction: multi-scan  
(*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.912$ ,  $T_{\max} = 1.00$   
(expected range = 0.908–0.996)

17487 measured reflections  
2199 independent reflections  
1380 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.115$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$   
 $wR(F^2) = 0.123$   
 $S = 1.07$   
2199 reflections

221 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.17$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.18$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N3}$	0.82	1.91	2.713 (4)	165
$\text{N4}-\text{H4}\cdots\text{O1}^{\text{i}}$	0.86	1.97	2.828 (4)	178
$\text{O2}-\text{H2}\cdots\text{N2}^{\text{ii}}$	0.82	1.93	2.743 (4)	170
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{iii}}$	0.86	1.93	2.751 (4)	160

Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ ; (ii)  $x, y, z - 1$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-III* (Burnett & Johnson, 1996), *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97*.

HJX acknowledges a Start-up Grant from Southeast University, P. R. China

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2341).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEP-III*. Report ORNL-6895. Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.
- Chen, J. & Ruan, Y. (2007). *Acta Cryst.* **E63**, m2964.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Garuti, L., Roberti, M. & Cermelli, C. (1999). *Bioorg. Med. Chem. Lett.* **9**, 2525–2530.
- Matsuno, T., Kato, M., Sasahara, H., Watanabe, T., Inaba, M., Takahashi, M., Yaguchi, S. I., Yoshioka, K., Sakato, M. & Kawashima, S. (2000). *Chem. Pharm. Bull.* **48**, 1778–1781.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Tlahuext, H., Tlahuext, M., López-Gómez, S. & Tapia-Benavides, A. R. (2007). *Acta Cryst.* **E63**, m1263–m1265.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1223 [ doi:10.1107/S1600536808013871 ]

## (±)-1-(1*H*-Benzimidazol-2-yl)ethanol

R. Xia and H.-J. Xu

### Comment

Imidazole and benzimidazole derivatives are important heteroaromatic compounds and have attracted considerable attention because of good biological and pharmaceutical activities (Matsuno *et al.*, 2000; Garuti *et al.*, 1999). These compounds also play an important role in the development of coordination chemistry. Many derivatives of benzimidazole have been prepared and their complexes have been studied (Tlahuext *et al.*, 2007; Chen & Ruan, 2007). In this paper, we report the crystal structure of the title compound.

There are two crystallographically independent molecules, A and B, linked by a O-H $\cdots$ N hydrogen bond in the asymmetric unit. The bond lengths and angles in A and B are within normal ranges (Allen *et al.*, 1987). The two fused benzene and imidazole rings are nearly planar with the largest deviations from the mean plane being 0.025 (3) Å at C7 and 0.018 (3) Å at C10. These two fused rings make a dihedral angle of 35.01 (9)°.

The molecules are further connected through O-H $\cdots$ N and N-H $\cdots$ O hydrogen bond building up a two dimensional network which is parallel to the (0 1 0) plane (Table 1, Fig. 2).

Only the relative absolute configuration could be determined, the C8 and C17 have the same absolute configuration (S,S) or (R,R). The (S,S) configuration is represented in Fig. 1.

### Experimental

All chemicals were obtained from commercial sources and used directly without further purification. Benzene-1, 2-diamine (2.16 g, 20 mmol) was dissolved in hydrochloric acid (25 mL, 4 M) at 100°C, and ethyl 2-hydroxypropanoate (2.48 g, 21 mmol) was added to the solution. The mixture were then heated to reflux for 7 h at 115°C. After cooling to room temperature, the product was divided by neutralizing the mixture solution using NaOH to make the pH 7–9. Solid product was collected by filtration and the yield was 80%. <sup>1</sup>H-NMR(CDCl<sub>3</sub>, 300 MHz): δ 1.72 (d, 3 H), 5.22 (q, 1 H), 7.47(m, 1 H), 7.58 (m, 2 H), 7.81 (m, 1 H). Esi-MS: calcd for C<sub>14</sub>H<sub>9</sub>N<sub>2</sub>O–H *m/z* 161.19, found 161.18. Deep-red single crystals of the title compound suitable for X-ray diffraction analysis were obtained from methanol solution by slow evaporation after a week.

### Refinement

All H atoms attached to C, O and N atom were fixed geometrically and treated as riding with C—H = 0.98 Å (methine), 0.96 Å (methyl or 0.93 Å (aromatic), O—H = 0.82 Å and N—H = 0.86 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$  or  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}}, \text{O})$ .

In the absence of significant anomalous scattering, the absolute configuration could not be reliably determined and then the Friedel pairs were merged and any references to the Flack parameter were removed.

## Figures




Fig. 1. A view of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen bond is shown as dashed line. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Partial packing view showing the hydrogen bonds network. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bondings have been omitted for clarity. [Symmetry codes: (i)  $x-1/2, -y+1/2, -z+1$ ; (ii)  $x, y, z-1$ ; (iii)  $x+1/2, -y+1/2, -z+1$  ]

## (±)-1-(1H-Benzimidazol-2-yl)ethanol

### Crystal data

$C_9H_{10}N_2O$

$M_r = 162.19$

Orthorhombic,  $P2_12_12$

Hall symbol: P 2 2ab

$a = 13.734$  (3) Å

$b = 15.376$  (3) Å

$c = 7.9163$  (16) Å

$V = 1671.7$  (6) Å<sup>3</sup>

$Z = 8$

$F_{000} = 688$

$D_x = 1.289$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 10749 reflections

$\theta = 2.4$ – $28.0^\circ$

$\mu = 0.09$  mm<sup>-1</sup>

$T = 293$  (2) K

Block, red

$0.20 \times 0.18 \times 0.05$  mm

### Data collection

Rigaku Mercury2  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 13.6612 pixels mm<sup>-1</sup>

$T = 293$ (2) K

$\omega$  scans

Absorption correction: multi-scan  
(CrystalClear; Rigaku, 2005)

$T_{\min} = 0.912$ ,  $T_{\max} = 1.00$

17487 measured reflections

2199 independent reflections

1380 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.115$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 3.0^\circ$

$h = -17 \rightarrow 17$

$k = -19 \rightarrow 19$

$l = -10 \rightarrow 10$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.059$	H-atom parameters constrained
$wR(F^2) = 0.123$	$w = 1/[\sigma^2(F_o^2) + (0.0395P)^2 + 0.22P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
2199 reflections	$(\Delta/\sigma)_{\max} < 0.001$
221 parameters	$\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6823 (3)	0.3355 (2)	1.1332 (4)	0.0383 (9)
C2	0.7541 (3)	0.3880 (2)	1.1995 (5)	0.0496 (10)
H2A	0.8168	0.3885	1.1551	0.060*
C3	0.7292 (3)	0.4388 (3)	1.3323 (5)	0.0605 (12)
H3	0.7760	0.4750	1.3801	0.073*
C4	0.6362 (4)	0.4384 (3)	1.3984 (6)	0.0652 (13)
H4A	0.6219	0.4740	1.4900	0.078*
C5	0.5647 (3)	0.3868 (3)	1.3328 (6)	0.0622 (12)
H5	0.5021	0.3872	1.3774	0.075*
C6	0.5887 (3)	0.3340 (2)	1.1975 (5)	0.0427 (9)
C7	0.5899 (3)	0.2474 (2)	0.9864 (5)	0.0428 (9)
C8	0.5593 (3)	0.1874 (3)	0.8498 (5)	0.0520 (11)
H8	0.4882	0.1823	0.8527	0.062*
C9	0.6018 (4)	0.0985 (3)	0.8690 (6)	0.0919 (18)
H9A	0.5832	0.0632	0.7743	0.138*
H9B	0.5781	0.0725	0.9713	0.138*
H9C	0.6715	0.1027	0.8737	0.138*
C10	0.3684 (3)	0.3451 (2)	0.6778 (5)	0.0423 (9)

## supplementary materials

---

C11	0.4085 (3)	0.3972 (3)	0.8015 (5)	0.0563 (11)
H11	0.4721	0.3888	0.8385	0.068*
C12	0.3517 (3)	0.4614 (3)	0.8676 (5)	0.0614 (12)
H12	0.3768	0.4974	0.9514	0.074*
C13	0.2571 (3)	0.4740 (3)	0.8120 (5)	0.0576 (11)
H13	0.2203	0.5186	0.8591	0.069*
C14	0.2166 (3)	0.4234 (2)	0.6914 (5)	0.0511 (10)
H14	0.1527	0.4317	0.6561	0.061*
C15	0.2740 (3)	0.3592 (2)	0.6231 (4)	0.0401 (9)
C16	0.3419 (2)	0.2520 (2)	0.4833 (4)	0.0380 (8)
C17	0.3509 (3)	0.1825 (2)	0.3527 (4)	0.0463 (10)
H17	0.2934	0.1450	0.3628	0.056*
C18	0.4389 (3)	0.1262 (3)	0.3790 (6)	0.0581 (11)
H18A	0.4414	0.0826	0.2924	0.087*
H18B	0.4349	0.0986	0.4876	0.087*
H18C	0.4965	0.1614	0.3739	0.087*
N1	0.68087 (19)	0.27939 (19)	0.9990 (3)	0.0451 (8)
H1A	0.7292	0.2669	0.9345	0.054*
N2	0.5316 (2)	0.2773 (2)	1.1042 (4)	0.0486 (8)
N3	0.4093 (2)	0.27755 (19)	0.5867 (4)	0.0443 (8)
N4	0.25902 (19)	0.29851 (18)	0.4998 (4)	0.0425 (7)
H4	0.2063	0.2912	0.4429	0.051*
O1	0.58658 (16)	0.2215 (2)	0.6915 (3)	0.0532 (7)
H1	0.5385	0.2415	0.6440	0.080*
O2	0.34919 (16)	0.21938 (19)	0.1892 (3)	0.0546 (7)
H2	0.4018	0.2426	0.1697	0.082*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.041 (2)	0.046 (2)	0.0280 (19)	-0.0019 (17)	0.0020 (17)	0.0067 (17)
C2	0.042 (2)	0.060 (3)	0.047 (2)	-0.0057 (19)	0.000 (2)	0.008 (2)
C3	0.065 (3)	0.065 (3)	0.052 (3)	-0.006 (2)	-0.009 (3)	-0.005 (2)
C4	0.086 (4)	0.060 (3)	0.050 (3)	0.001 (3)	0.002 (3)	-0.009 (2)
C5	0.059 (3)	0.073 (3)	0.055 (3)	0.000 (2)	0.018 (2)	-0.008 (3)
C6	0.041 (2)	0.051 (2)	0.036 (2)	-0.0025 (18)	0.0054 (19)	0.0037 (19)
C7	0.039 (2)	0.052 (2)	0.037 (2)	-0.0042 (19)	-0.0017 (19)	0.0043 (18)
C8	0.047 (2)	0.063 (3)	0.046 (2)	-0.0012 (19)	-0.002 (2)	0.000 (2)
C9	0.135 (5)	0.062 (3)	0.078 (4)	0.008 (3)	-0.040 (4)	-0.005 (3)
C10	0.042 (2)	0.044 (2)	0.040 (2)	-0.0017 (17)	-0.007 (2)	0.0045 (18)
C11	0.057 (3)	0.057 (3)	0.055 (3)	0.005 (2)	-0.015 (2)	-0.003 (2)
C12	0.076 (3)	0.057 (3)	0.051 (3)	-0.002 (2)	-0.009 (3)	-0.003 (2)
C13	0.064 (3)	0.056 (3)	0.053 (3)	0.003 (2)	0.006 (3)	-0.006 (2)
C14	0.043 (2)	0.054 (3)	0.057 (3)	0.0029 (19)	0.005 (2)	0.003 (2)
C15	0.040 (2)	0.048 (2)	0.0323 (19)	0.0002 (18)	-0.0009 (18)	0.0053 (18)
C16	0.0317 (19)	0.042 (2)	0.040 (2)	-0.0067 (16)	-0.0017 (18)	0.0069 (17)
C17	0.041 (2)	0.055 (2)	0.042 (2)	-0.0082 (18)	-0.0007 (19)	-0.0018 (19)
C18	0.053 (2)	0.056 (3)	0.065 (3)	0.009 (2)	-0.005 (2)	-0.005 (2)

N1	0.0335 (16)	0.066 (2)	0.0355 (16)	0.0029 (15)	0.0070 (15)	-0.0025 (17)
N2	0.0388 (17)	0.063 (2)	0.0443 (17)	-0.0022 (16)	0.0071 (15)	0.0033 (17)
N3	0.0394 (17)	0.048 (2)	0.0453 (17)	0.0020 (16)	-0.0106 (16)	-0.0019 (16)
N4	0.0292 (15)	0.0576 (19)	0.0407 (16)	0.0002 (15)	-0.0056 (15)	0.0033 (17)
O1	0.0353 (14)	0.088 (2)	0.0368 (14)	0.0083 (14)	-0.0012 (13)	0.0011 (16)
O2	0.0335 (14)	0.089 (2)	0.0413 (14)	-0.0056 (14)	-0.0032 (13)	-0.0031 (15)

*Geometric parameters (Å, °)*

C1—N1	1.368 (4)	C10—C15	1.385 (5)
C1—C2	1.379 (5)	C11—C12	1.364 (5)
C1—C6	1.383 (5)	C11—H11	0.9300
C2—C3	1.354 (5)	C12—C13	1.385 (5)
C2—H2A	0.9300	C12—H12	0.9300
C3—C4	1.381 (6)	C13—C14	1.352 (5)
C3—H3	0.9300	C13—H13	0.9300
C4—C5	1.364 (6)	C14—C15	1.374 (5)
C4—H4A	0.9300	C14—H14	0.9300
C5—C6	1.384 (5)	C15—N4	1.366 (4)
C5—H5	0.9300	C16—N3	1.297 (4)
C6—N2	1.386 (5)	C16—N4	1.350 (4)
C7—N2	1.312 (4)	C16—C17	1.492 (5)
C7—N1	1.347 (4)	C17—O2	1.413 (4)
C7—C8	1.482 (5)	C17—C18	1.501 (5)
C8—O1	1.410 (4)	C17—H17	0.9800
C8—C9	1.494 (6)	C18—H18A	0.9600
C8—H8	0.9800	C18—H18B	0.9600
C9—H9A	0.9600	C18—H18C	0.9600
C9—H9B	0.9600	N1—H1A	0.8600
C9—H9C	0.9600	N4—H4	0.8600
C10—C11	1.379 (5)	O1—H1	0.8200
C10—N3	1.384 (4)	O2—H2	0.8200
N1—C1—C2	132.5 (3)	C10—C11—H11	121.2
N1—C1—C6	105.2 (3)	C11—C12—C13	121.1 (4)
C2—C1—C6	122.3 (3)	C11—C12—H12	119.5
C3—C2—C1	116.9 (4)	C13—C12—H12	119.5
C3—C2—H2A	121.5	C14—C13—C12	122.0 (4)
C1—C2—H2A	121.5	C14—C13—H13	119.0
C2—C3—C4	121.7 (4)	C12—C13—H13	119.0
C2—C3—H3	119.1	C13—C14—C15	117.1 (4)
C4—C3—H3	119.1	C13—C14—H14	121.5
C5—C4—C3	121.6 (4)	C15—C14—H14	121.5
C5—C4—H4A	119.2	N4—C15—C14	133.3 (4)
C3—C4—H4A	119.2	N4—C15—C10	105.0 (3)
C4—C5—C6	117.6 (4)	C14—C15—C10	121.8 (4)
C4—C5—H5	121.2	N3—C16—N4	112.4 (3)
C6—C5—H5	121.2	N3—C16—C17	126.5 (3)
C1—C6—C5	119.8 (4)	N4—C16—C17	121.1 (3)
C1—C6—N2	109.9 (3)	O2—C17—C16	110.2 (3)

## supplementary materials

---

C5—C6—N2	130.3 (4)	O2—C17—C18	111.8 (3)
N2—C7—N1	112.6 (3)	C16—C17—C18	112.6 (3)
N2—C7—C8	124.3 (3)	O2—C17—H17	107.3
N1—C7—C8	123.0 (3)	C16—C17—H17	107.3
O1—C8—C7	110.0 (3)	C18—C17—H17	107.3
O1—C8—C9	109.1 (4)	C17—C18—H18A	109.5
C7—C8—C9	112.6 (3)	C17—C18—H18B	109.5
O1—C8—H8	108.4	H18A—C18—H18B	109.5
C7—C8—H8	108.4	C17—C18—H18C	109.5
C9—C8—H8	108.4	H18A—C18—H18C	109.5
C8—C9—H9A	109.5	H18B—C18—H18C	109.5
C8—C9—H9B	109.5	C7—N1—C1	107.5 (3)
H9A—C9—H9B	109.5	C7—N1—H1A	126.2
C8—C9—H9C	109.5	C1—N1—H1A	126.2
H9A—C9—H9C	109.5	C7—N2—C6	104.7 (3)
H9B—C9—H9C	109.5	C16—N3—C10	105.4 (3)
C11—C10—N3	130.1 (3)	C16—N4—C15	107.7 (3)
C11—C10—C15	120.4 (4)	C16—N4—H4	126.2
N3—C10—C15	109.5 (3)	C15—N4—H4	126.2
C12—C11—C10	117.6 (4)	C8—O1—H1	109.5
C12—C11—H11	121.2	C17—O2—H2	109.5

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 $\cdots$ N3	0.82	1.91	2.713 (4)	165
N4—H4 $\cdots$ O1 <sup>i</sup>	0.86	1.97	2.828 (4)	178
O2—H2 $\cdots$ N2 <sup>ii</sup>	0.82	1.93	2.743 (4)	170
N1—H1A $\cdots$ O2 <sup>iii</sup>	0.86	1.93	2.751 (4)	160

Symmetry codes: (i)  $x-1/2, -y+1/2, -z+1$ ; (ii)  $x, y, z-1$ ; (iii)  $x+1/2, -y+1/2, -z+1$ .



Fig. 1




Fig. 2

