

## Cytenamide–butyric acid (1/1)

Andrea Johnston,<sup>a</sup> Alastair J. Florence,<sup>a\*</sup> Francesca J. A. Fabbiani,<sup>b</sup> Kenneth Shankland<sup>c</sup> and Colin T. Bedford<sup>d</sup>

<sup>a</sup>Solid-State Research Group, Strathclyde Institute of Pharmacy and Biomedical Sciences, The John Arbuthnott Building, University of Strathclyde, Glasgow G4 0NR, Scotland, <sup>b</sup>GZG, Department of Crystallography, University of Göttingen, D-37077 Göttingen, Germany, <sup>c</sup>ISIS Facility, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, England, and <sup>d</sup>Department of Chemistry, University College London, London WC1H 0AJ, England  
Correspondence e-mail: alastair.florence@strath.ac.uk


Received 14 May 2008; accepted 13 June 2008

Key indicators: single-crystal X-ray study;  $T = 160$  K; mean  $\sigma(C-C) = 0.002$  Å;  $R$  factor = 0.041;  $wR$  factor = 0.089; data-to-parameter ratio = 18.0.

Cytenamide forms a 1:1 solvate with butyric acid [systematic name: 5*H*-dibenzo[*a,d*]cycloheptatriene-5-carboxamide–butanoic acid (1/1)],  $C_{16}H_{13}NO \cdot C_4H_8O_2$ . The title compound crystallizes with one molecule of cytenamide and one of butyric acid in the asymmetric unit; these molecules are linked by N–H···O and O–H···O hydrogen bonds to form an  $R_2^2(8)$  heterodimer motif. Pairs of adjacent motifs are further connected via N–H···O interactions to form a discrete centrosymmetric assembly.

### Related literature

For details on experimental methods used to obtain the title solvate, see: Davis *et al.* (1964); Florence *et al.* (2003); Florence, Johnston, Fernandes *et al.* (2006). For literature on cytenamide and related molecules, see: Florence, Bedford *et al.* (2008); Cyr *et al.* (1987); Fleischman *et al.* (2003); Florence, Johnston, Price *et al.* (2006); Florence, Leech *et al.* (2006); Bandoli *et al.* (1992); Harrison *et al.* (2006); Leech *et al.* (2007); Florence, Shankland *et al.* (2008). For other related literature, see: Etter (1990); Desiraju & Steiner (1999).



### Experimental

#### Crystal data

$C_{16}H_{13}NO \cdot C_4H_8O_2$   
 $M_r = 323.39$   
Monoclinic,  $P2_1/n$

$a = 5.9351(2)$  Å  
 $b = 16.3595(5)$  Å  
 $c = 17.6738(4)$  Å

$\beta = 98.046(2)^\circ$   
 $V = 1699.15(9)$  Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.09$  mm<sup>-1</sup>  
 $T = 160$  K  
 $0.35 \times 0.15 \times 0.12$  mm

#### Data collection

Oxford Diffraction Gemini diffractometer  
Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2007)  
 $T_{\min} = 0.91$ ,  $T_{\max} = 0.99$

18979 measured reflections  
4069 independent reflections  
2928 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.088$   
 $S = 0.95$   
4069 reflections  
226 parameters  
3 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.38$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N1–H11···O2	0.860 (14)	2.348 (14)	2.8761 (15)	120.0 (10)
N1–H12···O2 <sup>i</sup>	0.898 (13)	2.146 (13)	3.0167 (15)	163.2 (13)
O3–H311···O1 <sup>i</sup>	0.879 (17)	1.698 (17)	2.5658 (13)	168.8 (16)

Symmetry code: (i)  $-x + 2, -y + 1, -z + 1$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2007); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2007); data reduction: *CrysAlis RED* and *SORTAV* (Blessing, 1997); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *Mercury* (Macrae *et al.*, 2006) and *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2008).

The authors thank the Basic Technology programme of the UK Research Councils for funding this work under the project Control and Prediction of the Organic Solid State (<http://www.cpos.org.uk>).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2147).

### References

- Altomare, A., Cascarano, G., Giacovazzo, G., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Bandoli, G., Nicolini, M., Ongaro, A., Volpe, G. & Rubello, A. (1992). *J. Chem. Crystallogr.* **22**, 177–183.
- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Blessing, R. H. (1997). *J. Appl. Cryst.* **30**, 421–426.
- Cyr, T. D., Matsui, F., Sears, R. W., Curran, N. M. & Lovering, E. G. (1987). *J. Assoc. Off. Anal. Chem.* **70**, 836–840.
- Davis, M. A., Winthrop, S. O., Thomas, R. A., Herr, F., Charest, M.-P. & Gaudry, R. (1964). *J. Med. Chem.* **7**, 88–94.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology*. Oxford University Press.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fleischman, S. G., Kuduva, S. S., McMahon, J. A., Moulton, B., Walsh, R. D. B., Rodriguez-Hornedo, N. & Zaworotko, M. J. (2003). *Cryst. Growth Des.* **3**, 909–919.

# organic compounds

---

- Florence, A. J., Baumgartner, B., Weston, C., Shankland, N., Kennedy, A. R., Shankland, K. & David, W. I. F. (2003). *J. Pharm. Sci.* **92**, 1930–1938.
- Florence, A. J., Bedford, C. T., Fabbiani, F. P. A., Shankland, K., Gelbrich, T., Hursthouse, M. B., Shankland, N., Johnston, A. & Fernandes, P. (2008). *CrystEngComm*. doi:10.1039/b719717a.
- Florence, A. J., Johnston, A., Fernandes, P., Shankland, N. & Shankland, K. (2006). *J. Appl. Cryst.* **39**, 922–924.
- Florence, A. J., Johnston, A., Price, S. L., Nowell, H., Kennedy, A. R. & Shankland, N. (2006). *J. Pharm. Sci.* **95**, 1918–1930.
- Florence, A. J., Leech, C. K., Shankland, N., Shankland, K. & Johnston, A. (2006). *CrystEngComm*, **8**, 746–747.
- Florence, A. J., Shankland, K., Gelbrich, T., Hursthouse, M. B., Shankland, N., Johnston, A., Fernandes, P. & Leech, C. K. (2008). *CrystEngComm*, **10**, 26–28.
- Harrison, W. T. A., Yathirajan, H. S. & Anilkumar, H. G. (2006). *Acta Cryst. C* **62**, o240–o242.
- Leech, C. K., Florence, A. J., Shankland, K., Shankland, N. & Johnston, A. (2007). *Acta Cryst. E* **63**, o675–o677.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Oxford Diffraction (2007). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Westrip, S. P. (2008). *publCIF*. In preparation.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o1295–o1296 [doi:10.1107/S1600536808018059]

### Cytenamide-butyric acid (1/1)

**A. Johnston, A. J. Florence, F. J. A. Fabbiani, K. Shankland and C. T. Bedford**

#### Comment

Cytenamide (CYT) is an analogue of carbamazepine (CBZ), a dibenzazepine drug used to control seizures (Cyr *et al.*, 1987). CYT-butyric acid solvate was produced during an automated parallel crystallization study (Florence, Johnston, Fernandes *et al.*, 2006) of CYT as part of a wider investigation that couples automated parallel crystallization with crystal structure prediction methodology to investigate the basic science underlying the solid-state diversity in CBZ (Florence, Johnston, Price *et al.*, 2006; Florence, Leech *et al.*, 2006) and its closely related analogues, CYT (Florence, Bedford *et al.*, 2008), 10,11-dihydrocarbamazepine (Bandoli *et al.*, 1992; Harrison *et al.*, 2006; Leech *et al.*, 2007) and cyheptamide (Florence, Shankland *et al.*, 2008). The sample was identified as a new form using multi-sample foil transmission X-ray powder diffraction analysis (Florence *et al.*, 2003). Subsequent manual recrystallization from a saturated butyric acid solution by slow evaporation at 278 K yielded a sample suitable for single-crystal X-ray diffraction (Fig. 1).

The compound crystallizes in the monoclinic space group  $P2_1/n$  with one CYT and one solvent molecule in the asymmetric unit. The molecules adopt a hydrogen-bonded arrangement similar to that observed in CBZ-butyric acid solvate (1/1) (Fleischman *et al.*, 2003) whereby the CYT and butyric acid molecules are connected *via* N—H···O and O—H···O hydrogen bonds to form an  $R_2^2(8)$  dimer motif (Etter, 1990). Adjacent dimers are linked *via* a third contact (N1—H1···O2; Fig 2) to form an  $R_4^2(8)$  centrosymmetric double motif arrangement. The O1···O3 distance of 2.566 (1) Å lies within the expected range for strong hydrogen bonds (2.5–3.2 Å; Desiraju and Steiner, 1999).

CYT-butyric acid solvate structure reported here is essentially isostructural with both CBZ-formic acid and CBZ-acetic acid solvates (Fleischman *et al.*, 2003).

#### Experimental

A sample of cytenamide was synthesized according to a modification of the published method (Davis *et al.*, 1964). A single-crystal sample of cytenamide-butyric acid was grown from a saturated butyric acid solution by isothermal solvent evaporation at 278 K.

#### Refinement

H-atoms were found on a difference Fourier map and were initially refined with soft restraints on the bond lengths and angles to regularize their geometry and  $U_{\text{iso}}(\text{H})$  (in the range 1.2–1.5 times  $U_{\text{eq}}$  of the parent atom), after which the positions were refined with riding constraints. The positions of H-atoms involved in H-bonding were refined subject to distance restraints.

# supplementary materials

---

## Figures




Fig. 1. The molecular structure of CYT–butyric acid (1/1), showing 50% probability displacement ellipsoids.




Fig. 2. The hydrogen bonded  $R_2^{2}(8)$  motifs of CYT–butyric acid joined in a centrosymmetric arrangement *via* an  $R_4^{2}(8)$  motif. Hydrogen bonds are shown as dashed lines.

## 5H-dibenzo[a,d]cycloheptatriene-5-carboxamide–butanoic acid (1/1)

### Crystal data

$C_{16}H_{13}NO \cdot C_4H_8O_2$	$F_{000} = 688$
$M_r = 323.39$	$D_x = 1.264 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 216.2 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation
$a = 5.9351 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 16.3595 (5) \text{ \AA}$	Cell parameters from 6486 reflections
$c = 17.6738 (4) \text{ \AA}$	$\theta = 3\text{--}29^\circ$
$\beta = 98.046 (2)^\circ$	$\mu = 0.09 \text{ mm}^{-1}$
$V = 1699.15 (9) \text{ \AA}^3$	$T = 160 \text{ K}$
$Z = 4$	Block, colourless
	$0.35 \times 0.15 \times 0.12 \text{ mm}$

### Data collection

Oxford Diffraction Gemini diffractometer	4069 independent reflections
Radiation source: Enhance (Mo) X-ray Source	2928 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.031$
Detector resolution: 15.9745 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 28.7^\circ$
$T = 160 \text{ K}$	$\theta_{\text{min}} = 2.6^\circ$
$\omega$ scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2007)	$k = 0 \rightarrow 21$
$T_{\text{min}} = 0.91$ , $T_{\text{max}} = 0.99$	$l = 0 \rightarrow 23$
18979 measured reflections	

## *Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Hydrogen site location: geom+difmap
$R[F^2 > 2\sigma(F^2)] = 0.040$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.088$	Method = Modified Sheldrick $w = 1/[\sigma^2(F^2) + (0.03P)^2 + 0.5P]$ , where $P = [\max(F_o^2, 0) + 2F_c^2]/3$
$S = 0.95$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4069 reflections	$\Delta\rho_{\text{max}} = 0.38 \text{ e \AA}^{-3}$
226 parameters	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$
3 restraints	Extinction correction: none

## *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3604 (2)	0.32253 (7)	0.41898 (7)	0.0261
C2	0.1718 (2)	0.36489 (8)	0.38294 (7)	0.0323
C3	0.1256 (2)	0.36854 (9)	0.30423 (8)	0.0391
C4	0.2687 (2)	0.33058 (9)	0.25996 (7)	0.0392
C5	0.4591 (2)	0.28990 (8)	0.29445 (7)	0.0344
C6	0.5061 (2)	0.28359 (7)	0.37441 (7)	0.0279
C7	0.6996 (2)	0.23388 (8)	0.40716 (7)	0.0304
C8	0.7115 (2)	0.18488 (8)	0.46816 (7)	0.0301
C9	0.5383 (2)	0.17092 (8)	0.51782 (6)	0.0269
C10	0.5251 (2)	0.09343 (8)	0.55037 (7)	0.0341
C11	0.3603 (2)	0.07491 (9)	0.59538 (8)	0.0398
C12	0.2069 (2)	0.13407 (9)	0.61005 (8)	0.0389
C13	0.2205 (2)	0.21186 (8)	0.58012 (7)	0.0321
C14	0.3844 (2)	0.23123 (7)	0.53434 (6)	0.0258
C15	0.4034 (2)	0.31760 (7)	0.50530 (6)	0.0257
C16	0.6242 (2)	0.35710 (7)	0.54306 (7)	0.0276
C17	0.3158 (3)	0.57421 (11)	0.16558 (9)	0.0568
C18	0.5184 (3)	0.60108 (10)	0.22080 (8)	0.0455
C19	0.5957 (2)	0.53835 (9)	0.28103 (8)	0.0371
C20	0.8040 (2)	0.56040 (8)	0.33497 (7)	0.0295
O1	0.70946 (16)	0.33413 (6)	0.60746 (5)	0.0369
O2	0.84351 (15)	0.53413 (6)	0.39986 (5)	0.0346
N1	0.7091 (2)	0.41860 (7)	0.50777 (6)	0.0335
O3	0.94149 (17)	0.61035 (6)	0.30567 (5)	0.0435
H151	0.2828	0.3488	0.5249	0.0293*
H21	0.0713	0.3916	0.4145	0.0377*
H81	0.8456	0.1498	0.4786	0.0333*
H71	0.8293	0.2332	0.3790	0.0372*
H191	0.4766	0.5266	0.3115	0.0486*

## supplementary materials

---

H192	0.6295	0.4889	0.2564	0.0473*
H131	0.1155	0.2536	0.5914	0.0383*
H101	0.6291	0.0526	0.5389	0.0392*
H41	0.2338	0.3317	0.2057	0.0465*
H121	0.0935	0.1222	0.6419	0.0464*
H51	0.5637	0.2636	0.2641	0.0395*
H111	0.3507	0.0214	0.6150	0.0470*
H182	0.4770	0.6515	0.2459	0.0609*
H181	0.6503	0.6120	0.1946	0.0611*
H31	-0.0099	0.3971	0.2794	0.0463*
H172	0.2712	0.6162	0.1279	0.0868*
H173	0.1858	0.5625	0.1933	0.0867*
H171	0.3504	0.5240	0.1402	0.0875*
H12	0.834 (2)	0.4432 (9)	0.5320 (8)	0.0446*
H11	0.652 (2)	0.4352 (9)	0.4630 (8)	0.0433*
H311	1.058 (3)	0.6243 (11)	0.3395 (9)	0.0689*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0283 (6)	0.0234 (6)	0.0265 (6)	-0.0013 (5)	0.0032 (5)	-0.0008 (5)
C2	0.0315 (7)	0.0305 (7)	0.0348 (7)	0.0030 (5)	0.0040 (5)	0.0027 (5)
C3	0.0366 (8)	0.0396 (8)	0.0383 (7)	0.0032 (6)	-0.0041 (6)	0.0075 (6)
C4	0.0476 (8)	0.0422 (8)	0.0259 (6)	-0.0032 (7)	-0.0010 (6)	0.0030 (6)
C5	0.0414 (8)	0.0345 (8)	0.0279 (6)	-0.0020 (6)	0.0075 (6)	-0.0025 (5)
C6	0.0297 (6)	0.0258 (7)	0.0279 (6)	-0.0019 (5)	0.0031 (5)	-0.0021 (5)
C7	0.0287 (6)	0.0318 (7)	0.0315 (6)	0.0018 (5)	0.0065 (5)	-0.0068 (5)
C8	0.0270 (6)	0.0295 (7)	0.0321 (6)	0.0056 (5)	-0.0016 (5)	-0.0067 (5)
C9	0.0273 (6)	0.0284 (7)	0.0227 (6)	-0.0004 (5)	-0.0051 (5)	-0.0026 (5)
C10	0.0388 (7)	0.0291 (7)	0.0313 (6)	0.0021 (6)	-0.0057 (6)	-0.0015 (5)
C11	0.0507 (9)	0.0310 (8)	0.0347 (7)	-0.0078 (6)	-0.0051 (6)	0.0073 (6)
C12	0.0384 (8)	0.0448 (9)	0.0327 (7)	-0.0111 (7)	0.0021 (6)	0.0063 (6)
C13	0.0298 (7)	0.0379 (8)	0.0275 (6)	-0.0022 (6)	0.0010 (5)	-0.0008 (5)
C14	0.0264 (6)	0.0277 (7)	0.0215 (5)	-0.0016 (5)	-0.0027 (5)	-0.0024 (5)
C15	0.0270 (6)	0.0256 (7)	0.0252 (6)	0.0034 (5)	0.0055 (5)	-0.0028 (5)
C16	0.0334 (7)	0.0245 (6)	0.0252 (6)	0.0009 (5)	0.0059 (5)	-0.0041 (5)
C17	0.0518 (10)	0.0666 (12)	0.0470 (9)	0.0033 (8)	-0.0109 (7)	-0.0040 (8)
C18	0.0467 (9)	0.0430 (9)	0.0435 (8)	-0.0033 (7)	-0.0052 (7)	0.0042 (7)
C19	0.0398 (8)	0.0347 (8)	0.0361 (7)	-0.0062 (6)	0.0026 (6)	-0.0017 (6)
C20	0.0366 (7)	0.0246 (7)	0.0279 (6)	-0.0004 (5)	0.0066 (5)	-0.0014 (5)
O2	0.0421 (5)	0.0326 (5)	0.0290 (5)	-0.0029 (4)	0.0045 (4)	0.0035 (4)
N1	0.0402 (7)	0.0304 (6)	0.0291 (5)	-0.0056 (5)	0.0024 (5)	0.0018 (5)
O3	0.0457 (6)	0.0529 (7)	0.0297 (5)	-0.0205 (5)	-0.0027 (4)	0.0077 (4)
O1	0.0441 (5)	0.0373 (5)	0.0271 (4)	-0.0124 (4)	-0.0027 (4)	0.0021 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—C2	1.3924 (17)	C12—H121	0.956
C1—C6	1.4018 (17)	C13—C14	1.3860 (17)

C1—C15	1.5133 (15)	C13—H131	0.965
C2—C3	1.3811 (18)	C14—C15	1.5129 (17)
C2—H21	0.975	C15—C16	1.5282 (17)
C3—C4	1.380 (2)	C15—H151	0.981
C3—H31	0.980	C16—N1	1.3204 (16)
C4—C5	1.378 (2)	C16—O1	1.2376 (14)
C4—H41	0.952	C17—C18	1.504 (2)
C5—C6	1.4054 (17)	C17—H172	0.968
C5—H51	0.976	C17—H173	0.988
C6—C7	1.4591 (17)	C17—H171	0.971
C7—C8	1.3373 (18)	C18—C19	1.5034 (19)
C7—H71	0.974	C18—H182	0.984
C8—C9	1.4601 (18)	C18—H181	0.980
C8—H81	0.977	C19—C20	1.4956 (18)
C9—C10	1.3988 (18)	C19—H191	0.967
C9—C14	1.4026 (17)	C19—H192	0.954
C10—C11	1.3778 (19)	C20—O2	1.2165 (14)
C10—H101	0.950	C20—O3	1.3114 (15)
C11—C12	1.378 (2)	N1—H12	0.899 (14)
C11—H111	0.946	N1—H11	0.860 (14)
C12—C13	1.3848 (19)	O3—H311	0.880 (14)
C2—C1—C6	119.25 (11)	C14—C13—H131	119.0
C2—C1—C15	119.95 (11)	C9—C14—C13	119.42 (12)
C6—C1—C15	120.79 (10)	C9—C14—C15	120.31 (11)
C1—C2—C3	121.06 (12)	C13—C14—C15	120.22 (11)
C1—C2—H21	118.6	C1—C15—C14	112.44 (10)
C3—C2—H21	120.4	C1—C15—C16	115.54 (10)
C2—C3—C4	120.03 (13)	C14—C15—C16	110.29 (10)
C2—C3—H31	120.5	C1—C15—H151	107.5
C4—C3—H31	119.5	C14—C15—H151	105.8
C3—C4—C5	119.87 (12)	C16—C15—H151	104.4
C3—C4—H41	119.9	C15—C16—N1	118.47 (11)
C5—C4—H41	120.2	C15—C16—O1	119.24 (11)
C4—C5—C6	121.01 (12)	N1—C16—O1	122.14 (12)
C4—C5—H51	121.0	C18—C17—H172	110.8
C6—C5—H51	118.0	C18—C17—H173	110.1
C5—C6—C1	118.72 (11)	H172—C17—H173	108.7
C5—C6—C7	118.30 (11)	C18—C17—H171	110.2
C1—C6—C7	122.92 (11)	H172—C17—H171	109.7
C6—C7—C8	127.14 (12)	H173—C17—H171	107.2
C6—C7—H71	116.0	C17—C18—C19	113.39 (13)
C8—C7—H71	116.7	C17—C18—H182	108.1
C7—C8—C9	128.04 (12)	C19—C18—H182	108.9
C7—C8—H81	117.0	C17—C18—H181	111.4
C9—C8—H81	114.7	C19—C18—H181	105.9
C8—C9—C10	118.27 (11)	H182—C18—H181	109.0
C8—C9—C14	123.29 (11)	C18—C19—C20	115.46 (11)
C10—C9—C14	118.45 (11)	C18—C19—H191	110.9
C9—C10—C11	121.46 (13)	C20—C19—H191	107.3

## supplementary materials

---


C9—C10—H101	118.2	C18—C19—H192	108.6
C11—C10—H101	120.2	C20—C19—H192	106.7
C10—C11—C12	119.67 (13)	H191—C19—H192	107.6
C10—C11—H111	120.0	C19—C20—O2	123.28 (12)
C12—C11—H111	120.3	C19—C20—O3	113.77 (11)
C11—C12—C13	119.87 (13)	O2—C20—O3	122.95 (12)
C11—C12—H121	120.4	C16—N1—H12	117.5 (10)
C13—C12—H121	119.7	C16—N1—H11	123.2 (10)
C12—C13—C14	121.09 (13)	H12—N1—H11	119.3 (14)
C12—C13—H131	119.9	C20—O3—H311	111.5 (12)
C6—C1—C2—C3	0.48 (19)	C14—C9—C10—C11	2.43 (18)
C15—C1—C2—C3	-178.34 (12)	C8—C9—C14—C13	177.89 (11)
C2—C1—C6—C5	1.17 (17)	C8—C9—C14—C15	-4.73 (17)
C2—C1—C6—C7	-176.25 (11)	C10—C9—C14—C13	-1.96 (17)
C15—C1—C6—C5	180.00 (12)	C10—C9—C14—C15	175.42 (10)
C15—C1—C6—C7	2.57 (18)	C9—C10—C11—C12	-1.2 (2)
C2—C1—C15—C14	114.81 (12)	C10—C11—C12—C13	-0.6 (2)
C2—C1—C15—C16	-117.37 (12)	C11—C12—C13—C14	1.0 (2)
C6—C1—C15—C14	-63.99 (14)	C12—C13—C14—C9	0.28 (18)
C6—C1—C15—C16	63.83 (14)	C12—C13—C14—C15	-177.10 (11)
C1—C2—C3—C4	-0.7 (2)	C9—C14—C15—C1	64.61 (14)
C2—C3—C4—C5	-0.8 (2)	C9—C14—C15—C16	-65.93 (13)
C3—C4—C5—C6	2.5 (2)	C13—C14—C15—C1	-118.03 (12)
C4—C5—C6—C1	-2.67 (18)	C13—C14—C15—C16	111.43 (12)
C4—C5—C6—C7	174.87 (12)	C1—C15—C16—O1	-156.56 (11)
C1—C6—C7—C8	36.0 (2)	C1—C15—C16—N1	27.97 (15)
C5—C6—C7—C8	-141.46 (14)	C14—C15—C16—O1	-27.68 (15)
C6—C7—C8—C9	-1.9 (2)	C14—C15—C16—N1	156.85 (11)
C7—C8—C9—C10	147.22 (13)	C17—C18—C19—C20	177.01 (13)
C7—C8—C9—C14	-32.6 (2)	C18—C19—C20—O2	152.97 (13)
C8—C9—C10—C11	-177.43 (12)	C18—C19—C20—O3	-27.50 (17)

### *Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H11···O2	0.860 (14)	2.348 (14)	2.8761 (15)	120.0 (10)
N1—H12···O2 <sup>i</sup>	0.898 (13)	2.146 (13)	3.0167 (15)	163.2 (13)
O3—H311···O1 <sup>i</sup>	0.879 (17)	1.698 (17)	2.5658 (13)	168.8 (16)

Symmetry codes: (i)  $-x+2, -y+1, -z+1$ .

Fig. 1



## supplementary materials

---

Fig. 2

