

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-[5-(4-Fluorophenyl)-1,3,4-thiadiazol-2-yl]-2-(4-methoxyphenyl)-1,3-thiazolidin-4-one

Li-He Yin, Rong Wan,* Feng Han, Bin Wang and Jin-Tang Wang

Department of Applied Chemistry, College of Science, Nanjing University of Technology, No. 5 Xinmofan Road, Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: rwan@njut.edu.cn


Received 22 June 2008; accepted 24 June 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.071; wR factor = 0.186; data-to-parameter ratio = 13.6.

The title compound, $\text{C}_{18}\text{H}_{14}\text{FN}_3\text{O}_2\text{S}_2$, was synthesized by the reaction of 5-(4-fluorophenyl)-*N*-(4-methoxybenzylidene)-1,3,4-thiadiazol-2-amine and mercaptoacetic acid. The thiazolidinone ring adopts a twist conformation. The 4-methoxyphenyl ring is almost perpendicular to the thiadiazole ring, making a dihedral angle of $88.4(3)^\circ$. The 4-fluorophenyl ring is nearly coplanar with the thiadiazole ring, the dihedral angle being $6.8(3)^\circ$. The crystal structure involves $\text{C}-\text{H}\cdots\text{N}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds.

Related literature

For related literature, see: Arun *et al.* (1999); Chen *et al.* (2000); Kidwai *et al.* (2000); Vicentini *et al.* (1998); Wasfy *et al.* (1996); Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{18}\text{H}_{14}\text{FN}_3\text{O}_2\text{S}_2$ $M_r = 387.44$ Triclinic, $P\bar{1}$ $a = 6.4550(13)$ Å $b = 8.9200(18)$ Å $c = 16.483(3)$ Å $\alpha = 75.78(3)^\circ$ $\beta = 82.44(3)^\circ$ $\gamma = 71.11(3)^\circ$ $V = 869.0(3)$ Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 0.34$ mm⁻¹ $T = 298(2)$ K $0.10 \times 0.05 \times 0.05$ mm

Data collection

Enraf-Nonius CAD-4 diffractometer

Absorption correction: ψ scan (North *et al.*, 1968) $T_{\min} = 0.967$, $T_{\max} = 0.984$

3421 measured reflections

3120 independent reflections

2054 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.023$

3 standard reflections

every 200 reflections

intensity decay: none

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.070$ $wR(F^2) = 0.186$ $S = 1.00$

3120 reflections

229 parameters

48 restraints

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.41$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6A}\cdots\text{N1}$	0.93	2.60	2.917 (9)	101
$\text{C8}-\text{H8A}\cdots\text{O2}^i$	0.98	2.52	3.233 (7)	129
$\text{C14}-\text{H14A}\cdots\text{S2}$	0.93	2.74	3.146 (6)	107
$\text{C18}-\text{H18A}\cdots\text{N3}$	0.93	2.57	2.881 (7)	100

Symmetry code: (i) $x - 1, y, z$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor Hua-Qin Wang of Nanjing University for carrying out the X-ray crystallographic analysis.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2579).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Arun, K. P., Nag, V. L. & Panda, C. S. (1999). *Indian J. Chem. Sect. B*, **38**, 998–1001.
- Chen, H. S., Li, Z. M. & Han, Y. F. (2000). *J. Agric. Food Chem.* **48**, 5312–5315.
- Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Kidwai, M., Negi, N. & Misra, P. (2000). *J. Indian Chem. Soc.* **77**, 46–48.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vicentini, C. B., Manfrini, M., Veronese, A. C. & Guarnieri, M. (1998). *J. Heterocycl. Chem.* **35**, 29–36.
- Wasfy, A. A., Nassar, S. A. & Eissa, A. M. (1996). *Indian J. Chem. Sect. B*, **35**, 1218–1220.

supplementary materials

Acta Cryst. (2008). E64, o1359 [doi:10.1107/S1600536808019089]

3-[5-(4-Fluorophenyl)-1,3,4-thiadiazol-2-yl]-2-(4-methoxyphenyl)-1,3-thiazolidin-4-one

L.-H. Yin, R. Wan, F. Han, B. Wang and J.-T. Wang

Comment

1,3,4-Thiadiazole derivatives containing the thiazolidinone unit are of great interest because of their chemical and pharmaceutical properties. Some derivatives have fungicidal activities and exhibit certain herbicidal activities (Chen *et al.*, 2000; Kidwai *et al.*, 2000; Vicentini *et al.*, 1998). Some show insecticidal activities (Arun *et al.*, 1999; Wasfy *et al.*, 1996).

We report here the crystal structure of the titled compound, (I). The molecular structure of (I) is shown in Fig.1. In this structure, the thiazolidinone adopts a twist conformation, the dihedral angle between the C9/S1/C10 and C9/N1/C10 is 15.5 (7)°. The thiadiazole ring is an aromatic heterocyclic ring, all atoms are in the same plane. The angle between the thiadiazole ring and the 4-fluorophenyl ring is 6.8 (3)°. The 4-methoxyphenyl ring is nearly perpendicular to the thiadiazole ring, with the dihedral angle being 88.4 (3)°. There are intramolecular C—H···S and C—H···N hydrogen bonding interactions in the molecule structure. In the crystal structure, intermolecular C—H···O hydrogen bonding interactions link the molecules (Table 1 and Fig. 2).

Experimental

5-(4-Fluorophenyl)-*N*-(4-methoxybenzylidene)-1,3,4-thiadiazol-2-amine (5 mmol) and mercapto-acetic acid (5 mmol) were added in toluene (50 ml). The water was removed by distillation for 5 h. The reaction mixture was left to cool to room temperature, filtered, and the filter cake was crystallized from acetone to give pure compound (I) [m.p. 341–345 K]. Crystals of (I) suitable for X-ray diffraction were obtained by slow evaporation of an acetone solution.

Refinement

All H atoms were positioned geometrically, with C—H = 0.98, 0.97, 0.96 and 0.93 Å for methine, methylene, methyl and aromatic H atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H atoms and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. A view of the molecular structure of (I). Displacement ellipsoids are drawn at the 50% probability level. Dashed lines indicate intramolecular C—H···S and C—H···N hydrogen bonding interactions.




Fig. 2. A packing diagram for (I). Dashed lines indicate intramolecular C—H...S and C—H...N, and intermolecular C—H...O hydrogen bonding interactions.

3-[5-(4-Fluorophenyl)-1,3,4-thiadiazol-2-yl]-2-(4-methoxyphenyl)-1,3-thiazolidin-4-one

Crystal data

$C_{18}H_{14}FN_3O_2S_2$	$Z = 2$
$M_r = 387.44$	$F_{000} = 400$
Triclinic, $P\bar{1}$	$D_x = 1.481 \text{ Mg m}^{-3}$
Hall symbol: $-P\ 1$	Melting point = 341–345 K
$a = 6.4550 (13) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.9200 (18) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$c = 16.483 (3) \text{ \AA}$	Cell parameters from 25 reflections
$\alpha = 75.78 (3)^\circ$	$\theta = 9\text{--}12^\circ$
$\beta = 82.44 (3)^\circ$	$\mu = 0.34 \text{ mm}^{-1}$
$\gamma = 71.11 (3)^\circ$	$T = 298 (2) \text{ K}$
$V = 869.0 (3) \text{ \AA}^3$	Block, colourless
	$0.10 \times 0.05 \times 0.05 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.023$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.2^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 1.3^\circ$
$T = 298(2) \text{ K}$	$h = -7 \rightarrow 7$
$\omega/2\theta$ scans	$k = -10 \rightarrow 10$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = 0 \rightarrow 19$
$T_{\text{min}} = 0.967$, $T_{\text{max}} = 0.984$	3 standard reflections
3421 measured reflections	every 200 reflections
3120 independent reflections	intensity decay: none
2054 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.070$$

$$wR(F^2) = 0.186$$

$$S = 1.00$$

3120 reflections

229 parameters

48 restraints

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.05P)^2 + 2.5P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.1232 (3)	1.32787 (19)	0.18123 (10)	0.0698 (5)
S2	0.7085 (2)	0.94841 (16)	0.40097 (8)	0.0525 (4)
F	1.1681 (6)	0.3585 (4)	0.7202 (2)	0.0868 (11)
O1	0.3391 (7)	0.6926 (6)	0.0237 (3)	0.0873 (14)
O2	0.6785 (6)	1.2181 (5)	0.2766 (2)	0.0669 (11)
N1	0.4009 (7)	1.1081 (5)	0.2832 (3)	0.0524 (10)
N2	0.3572 (7)	0.8821 (6)	0.3839 (3)	0.0624 (12)
N3	0.4550 (7)	0.7685 (6)	0.4514 (3)	0.0615 (12)
C1	0.1565 (11)	0.6619 (9)	-0.0019 (4)	0.084 (2)
H1A	0.2071	0.5921	-0.0410	0.126*
H1B	0.0557	0.7626	-0.0281	0.126*
H1C	0.0839	0.6105	0.0462	0.126*
C2	0.2971 (9)	0.7920 (7)	0.0792 (4)	0.0628 (14)
C3	0.0998 (10)	0.8504 (8)	0.1184 (4)	0.0731 (16)
H3B	-0.0205	0.8231	0.1084	0.088*
C4	0.0759 (9)	0.9502 (8)	0.1733 (4)	0.0687 (16)
H4A	-0.0604	0.9869	0.2006	0.082*
C5	0.2469 (8)	0.9971 (7)	0.1889 (3)	0.0550 (12)
C6	0.4424 (11)	0.9355 (10)	0.1499 (5)	0.107 (3)
H6A	0.5630	0.9621	0.1602	0.128*
C7	0.4704 (12)	0.8350 (10)	0.0956 (5)	0.109
H7A	0.6079	0.7959	0.0697	0.131*
C8	0.2051 (8)	1.1190 (6)	0.2422 (3)	0.0544 (13)

supplementary materials

H8A	0.0890	1.1042	0.2852	0.065*
C9	0.3907 (10)	1.3489 (8)	0.1783 (4)	0.0770 (18)
H9A	0.4707	1.3304	0.1258	0.092*
H9B	0.3797	1.4573	0.1835	0.092*
C10	0.5063 (9)	1.2245 (7)	0.2508 (3)	0.0567 (13)
C11	0.4717 (8)	0.9827 (6)	0.3532 (3)	0.0505 (12)
C12	0.6418 (8)	0.7842 (6)	0.4677 (3)	0.0470 (11)
C13	0.7762 (8)	0.6774 (6)	0.5357 (3)	0.0480 (11)
C14	0.9628 (9)	0.7058 (7)	0.5547 (3)	0.0608 (14)
H14A	1.0009	0.7966	0.5243	0.073*
C15	1.0908 (10)	0.6009 (7)	0.6181 (4)	0.0668 (15)
H15A	1.2114	0.6223	0.6324	0.080*
C16	1.0368 (9)	0.4640 (7)	0.6596 (3)	0.0613 (15)
C17	0.8542 (10)	0.4310 (7)	0.6425 (3)	0.0647 (15)
H17A	0.8189	0.3390	0.6728	0.078*
C18	0.7262 (9)	0.5372 (6)	0.5799 (3)	0.0584 (14)
H18A	0.6044	0.5156	0.5668	0.070*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0724 (10)	0.0617 (9)	0.0743 (10)	-0.0157 (8)	-0.0163 (8)	-0.0123 (8)
S2	0.0476 (7)	0.0587 (8)	0.0565 (8)	-0.0219 (6)	-0.0011 (6)	-0.0152 (6)
F	0.092 (3)	0.081 (2)	0.075 (2)	-0.008 (2)	-0.022 (2)	-0.0116 (19)
O1	0.085 (3)	0.102 (3)	0.098 (3)	-0.038 (3)	0.013 (3)	-0.059 (3)
O2	0.054 (2)	0.069 (3)	0.081 (3)	-0.0299 (19)	-0.005 (2)	-0.008 (2)
N1	0.052 (2)	0.056 (3)	0.056 (3)	-0.023 (2)	0.000 (2)	-0.017 (2)
N2	0.057 (3)	0.064 (3)	0.069 (3)	-0.029 (2)	-0.007 (2)	-0.003 (2)
N3	0.058 (3)	0.068 (3)	0.063 (3)	-0.028 (2)	-0.007 (2)	-0.008 (2)
C1	0.100 (5)	0.094 (5)	0.075 (4)	-0.038 (4)	-0.012 (4)	-0.032 (4)
C2	0.064 (3)	0.061 (3)	0.067 (3)	-0.015 (3)	-0.002 (3)	-0.027 (3)
C3	0.064 (3)	0.089 (4)	0.077 (4)	-0.026 (3)	0.003 (3)	-0.038 (3)
C4	0.051 (3)	0.090 (4)	0.072 (4)	-0.024 (3)	0.011 (3)	-0.035 (3)
C5	0.050 (3)	0.062 (3)	0.054 (3)	-0.016 (2)	-0.004 (2)	-0.014 (2)
C6	0.062 (4)	0.139 (6)	0.152 (6)	-0.034 (4)	0.011 (4)	-0.096 (5)
C7	0.070	0.148	0.146	-0.040	0.021	-0.105
C8	0.046 (3)	0.059 (3)	0.061 (3)	-0.018 (2)	-0.003 (2)	-0.016 (3)
C9	0.083 (4)	0.081 (4)	0.065 (4)	-0.037 (4)	0.005 (3)	-0.001 (3)
C10	0.055 (3)	0.061 (3)	0.058 (3)	-0.024 (3)	0.008 (3)	-0.017 (3)
C11	0.049 (3)	0.057 (3)	0.051 (3)	-0.021 (2)	0.004 (2)	-0.018 (2)
C12	0.046 (3)	0.047 (3)	0.047 (3)	-0.013 (2)	0.005 (2)	-0.014 (2)
C13	0.042 (3)	0.050 (3)	0.048 (3)	-0.007 (2)	0.006 (2)	-0.016 (2)
C14	0.060 (3)	0.072 (4)	0.057 (3)	-0.028 (3)	0.001 (3)	-0.016 (3)
C15	0.065 (4)	0.075 (4)	0.061 (4)	-0.014 (3)	-0.015 (3)	-0.019 (3)
C16	0.065 (4)	0.059 (3)	0.045 (3)	0.003 (3)	-0.012 (3)	-0.008 (3)
C17	0.074 (4)	0.058 (3)	0.058 (3)	-0.018 (3)	-0.004 (3)	-0.005 (3)
C18	0.060 (3)	0.059 (3)	0.063 (3)	-0.027 (3)	0.000 (3)	-0.017 (3)

Geometric parameters (Å, °)

S1—C9	1.790 (6)	C4—H4A	0.9300
S1—C8	1.828 (5)	C5—C6	1.349 (8)
S2—C11	1.716 (5)	C5—C8	1.497 (7)
S2—C12	1.741 (5)	C6—C7	1.373 (9)
F—C16	1.359 (6)	C6—H6A	0.9300
O1—C2	1.368 (6)	C7—H7A	0.9300
O1—C1	1.428 (7)	C8—H8A	0.9800
O2—C10	1.221 (6)	C9—C10	1.504 (8)
N1—C10	1.384 (6)	C9—H9A	0.9700
N1—C11	1.402 (6)	C9—H9B	0.9700
N1—C8	1.475 (6)	C12—C13	1.454 (7)
N2—C11	1.309 (6)	C13—C14	1.396 (7)
N2—N3	1.371 (6)	C13—C18	1.396 (7)
N3—C12	1.326 (6)	C14—C15	1.377 (8)
C1—H1A	0.9600	C14—H14A	0.9300
C1—H1B	0.9600	C15—C16	1.373 (8)
C1—H1C	0.9600	C15—H15A	0.9300
C2—C3	1.354 (8)	C16—C17	1.381 (8)
C2—C7	1.371 (8)	C17—C18	1.372 (7)
C3—C4	1.380 (8)	C17—H17A	0.9300
C3—H3B	0.9300	C18—H18A	0.9300
C4—C5	1.373 (7)		
C9—S1—C8	93.5 (3)	C5—C8—H8A	109.0
C11—S2—C12	85.8 (2)	S1—C8—H8A	109.0
C2—O1—C1	117.6 (5)	C10—C9—S1	106.9 (4)
C10—N1—C11	122.7 (4)	C10—C9—H9A	110.3
C10—N1—C8	118.0 (4)	S1—C9—H9A	110.3
C11—N1—C8	119.3 (4)	C10—C9—H9B	110.3
C11—N2—N3	110.4 (4)	S1—C9—H9B	110.3
C12—N3—N2	113.4 (4)	H9A—C9—H9B	108.6
O1—C1—H1A	109.5	O2—C10—N1	122.5 (5)
O1—C1—H1B	109.5	O2—C10—C9	125.7 (5)
H1A—C1—H1B	109.5	N1—C10—C9	111.7 (5)
O1—C1—H1C	109.5	N2—C11—N1	119.7 (4)
H1A—C1—H1C	109.5	N2—C11—S2	116.8 (4)
H1B—C1—H1C	109.5	N1—C11—S2	123.4 (4)
C3—C2—O1	125.2 (5)	N3—C12—C13	123.5 (5)
C3—C2—C7	118.1 (6)	N3—C12—S2	113.5 (4)
O1—C2—C7	116.7 (5)	C13—C12—S2	123.0 (4)
C2—C3—C4	120.4 (6)	C14—C13—C18	118.8 (5)
C2—C3—H3B	119.8	C14—C13—C12	121.6 (5)
C4—C3—H3B	119.8	C18—C13—C12	119.5 (5)
C5—C4—C3	122.1 (5)	C15—C14—C13	120.7 (5)
C5—C4—H4A	118.9	C15—C14—H14A	119.7
C3—C4—H4A	118.9	C13—C14—H14A	119.7
C6—C5—C4	116.3 (5)	C16—C15—C14	118.6 (5)

supplementary materials

C6—C5—C8	124.1 (5)	C16—C15—H15A	120.7
C4—C5—C8	119.5 (5)	C14—C15—H15A	120.7
C5—C6—C7	122.5 (6)	F—C16—C15	118.2 (5)
C5—C6—H6A	118.7	F—C16—C17	119.3 (6)
C7—C6—H6A	118.7	C15—C16—C17	122.5 (5)
C2—C7—C6	120.5 (6)	C18—C17—C16	118.5 (5)
C2—C7—H7A	119.7	C18—C17—H17A	120.8
C6—C7—H7A	119.7	C16—C17—H17A	120.8
N1—C8—C5	113.5 (4)	C17—C18—C13	120.9 (5)
N1—C8—S1	103.5 (3)	C17—C18—H18A	119.6
C5—C8—S1	112.5 (4)	C13—C18—H18A	119.6
N1—C8—H8A	109.0		
C11—N2—N3—C12	-1.9 (6)	S1—C9—C10—O2	-168.3 (5)
C1—O1—C2—C3	7.2 (9)	S1—C9—C10—N1	15.5 (6)
C1—O1—C2—C7	-172.7 (7)	N3—N2—C11—N1	179.5 (4)
O1—C2—C3—C4	-180.0 (6)	N3—N2—C11—S2	1.2 (6)
C7—C2—C3—C4	0.0 (10)	C10—N1—C11—N2	175.9 (5)
C2—C3—C4—C5	1.5 (10)	C8—N1—C11—N2	-3.5 (7)
C3—C4—C5—C6	-2.3 (10)	C10—N1—C11—S2	-5.9 (7)
C3—C4—C5—C8	173.7 (6)	C8—N1—C11—S2	174.7 (4)
C4—C5—C6—C7	1.8 (12)	C12—S2—C11—N2	-0.2 (4)
C8—C5—C6—C7	-174.0 (7)	C12—S2—C11—N1	-178.4 (4)
C3—C2—C7—C6	-0.5 (12)	N2—N3—C12—C13	-179.5 (4)
O1—C2—C7—C6	179.5 (8)	N2—N3—C12—S2	1.8 (6)
C5—C6—C7—C2	-0.4 (14)	C11—S2—C12—N3	-0.9 (4)
C10—N1—C8—C5	104.3 (5)	C11—S2—C12—C13	-179.7 (4)
C11—N1—C8—C5	-76.2 (6)	N3—C12—C13—C14	-174.9 (5)
C10—N1—C8—S1	-17.9 (5)	S2—C12—C13—C14	3.7 (7)
C11—N1—C8—S1	161.6 (4)	N3—C12—C13—C18	9.1 (7)
C6—C5—C8—N1	-28.9 (9)	S2—C12—C13—C18	-172.3 (4)
C4—C5—C8—N1	155.4 (5)	C18—C13—C14—C15	-2.4 (8)
C6—C5—C8—S1	88.3 (7)	C12—C13—C14—C15	-178.4 (5)
C4—C5—C8—S1	-87.4 (6)	C13—C14—C15—C16	2.8 (8)
C9—S1—C8—N1	22.4 (4)	C14—C15—C16—F	177.9 (5)
C9—S1—C8—C5	-100.6 (4)	C14—C15—C16—C17	-2.6 (9)
C8—S1—C9—C10	-22.2 (5)	F—C16—C17—C18	-178.7 (5)
C11—N1—C10—O2	6.3 (8)	C15—C16—C17—C18	1.9 (9)
C8—N1—C10—O2	-174.3 (5)	C16—C17—C18—C13	-1.4 (8)
C11—N1—C10—C9	-177.4 (5)	C14—C13—C18—C17	1.6 (8)
C8—N1—C10—C9	2.1 (7)	C12—C13—C18—C17	177.7 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6A \cdots N1	0.93	2.60	2.917 (9)	101
C8—H8A \cdots O2 ⁱ	0.98	2.52	3.233 (7)	129
C14—H14A \cdots S2	0.93	2.74	3.146 (6)	107
C18—H18A \cdots N3	0.93	2.57	2.881 (7)	100

Symmetry codes: (i) $x-1, y, z$.

Fig. 1




Fig. 2

