

Hexa- μ_2 -acetato-triaqua- μ_3 -oxido-triiron(III) nitrate acetic acid solvate

Sumei Yao,^a Jianhua Liu^b and Qiuxia Han^{b*}

^aMedical College of Henan University, Henan University, Kaifeng 475004, People's Republic of China, and ^bBasic Experiment Teaching Center, Henan University, Kaifeng 475004, People's Republic of China
Correspondence e-mail: hdhqx@henu.edu.cn


Received 28 May 2008; accepted 29 June 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å;
R factor = 0.044; wR factor = 0.106; data-to-parameter ratio = 13.1.

The asymmetric unit of the title compound, $[\text{Fe}_3(\text{CH}_3\text{COO})_6\cdot(\text{H}_2\text{O})_3]\text{NO}_3\cdot\text{CH}_3\text{COOH}$, consists of a hexa- μ_2 -acetato-triaqua- μ_3 -oxido-triiron(III) macrocation, a nitrate ion and an acetic acid solvent molecule. In the cation, each Fe^{3+} ion is coordinated by four carboxylate O atoms, one central bridged O atom and one water molecule, resulting in distorted FeO_6 octahedra. A network of O—H···O hydrogen bonds helps to establish the packing.

Related literature

For related literature, see: Fujihara *et al.* (1998); Ren *et al.* (2004); Thirumurugan & Natarajan (2004); Vrubel *et al.* (2006); Zhang *et al.* (2005).



Experimental

Crystal data

$[\text{Fe}_3(\text{C}_2\text{H}_3\text{O}_2)_6\text{O}(\text{H}_2\text{O})_3]\text{NO}_3\cdot\text{C}_2\text{H}_4\text{O}_2$

$M_r = 713.92$

Monoclinic, $P2_1/c$

$a = 11.835$ (3) Å

$b = 14.755$ (4) Å

$c = 15.250$ (4) Å

$\beta = 90.851$ (5)°

$V = 2662.8$ (12) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.71$ mm⁻¹

$T = 296$ (2) K

$0.18 \times 0.13 \times 0.10$ mm

Data collection

Bruker SMART CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 2001)

$T_{\min} = 0.750$, $T_{\max} = 0.848$

14072 measured reflections

4953 independent reflections

3355 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.106$

$S = 1.00$

4953 reflections

378 parameters

9 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.51$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.41$ e Å⁻³

Table 1
Selected bond lengths (Å).

Fe1—O13	1.897 (2)	Fe2—O2	2.030 (2)
Fe1—O1	1.987 (2)	Fe2—O4	2.030 (2)
Fe1—O10	1.995 (2)	Fe2—O2W	2.126 (3)
Fe1—O12	2.005 (3)	Fe3—O13	1.916 (2)
Fe1—O3	2.063 (2)	Fe3—O11	2.011 (3)
Fe1—O1W	2.104 (2)	Fe3—O6	2.013 (2)
Fe2—O13	1.900 (2)	Fe3—O8	2.013 (2)
Fe2—O5	1.985 (2)	Fe3—O9	2.017 (2)
Fe2—O7	2.021 (2)	Fe3—O3W	2.048 (2)

Table 2
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O14—H14···O17 ⁱ	0.82	1.82	2.642 (4)	178
O3W—H3AW···O15 ⁱⁱ	0.816 (9)	1.894 (9)	2.697 (4)	168 (2)
O1W—H1AW···O18 ⁱⁱⁱ	0.815 (9)	2.008 (10)	2.821 (4)	176 (2)
O3W—H3BW···O17 ^{iv}	0.818 (9)	1.938 (13)	2.742 (4)	167 (3)
O2W—H2AW···O15 ^v	0.816 (9)	2.28 (2)	2.904 (4)	134 (2)
O1W—H1BW···O3 ^{vi}	0.814 (9)	2.188 (12)	2.948 (3)	155 (2)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x - 1, y, z$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$;
(iv) $x, y + 1, z$; (v) $x - 1, -y + \frac{3}{2}, z - \frac{1}{2}$; (vi) $-x + 1, -y + 2, -z + 1$.

Data collection: *SMART* (Bruker, 2001); cell refinement and data reduction: *SAINT-Plus* (Bruker, 2001); structure solution: *SHELXS97* (Sheldrick, 2008); structure refinement: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *PLATON*.

This work was supported by the Basic Research Foundation for Natural Science of Henan University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2740).

References

- Bruker (2001). *SAINT-Plus* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fujihara, T., Aonahata, J., Kumakura, S., Nagasawa, A., Murakami, K. & Ito, T. (1998). *Inorg. Chem.* **37**, 3779–3784.
- Ren, X. M., Okudera, H. & Kremer, R. K. (2004). *Acta Cryst. E60*, m14–m16.
- Sheldrick, G. M. (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Thirumurugan, A. & Natarajan, S. (2004). *Dalton Trans.* pp. 2923–2928.
- Vrubel, H., Hasegawa, T., Oliverira, E. & Nunes, F. S. (2006). *Inorg. Chem. Commun.* **9**, 208–211.
- Zhang, H. T., Li, Y. Z., Wang, H. Q., Nfor, E. N. & You, X. Z. (2005). *CrysfEngComm*, **7**, 578–585.

supplementary materials

Acta Cryst. (2008). E64, m989 [doi:10.1107/S1600536808019806]

Hexa- μ_2 -acetato-triaqua- μ_3 -oxido-triiron(III) nitrate acetic acid solvate

S. Yao, J. Liu and Q. Han

Comment

Transiton-metal coordination complexes based on carboxylates have been attracting chemist's interests and constitutes one of the widest families of research (Thirumurugan & Natarajan, 2004). During the past years, lots of novel carboxylates compounds have been reported (Zhang *et al.*, 2005), in which carboxlate-supported Cr₃(/m₃-O) (Fujihara *et al.*, 1998) and Fe₃(/m₃-O) core (Ren *et al.*, 2004; Vrubel *et al.*, 2006), present two large kinds of widely investigated transtion-metal complexes. Herein, we report the title compound (I).

The title compound, (I), presents a macrocation of [Fe₃O(CH₃COO)₆(H₂O)₃]⁺, in which Fe³⁺ is coordinated by four oxygen atoms from four carboxylates of four acetate anions, one central bridged oxygen atom, and one water molecule. The environment of all the Fe ions are distorted octahedral geometry (Fig. 1). The three Fe atoms approximatively reside in an equilateral triangle with an oxide ion in the center [Fe₃O]. The Fe—O distances range from 1.897 (2) to 2.126 (3) Å (Table 1).

In the crystal, the components are linked by O—H···O hydrogen bonds generating a three-dimensional framework (Fig. 2 and Table 2).

Experimental

Fe(NO₃)₃.9H₂O (1 mmol, 0.404 g) was suspended in 5 ml water and 3 ml (1 mol/L) NaOH solution was added dropwise to produce a brown precipitate, then 25 ml acetic acid were added to the mixture. It was stirred under reflux for 3 h. The solution was filtered, and the filtrate was kept at the room temperature. After one weeks, xxx blocks of (I) were obtained.

Refinement

H atoms were treated as riding, with C—H distances in the range of 0.93–0.98 Å and O—H distances of 0.82 Å, and were refined as riding with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C}_\text{methylene}$ and $\text{C}_\text{methyldyne})$ and $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{O}$ or $\text{C}_\text{methyl})$.

supplementary materials

Figures




Fig. 1. The molecular structure of (I), with displacement ellipsoids for the non-hydrogen atoms drawn at the 50% probability level.




Fig. 2. Three-dimensional structure of (I), with hydrogen bonds shown as dashed lines.

Hexa- μ_2 -acetato-triaqua- μ_3 -oxido-triiron(III) nitrate acetic acid solvate

Crystal data

$[Fe_3(C_2H_3O_2)_6O(H_2O)_3]NO_3 \cdot C_2H_4O_2$	$F_{000} = 1460$
$M_r = 713.92$	$D_x = 1.781 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.835 (3) \text{ \AA}$	Cell parameters from 2092 reflections
$b = 14.755 (4) \text{ \AA}$	$\theta = 2.2\text{--}23.2^\circ$
$c = 15.250 (4) \text{ \AA}$	$\mu = 1.71 \text{ mm}^{-1}$
$\beta = 90.851 (5)^\circ$	$T = 296 (2) \text{ K}$
$V = 2662.8 (12) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.18 \times 0.13 \times 0.10 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	4953 independent reflections
Radiation source: fine-focus sealed tube	3355 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.054$
$T = 296(2) \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan	$h = -14 \rightarrow 11$

(SADABS; Sheldrick, 2001)
 $T_{\min} = 0.750$, $T_{\max} = 0.848$
 14072 measured reflections

$k = -17 \rightarrow 17$
 $l = -18 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.106$	$w = 1/[\sigma^2(F_o^2) + (0.048P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\max} = 0.001$
4953 reflections	$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$
378 parameters	$\Delta\rho_{\min} = -0.41 \text{ e \AA}^{-3}$
9 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.40305 (4)	0.89656 (3)	0.62790 (3)	0.02692 (12)
Fe2	0.17864 (4)	0.76827 (3)	0.59152 (3)	0.02751 (12)
Fe3	0.16879 (4)	0.93399 (3)	0.73634 (3)	0.02779 (13)
O1W	0.5730 (2)	0.92274 (16)	0.59831 (17)	0.0390 (7)
O2W	0.1067 (2)	0.66185 (17)	0.51505 (18)	0.0536 (8)
O3W	0.0853 (2)	1.01043 (16)	0.82684 (16)	0.0414 (7)
O1	0.4549 (2)	0.76906 (16)	0.61642 (19)	0.0505 (8)
O2	0.30533 (19)	0.67883 (15)	0.61664 (16)	0.0373 (6)
O3	0.3729 (2)	0.90801 (17)	0.49476 (15)	0.0433 (7)
O4	0.2469 (2)	0.79746 (15)	0.47350 (15)	0.0375 (6)
O5	0.0995 (2)	0.71685 (15)	0.69403 (15)	0.0379 (6)
O6	0.0942 (2)	0.82725 (15)	0.79378 (16)	0.0461 (7)
O7	0.0395 (2)	0.84025 (16)	0.55730 (15)	0.0387 (7)

supplementary materials

O8	0.03183 (19)	0.94965 (16)	0.65798 (16)	0.0406 (7)
O9	0.2290 (2)	1.05548 (15)	0.69826 (17)	0.0430 (7)
O10	0.38902 (19)	1.03126 (16)	0.62901 (17)	0.0456 (7)
O11	0.2928 (2)	0.92041 (18)	0.82695 (16)	0.0492 (8)
O12	0.4514 (2)	0.89345 (18)	0.75438 (16)	0.0481 (7)
O13	0.25047 (17)	0.86634 (13)	0.65078 (14)	0.0254 (5)
O17	0.1754 (3)	0.16154 (19)	0.9058 (2)	0.0782 (11)
O18	0.2920 (3)	0.26869 (19)	0.9325 (2)	0.0674 (9)
O14	0.7459 (2)	0.9001 (2)	0.9425 (2)	0.0738 (10)
H14	0.7698	0.8798	0.9893	0.111*
O15	0.9248 (2)	0.9381 (2)	0.92925 (19)	0.0637 (9)
C1	0.4856 (3)	0.6134 (2)	0.6411 (3)	0.0472 (11)
H1A	0.4414	0.5590	0.6445	0.071*
H1B	0.5258	0.6224	0.6956	0.071*
H1C	0.5386	0.6080	0.5943	0.071*
C2	0.4094 (3)	0.6923 (2)	0.6240 (2)	0.0302 (8)
C3	0.3290 (3)	0.8632 (3)	0.3488 (2)	0.0479 (11)
H3A	0.3837	0.9093	0.3361	0.072*
H3B	0.2576	0.8792	0.3225	0.072*
H3C	0.3539	0.8063	0.3256	0.072*
C4	0.3164 (3)	0.8553 (2)	0.4464 (2)	0.0305 (9)
C5	0.0056 (3)	0.6886 (3)	0.8275 (2)	0.0458 (11)
H5A	-0.0075	0.6307	0.8004	0.069*
H5B	-0.0655	0.7167	0.8400	0.069*
H5C	0.0478	0.6803	0.8811	0.069*
C6	0.0711 (3)	0.7479 (2)	0.7666 (2)	0.0317 (9)
C7	-0.1126 (3)	0.9443 (3)	0.5496 (3)	0.0476 (11)
H7A	-0.1380	0.9960	0.5820	0.071*
H7B	-0.1696	0.8980	0.5504	0.071*
H7C	-0.0987	0.9618	0.4901	0.071*
C8	-0.0058 (3)	0.9087 (2)	0.5905 (2)	0.0317 (9)
C9	0.3287 (3)	1.1823 (2)	0.6455 (3)	0.0512 (12)
H9A	0.3975	1.1935	0.6147	0.077*
H9B	0.3322	1.2118	0.7016	0.077*
H9C	0.2659	1.2056	0.6120	0.077*
C10	0.3142 (3)	1.0824 (2)	0.6584 (2)	0.0303 (9)
C11	0.4539 (3)	0.8788 (3)	0.9092 (2)	0.0487 (11)
H11A	0.4025	0.8865	0.9567	0.073*
H11B	0.5168	0.9193	0.9168	0.073*
H11C	0.4807	0.8174	0.9084	0.073*
C12	0.3942 (3)	0.8995 (2)	0.8241 (2)	0.0324 (9)
O16	0.1840 (3)	0.2745 (2)	0.8165 (2)	0.0802 (11)
N1	0.2187 (3)	0.2358 (2)	0.8830 (2)	0.0534 (10)
C13	0.7984 (4)	0.9702 (3)	0.8110 (3)	0.0624 (13)
H13A	0.7694	1.0306	0.8163	0.094*
H13B	0.7415	0.9318	0.7851	0.094*
H13C	0.8637	0.9707	0.7744	0.094*
C14	0.8301 (3)	0.9353 (3)	0.8986 (3)	0.0470 (11)
H3AW	0.0417 (12)	0.9921 (13)	0.8637 (11)	0.083 (16)*

H1AW	0.6123 (14)	0.8779 (8)	0.5921 (16)	0.048 (12)*
H3BW	0.1149 (16)	1.0579 (9)	0.8427 (17)	0.085 (16)*
H2AW	0.0434 (9)	0.6407 (19)	0.5202 (14)	0.106 (19)*
H2BW	0.122 (2)	0.667 (3)	0.4630 (7)	0.16 (3)*
H1BW	0.584 (3)	0.9594 (9)	0.5594 (11)	0.079 (16)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0246 (3)	0.0245 (2)	0.0317 (3)	-0.0017 (2)	0.0026 (2)	-0.0008 (2)
Fe2	0.0289 (3)	0.0224 (2)	0.0313 (3)	-0.0028 (2)	0.0011 (2)	0.0004 (2)
Fe3	0.0276 (3)	0.0232 (2)	0.0327 (3)	-0.0019 (2)	0.0055 (2)	-0.0002 (2)
O1W	0.0293 (13)	0.0336 (13)	0.0542 (16)	0.0006 (11)	0.0079 (12)	0.0048 (12)
O2W	0.0598 (18)	0.0415 (16)	0.0590 (19)	-0.0159 (14)	-0.0129 (15)	-0.0030 (14)
O3W	0.0436 (15)	0.0338 (14)	0.0472 (15)	-0.0071 (12)	0.0163 (13)	-0.0110 (12)
O1	0.0332 (14)	0.0288 (13)	0.090 (2)	0.0000 (12)	0.0072 (14)	-0.0050 (14)
O2	0.0375 (14)	0.0235 (12)	0.0506 (15)	0.0018 (11)	-0.0025 (12)	-0.0014 (11)
O3	0.0430 (15)	0.0563 (16)	0.0306 (13)	-0.0180 (13)	0.0012 (12)	0.0004 (12)
O4	0.0425 (14)	0.0368 (13)	0.0333 (13)	-0.0113 (12)	0.0046 (11)	-0.0004 (11)
O5	0.0445 (15)	0.0337 (13)	0.0356 (14)	-0.0096 (12)	0.0077 (12)	0.0040 (11)
O6	0.0565 (16)	0.0293 (14)	0.0530 (16)	-0.0106 (12)	0.0223 (13)	0.0000 (12)
O7	0.0382 (14)	0.0376 (14)	0.0399 (14)	0.0067 (12)	-0.0069 (12)	-0.0032 (12)
O8	0.0323 (14)	0.0413 (14)	0.0482 (16)	0.0094 (12)	-0.0028 (12)	-0.0082 (12)
O9	0.0410 (15)	0.0247 (13)	0.0638 (17)	-0.0030 (11)	0.0204 (13)	0.0019 (12)
O10	0.0341 (14)	0.0250 (13)	0.0784 (19)	-0.0050 (11)	0.0198 (14)	-0.0003 (13)
O11	0.0411 (15)	0.0685 (18)	0.0376 (15)	0.0119 (14)	-0.0061 (13)	-0.0085 (13)
O12	0.0355 (14)	0.0763 (19)	0.0324 (14)	-0.0086 (14)	-0.0018 (12)	0.0012 (14)
O13	0.0236 (12)	0.0226 (11)	0.0303 (12)	-0.0007 (9)	0.0018 (10)	-0.0004 (10)
O17	0.116 (3)	0.0492 (18)	0.069 (2)	-0.0405 (18)	-0.019 (2)	0.0065 (16)
O18	0.0604 (19)	0.0547 (18)	0.087 (2)	-0.0155 (16)	-0.0065 (18)	0.0070 (17)
O14	0.0485 (18)	0.096 (2)	0.076 (2)	-0.0236 (18)	-0.0037 (16)	0.0325 (19)
O15	0.0425 (17)	0.089 (2)	0.0599 (19)	-0.0098 (16)	0.0070 (15)	0.0113 (17)
C1	0.050 (2)	0.038 (2)	0.053 (2)	0.0169 (19)	0.000 (2)	0.0056 (19)
C2	0.034 (2)	0.0310 (18)	0.0254 (18)	0.0068 (17)	0.0036 (15)	-0.0009 (15)
C3	0.053 (2)	0.060 (3)	0.031 (2)	-0.006 (2)	0.0001 (19)	0.0040 (19)
C4	0.0294 (19)	0.0318 (19)	0.0305 (19)	0.0023 (16)	0.0027 (16)	0.0025 (16)
C5	0.047 (2)	0.047 (2)	0.043 (2)	-0.017 (2)	0.0054 (19)	0.0111 (19)
C6	0.0297 (19)	0.0257 (19)	0.040 (2)	-0.0044 (15)	-0.0030 (16)	0.0085 (15)
C7	0.039 (2)	0.049 (2)	0.054 (3)	0.0014 (19)	-0.008 (2)	0.010 (2)
C8	0.0255 (18)	0.035 (2)	0.035 (2)	-0.0030 (16)	0.0037 (16)	0.0117 (17)
C9	0.046 (2)	0.032 (2)	0.076 (3)	-0.0016 (19)	0.013 (2)	0.010 (2)
C10	0.0274 (19)	0.0260 (18)	0.038 (2)	-0.0042 (15)	0.0007 (16)	0.0009 (15)
C11	0.054 (3)	0.050 (2)	0.041 (2)	-0.005 (2)	-0.008 (2)	0.0116 (19)
C12	0.040 (2)	0.0230 (17)	0.034 (2)	-0.0067 (16)	-0.0057 (17)	0.0055 (16)
O16	0.120 (3)	0.070 (2)	0.0502 (19)	-0.002 (2)	0.002 (2)	0.0049 (17)
N1	0.063 (2)	0.045 (2)	0.053 (2)	-0.0038 (19)	0.0088 (19)	-0.0055 (18)
C13	0.063 (3)	0.066 (3)	0.058 (3)	0.007 (3)	-0.001 (2)	0.006 (2)
C14	0.039 (2)	0.042 (2)	0.061 (3)	0.0000 (19)	0.005 (2)	0.001 (2)

supplementary materials

Geometric parameters (\AA , $^\circ$)

Fe1—O13	1.897 (2)	O12—C12	1.272 (4)
Fe1—O1	1.987 (2)	O17—N1	1.260 (4)
Fe1—O10	1.995 (2)	O18—N1	1.240 (4)
Fe1—O12	2.005 (3)	O14—C14	1.316 (4)
Fe1—O3	2.063 (2)	O14—H14	0.8200
Fe1—O1W	2.104 (2)	O15—C14	1.209 (5)
Fe2—O13	1.900 (2)	C1—C2	1.494 (5)
Fe2—O5	1.985 (2)	C1—H1A	0.9600
Fe2—O7	2.021 (2)	C1—H1B	0.9600
Fe2—O2	2.030 (2)	C1—H1C	0.9600
Fe2—O4	2.030 (2)	C3—C4	1.501 (5)
Fe2—O2W	2.126 (3)	C3—H3A	0.9600
Fe3—O13	1.916 (2)	C3—H3B	0.9600
Fe3—O11	2.011 (3)	C3—H3C	0.9600
Fe3—O6	2.013 (2)	C5—C6	1.500 (5)
Fe3—O8	2.013 (2)	C5—H5A	0.9600
Fe3—O9	2.017 (2)	C5—H5B	0.9600
Fe3—O3W	2.048 (2)	C5—H5C	0.9600
O1W—H1AW	0.815 (9)	C7—C8	1.497 (5)
O1W—H1BW	0.814 (9)	C7—H7A	0.9600
O2W—H2AW	0.816 (9)	C7—H7B	0.9600
O2W—H2BW	0.819 (9)	C7—H7C	0.9600
O3W—H3AW	0.816 (9)	C9—C10	1.497 (5)
O3W—H3BW	0.818 (9)	C9—H9A	0.9600
O1—C2	1.260 (4)	C9—H9B	0.9600
O2—C2	1.251 (4)	C9—H9C	0.9600
O3—C4	1.258 (4)	C11—C12	1.499 (5)
O4—C4	1.260 (4)	C11—H11A	0.9600
O5—C6	1.249 (4)	C11—H11B	0.9600
O6—C6	1.270 (4)	C11—H11C	0.9600
O7—C8	1.254 (4)	O16—N1	1.229 (5)
O8—C8	1.268 (4)	C13—C14	1.475 (6)
O9—C10	1.251 (4)	C13—H13A	0.9600
O10—C10	1.251 (4)	C13—H13B	0.9600
O11—C12	1.241 (4)	C13—H13C	0.9600
O13—Fe1—O1	95.13 (10)	Fe2—O13—Fe3	119.60 (11)
O13—Fe1—O10	98.80 (9)	C14—O14—H14	109.5
O1—Fe1—O10	165.92 (10)	C2—C1—H1A	109.5
O13—Fe1—O12	94.40 (10)	C2—C1—H1B	109.5
O1—Fe1—O12	88.79 (12)	H1A—C1—H1B	109.5
O10—Fe1—O12	92.13 (11)	C2—C1—H1C	109.5
O13—Fe1—O3	92.81 (9)	H1A—C1—H1C	109.5
O1—Fe1—O3	92.29 (11)	H1B—C1—H1C	109.5
O10—Fe1—O3	85.04 (11)	O2—C2—O1	123.8 (3)
O12—Fe1—O3	172.58 (10)	O2—C2—C1	118.9 (3)
O13—Fe1—O1W	176.56 (9)	O1—C2—C1	117.4 (3)

O1—Fe1—O1W	81.82 (10)	C4—C3—H3A	109.5
O10—Fe1—O1W	84.19 (9)	C4—C3—H3B	109.5
O12—Fe1—O1W	87.14 (10)	H3A—C3—H3B	109.5
O3—Fe1—O1W	85.75 (10)	C4—C3—H3C	109.5
O13—Fe2—O5	97.45 (9)	H3A—C3—H3C	109.5
O13—Fe2—O7	94.56 (9)	H3B—C3—H3C	109.5
O5—Fe2—O7	90.68 (10)	O3—C4—O4	124.8 (3)
O13—Fe2—O2	94.67 (9)	O3—C4—C3	118.2 (3)
O5—Fe2—O2	87.64 (10)	O4—C4—C3	117.0 (3)
O7—Fe2—O2	170.76 (10)	C6—C5—H5A	109.5
O13—Fe2—O4	94.54 (9)	C6—C5—H5B	109.5
O5—Fe2—O4	167.95 (10)	H5A—C5—H5B	109.5
O7—Fe2—O4	89.59 (10)	C6—C5—H5C	109.5
O2—Fe2—O4	90.17 (10)	H5A—C5—H5C	109.5
O13—Fe2—O2W	174.84 (10)	H5B—C5—H5C	109.5
O5—Fe2—O2W	87.69 (11)	O5—C6—O6	124.6 (3)
O7—Fe2—O2W	85.90 (10)	O5—C6—C5	118.9 (3)
O2—Fe2—O2W	84.95 (10)	O6—C6—C5	116.5 (3)
O4—Fe2—O2W	80.32 (10)	C8—C7—H7A	109.5
O13—Fe3—O11	92.62 (10)	C8—C7—H7B	109.5
O13—Fe3—O6	96.76 (9)	H7A—C7—H7B	109.5
O11—Fe3—O6	86.74 (11)	C8—C7—H7C	109.5
O13—Fe3—O8	93.76 (10)	H7A—C7—H7C	109.5
O11—Fe3—O8	173.00 (10)	H7B—C7—H7C	109.5
O6—Fe3—O8	89.64 (11)	O7—C8—O8	124.4 (3)
O13—Fe3—O9	94.77 (9)	O7—C8—C7	118.5 (3)
O11—Fe3—O9	91.68 (11)	O8—C8—C7	117.1 (3)
O6—Fe3—O9	168.42 (10)	C10—C9—H9A	109.5
O8—Fe3—O9	90.67 (10)	C10—C9—H9B	109.5
O13—Fe3—O3W	177.83 (10)	H9A—C9—H9B	109.5
O11—Fe3—O3W	86.91 (10)	C10—C9—H9C	109.5
O6—Fe3—O3W	85.33 (10)	H9A—C9—H9C	109.5
O8—Fe3—O3W	86.82 (10)	H9B—C9—H9C	109.5
O9—Fe3—O3W	83.13 (10)	O9—C10—O10	124.3 (3)
Fe1—O1W—H1AW	115.2 (13)	O9—C10—C9	118.1 (3)
Fe1—O1W—H1BW	116 (2)	O10—C10—C9	117.6 (3)
H1AW—O1W—H1BW	111.0 (16)	C12—C11—H11A	109.5
Fe2—O2W—H2AW	126.2 (19)	C12—C11—H11B	109.5
Fe2—O2W—H2BW	112 (3)	H11A—C11—H11B	109.5
H2AW—O2W—H2BW	109.8 (15)	C12—C11—H11C	109.5
Fe3—O3W—H3AW	126.8 (15)	H11A—C11—H11C	109.5
Fe3—O3W—H3BW	117.5 (16)	H11B—C11—H11C	109.5
H3AW—O3W—H3BW	110.7 (16)	O11—C12—O12	125.1 (3)
C2—O1—Fe1	135.3 (2)	O11—C12—C11	117.6 (3)
C2—O2—Fe2	129.6 (2)	O12—C12—C11	117.3 (3)
C4—O3—Fe1	127.6 (2)	O16—N1—O18	122.8 (4)
C4—O4—Fe2	134.9 (2)	O16—N1—O17	119.9 (4)
C6—O5—Fe2	134.0 (2)	O18—N1—O17	117.2 (4)
C6—O6—Fe3	132.3 (2)	C14—C13—H13A	109.5

supplementary materials


C8—O7—Fe2	132.1 (2)	C14—C13—H13B	109.5
C8—O8—Fe3	134.2 (2)	H13A—C13—H13B	109.5
C10—O9—Fe3	135.5 (2)	C14—C13—H13C	109.5
C10—O10—Fe1	131.6 (2)	H13A—C13—H13C	109.5
C12—O11—Fe3	134.2 (2)	H13B—C13—H13C	109.5
C12—O12—Fe1	130.9 (2)	O15—C14—O14	121.4 (4)
Fe1—O13—Fe2	120.78 (11)	O15—C14—C13	124.1 (4)
Fe1—O13—Fe3	119.60 (11)	O14—C14—C13	114.5 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O14—H14 \cdots O17 ⁱ	0.82	1.82	2.642 (4)	178
O3W—H3AW \cdots O15 ⁱⁱ	0.816 (9)	1.894 (9)	2.697 (4)	168 (2)
O1W—H1AW \cdots O18 ⁱⁱⁱ	0.815 (9)	2.008 (10)	2.821 (4)	176 (2)
O3W—H3BW \cdots O17 ^{iv}	0.818 (9)	1.938 (13)	2.742 (4)	167 (3)
O2W—H2AW \cdots O15 ^v	0.816 (9)	2.28 (2)	2.904 (4)	134 (2)
O1W—H1BW \cdots O3 ^{vi}	0.814 (9)	2.188 (12)	2.948 (3)	155 (2)

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x-1, y, z$; (iii) $-x+1, y+1/2, -z+3/2$; (iv) $x, y+1, z$; (v) $x-1, -y+3/2, z-1/2$; (vi) $-x+1, -y+2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

