

catena-Poly[[bis(3-benzoylpyridine- κ N)-zinc(II)]-di- μ -dicyanamido- κ^4 N¹:N⁵]

Fei Yu, Zhong-Shu Li and Bai-Wang Sun*

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: chmsunbw@seu.edu.cn


Received 15 June 2008; accepted 30 June 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.050; wR factor = 0.130; data-to-parameter ratio = 16.2.

The title compound, $[\text{Zn}(\text{C}_2\text{N}_3)_2(\text{C}_{12}\text{H}_9\text{NO})_2]_n$, is a polymeric zinc(II) complex with the metal ion located on an inversion centre. The Zn^{II} ion is six-coordinated by two N atoms of two 3-benzoylpyridine ligands and four N atoms from four dicyanamide ligands, forming a slightly distorted octahedral configuration. In the crystal structure, neighboring Zn atoms are linked together by double dicyanamide bridges to form a polymeric zinc(II) complex.

Related literature

For related literature, see: Armentano *et al.* (2006); Claramunt *et al.* (2000); Manson *et al.* (1998); Miller (2006).



Experimental

Crystal data

$[\text{Zn}(\text{C}_2\text{N}_3)_2(\text{C}_{12}\text{H}_9\text{NO})_2]$
 $M_r = 563.89$
Monoclinic, $P2_1/c$
 $a = 6.463$ (4) Å
 $b = 7.490$ (4) Å
 $c = 26.300$ (15) Å
 $\beta = 98.399$ (16)°

$V = 1259.5$ (13) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.02$ mm⁻¹
 $T = 293$ (2) K
 $0.07 \times 0.04 \times 0.03$ mm

Data collection

Rigaku Scxmini 1K CCD area-detector diffractometer
Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\text{min}} = 0.867$, $T_{\text{max}} = 1.000$
(expected range = 0.841–0.970)

12079 measured reflections
2880 independent reflections
2342 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.129$
 $S = 1.09$
2880 reflections

178 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.60$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.56$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Zn1–N2	2.162 (3)	Zn1–N4	2.172 (3)
Zn1–N3	2.169 (2)		
N2–Zn1–N3	92.27 (11)	N3–Zn1–N4	89.87 (9)
N2–Zn1–N4	90.79 (9)		

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2086).

References

- Armentano, D., De Munno, G., Guerra, F., Julve, M. & Lloret, F. (2006). *Inorg. Chem.* **45**, 4626–4636.
Claramunt, A., Escuer, A., Mautner, F. A., Sanz, N. & Vicente, R. (2000). *J. Chem. Soc. Dalton Trans.* pp. 2627–2630.
Manson, J. L., Lee, D. W., Rheingold, A. L. & Miller, J. S. (1998). *Inorg. Chem.* **37**, 5966–5967.
Miller, J. S. (2006). *Pramana*, **67**, 1–16.
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m993 [doi:10.1107/S1600536808019880]

***catena*-Poly[[bis(3-benzoylpyridine- κ N)zinc(II)]-di- μ -dicyanamido- κ^4 N¹:N⁵]**

F. Yu, Z.-S. Li and B.-W. Sun

Comment

The dicyanamide ligand has frequently been used to bridge polynuclear transition metal complexes in the study of multidimensional molecule-based magnetic materials and other areas. Many such compounds have been reported (Manson *et al.*, 1998; Claramunt *et al.*, 2000; Armentano *et al.*, 2006; Miller, 2006;). Here, we report the structure of the title Zn^{II} compound, (I). The structure of (I) is illustrated in Fig. 1, and bond distances and angles are given in Table 1. The Zn^{II} ion, which lies on the inversion centre, is in an octahedral geometry and is six-coordinated by six N atoms, from four dicyanamide ligands and two 3-benzoylpyridine ligands in a *trans* arrangement. The resulting coordination geometry is very close to that expected for an ideal octahedral complex. In the crystal structure, the Zn^{II} ions are bridged to form a one-dimensional chain by dicyanamide ligands, through single end-to-end coordination, the dicyanamide ligand acts as a bidentate bridging ligand by coordinating to adjacent Zn^{II} centres through its two terminal nitrile N atoms. No significant contacts are observed between adjacent chains in the crystal structure.

Experimental

All chemicals used (reagent grade) were commercially available. 3-benzoylpyridine (18.3 mg, 0.1 mmol) was added slowly with stirring in aqueous solution (5 ml) of Zn(CH₃COO)₂·2H₂O (21.9 mg, 0.1 mmol) and then sodium dicyanamide (17.8 mg, 0.2 mmol) in aqueous solution (5 ml) was added slowly. The resulting colorless solution was continuously stirred for about 30 min at room temperature and then filtered. The filtrate was slowly evaporated at room temperature over several days, and colorless needles crystals suitable for X-ray analysis were obtained.

Refinement

Positional parameters of all H atoms were calculated geometrically.

Figures




Fig. 1. The molecular structure of the title compound with the atom-numbering scheme and all hydrogen atoms. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. One-dimensional structure in the title compound. Displacement ellipsoids are drawn at the 30% probability level and all hydrogen atoms.

catena-Poly[[bis(3-benzoylpyridine-κN)zinc(II)]- di-μ-dicyanamido-κ⁴N¹:N⁵]

Crystal data

$[\text{Zn}(\text{C}_2\text{N}_3)_2(\text{C}_{12}\text{H}_9\text{NO})_2]$

$M_r = 563.89$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 6.463$ (4) Å

$b = 7.490$ (4) Å

$c = 26.300$ (15) Å

$\beta = 98.399$ (16)°

$V = 1259.5$ (13) Å³

$Z = 2$

$F_{000} = 576$

$D_x = 1.487$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2637 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 1.02$ mm⁻¹

$T = 293$ (2) K

Block, colorless

$0.07 \times 0.04 \times 0.03$ mm

Data collection

Rigaku Scxmini 1K CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 8.192 pixels mm⁻¹

$T = 293$ (2) K

thin-slice ω scans

Absorption correction: Multi-scan (CrystalClear; Rigaku, 2005)

$T_{\min} = 0.867$, $T_{\max} = 1.000$

12079 measured reflections

2880 independent reflections

2342 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.045$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 3.1^\circ$

$h = -8 \rightarrow 8$

$k = -9 \rightarrow 9$

$l = -34 \rightarrow 33$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.129$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0563P)^2 + 0.6329P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.09$ $(\Delta/\sigma)_{\max} < 0.001$
 2880 reflections $\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$
 178 parameters $\Delta\rho_{\min} = -0.56 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	-0.5000	0.0000	0.03523 (16)
O1	0.1648 (4)	-0.5269 (5)	-0.20379 (11)	0.1080 (15)
N2	0.7192 (4)	-0.2827 (3)	0.01736 (10)	0.0467 (6)
N3	0.7294 (4)	-0.6966 (3)	0.03187 (9)	0.0441 (6)
N4	0.5973 (4)	-0.5346 (3)	-0.07500 (9)	0.0374 (5)
N5	0.8571 (7)	-0.9980 (3)	0.05024 (17)	0.0911 (15)
C1	0.7821 (4)	-0.8411 (4)	0.03808 (11)	0.0405 (6)
C2	0.7750 (4)	-0.1441 (4)	0.03062 (11)	0.0411 (6)
C4	0.7925 (4)	-0.5898 (4)	-0.07850 (11)	0.0442 (7)
H4A	0.8899	-0.5924	-0.0487	0.053*
C5	0.8557 (5)	-0.6427 (4)	-0.12406 (11)	0.0488 (7)
H5C	0.9914	-0.6831	-0.1247	0.059*
C6	0.7130 (5)	-0.6347 (4)	-0.16892 (11)	0.0465 (7)
H6A	0.7504	-0.6728	-0.2000	0.056*
C7	0.5141 (5)	-0.5690 (4)	-0.16669 (11)	0.0435 (6)
C8	0.4633 (4)	-0.5223 (3)	-0.11864 (11)	0.0395 (6)
H8A	0.3290	-0.4805	-0.1169	0.047*
C9	0.3436 (5)	-0.5503 (5)	-0.21175 (12)	0.0568 (8)
C10	0.3845 (5)	-0.5665 (4)	-0.26610 (11)	0.0491 (7)
C11	0.5699 (6)	-0.5147 (4)	-0.28267 (13)	0.0553 (8)
H11A	0.6808	-0.4735	-0.2591	0.066*
C12	0.2197 (6)	-0.6285 (5)	-0.30199 (13)	0.0635 (9)
H12A	0.0948	-0.6634	-0.2913	0.076*
C13	0.2419 (8)	-0.6382 (5)	-0.35361 (14)	0.0783 (13)
H13A	0.1321	-0.6804	-0.3774	0.094*
C14	0.4252 (8)	-0.5857 (6)	-0.36965 (15)	0.0811 (14)
H14A	0.4386	-0.5915	-0.4043	0.097*

supplementary materials

C15	0.5896 (8)	-0.5245 (5)	-0.33469 (14)	0.0707 (12)
H15A	0.7137	-0.4898	-0.3458	0.085*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0407 (3)	0.0283 (2)	0.0361 (3)	0.00125 (17)	0.00358 (18)	-0.00050 (17)
O1	0.0444 (14)	0.232 (5)	0.0468 (15)	0.0126 (19)	0.0035 (12)	-0.0020 (19)
N2	0.0456 (13)	0.0353 (13)	0.0569 (15)	-0.0031 (10)	0.0002 (11)	-0.0015 (11)
N3	0.0486 (14)	0.0348 (13)	0.0479 (14)	0.0061 (10)	0.0033 (11)	0.0023 (10)
N4	0.0400 (12)	0.0358 (12)	0.0364 (11)	0.0012 (9)	0.0055 (9)	0.0000 (9)
N5	0.097 (3)	0.0326 (15)	0.121 (3)	0.0078 (15)	-0.060 (2)	-0.0090 (15)
C1	0.0391 (14)	0.0354 (15)	0.0438 (15)	0.0003 (11)	-0.0044 (12)	-0.0041 (11)
C2	0.0414 (15)	0.0348 (15)	0.0444 (15)	0.0057 (12)	-0.0024 (12)	0.0006 (12)
C4	0.0394 (14)	0.0520 (18)	0.0407 (15)	0.0017 (13)	0.0047 (12)	0.0034 (13)
C5	0.0421 (15)	0.0588 (19)	0.0459 (16)	0.0102 (14)	0.0078 (13)	0.0037 (14)
C6	0.0481 (16)	0.0505 (17)	0.0432 (15)	0.0007 (13)	0.0142 (13)	-0.0029 (13)
C7	0.0442 (15)	0.0478 (16)	0.0387 (15)	-0.0025 (13)	0.0065 (12)	-0.0015 (12)
C8	0.0390 (14)	0.0392 (15)	0.0402 (14)	0.0013 (11)	0.0052 (11)	0.0001 (11)
C9	0.0495 (18)	0.079 (2)	0.0413 (17)	0.0025 (17)	0.0050 (14)	0.0018 (16)
C10	0.0579 (19)	0.0500 (17)	0.0378 (15)	0.0099 (15)	0.0017 (13)	-0.0014 (13)
C11	0.071 (2)	0.0519 (19)	0.0445 (17)	0.0105 (16)	0.0125 (16)	0.0005 (14)
C12	0.068 (2)	0.066 (2)	0.0512 (19)	0.0146 (18)	-0.0106 (16)	-0.0011 (16)
C13	0.105 (3)	0.072 (3)	0.048 (2)	0.034 (2)	-0.021 (2)	-0.0112 (18)
C14	0.119 (4)	0.081 (3)	0.043 (2)	0.044 (3)	0.011 (2)	0.0005 (19)
C15	0.097 (3)	0.070 (3)	0.049 (2)	0.028 (2)	0.025 (2)	0.0075 (17)

Geometric parameters (\AA , $^\circ$)

Zn1—N2	2.162 (3)	C6—C7	1.385 (4)
Zn1—N2 ⁱ	2.162 (3)	C6—H6A	0.9300
Zn1—N3 ⁱ	2.169 (2)	C7—C8	1.396 (4)
Zn1—N3	2.169 (2)	C7—C9	1.502 (4)
Zn1—N4	2.172 (3)	C8—H8A	0.9300
Zn1—N4 ⁱ	2.172 (3)	C9—C10	1.496 (4)
O1—C9	1.217 (4)	C10—C11	1.389 (5)
N2—C2	1.137 (4)	C10—C12	1.395 (5)
N3—C1	1.139 (4)	C11—C15	1.395 (5)
N4—C8	1.337 (4)	C11—H11A	0.9300
N4—C4	1.343 (4)	C12—C13	1.388 (5)
N5—C2 ⁱⁱ	1.290 (4)	C12—H12A	0.9300
N5—C1	1.293 (4)	C13—C14	1.372 (6)
C2—N5 ⁱⁱⁱ	1.290 (4)	C13—H13A	0.9300
C4—C5	1.379 (4)	C14—C15	1.378 (6)
C4—H4A	0.9300	C14—H14A	0.9300
C5—C6	1.389 (4)	C15—H15A	0.9300
C5—H5C	0.9300		

N2—Zn1—N2 ⁱ	180.00 (9)	C7—C6—H6A	120.5
N2—Zn1—N3 ⁱ	87.73 (11)	C5—C6—H6A	120.5
N2 ⁱ —Zn1—N3 ⁱ	92.27 (11)	C6—C7—C8	118.0 (3)
N2—Zn1—N3	92.27 (11)	C6—C7—C9	125.3 (3)
N2 ⁱ —Zn1—N3	87.73 (11)	C8—C7—C9	116.6 (3)
N3 ⁱ —Zn1—N3	180.0	N4—C8—C7	123.5 (3)
N2—Zn1—N4	90.79 (9)	N4—C8—H8A	118.3
N2 ⁱ —Zn1—N4	89.21 (9)	C7—C8—H8A	118.3
N3 ⁱ —Zn1—N4	90.13 (9)	O1—C9—C10	118.7 (3)
N3—Zn1—N4	89.87 (9)	O1—C9—C7	118.9 (3)
N2—Zn1—N4 ⁱ	89.21 (9)	C10—C9—C7	122.3 (3)
N2 ⁱ —Zn1—N4 ⁱ	90.79 (9)	C11—C10—C12	119.3 (3)
N3 ⁱ —Zn1—N4 ⁱ	89.87 (9)	C11—C10—C9	123.9 (3)
N3—Zn1—N4 ⁱ	90.13 (9)	C12—C10—C9	116.7 (3)
N4—Zn1—N4 ⁱ	180.00 (4)	C10—C11—C15	120.0 (4)
C2—N2—Zn1	157.0 (2)	C10—C11—H11A	120.0
C1—N3—Zn1	150.9 (2)	C15—C11—H11A	120.0
C8—N4—C4	117.4 (2)	C13—C12—C10	120.1 (4)
C8—N4—Zn1	122.30 (19)	C13—C12—H12A	120.0
C4—N4—Zn1	119.90 (18)	C10—C12—H12A	120.0
C2 ⁱⁱ —N5—C1	123.7 (3)	C14—C13—C12	120.2 (4)
N3—C1—N5	172.8 (3)	C14—C13—H13A	119.9
N2—C2—N5 ⁱⁱⁱ	172.0 (3)	C12—C13—H13A	119.9
N4—C4—C5	123.2 (3)	C13—C14—C15	120.4 (4)
N4—C4—H4A	118.4	C13—C14—H14A	119.8
C5—C4—H4A	118.4	C15—C14—H14A	119.8
C4—C5—C6	118.8 (3)	C14—C15—C11	120.0 (4)
C4—C5—H5C	120.6	C14—C15—H15A	120.0
C6—C5—H5C	120.6	C11—C15—H15A	120.0
C7—C6—C5	119.0 (3)		
N3 ⁱ —Zn1—N2—C2	40.5 (6)	C4—N4—C8—C7	2.1 (4)
N3—Zn1—N2—C2	-139.5 (6)	Zn1—N4—C8—C7	-170.4 (2)
N4—Zn1—N2—C2	130.6 (6)	C6—C7—C8—N4	1.3 (4)
N4 ⁱ —Zn1—N2—C2	-49.4 (6)	C9—C7—C8—N4	178.9 (3)
N2—Zn1—N3—C1	-168.8 (5)	C6—C7—C9—O1	164.8 (4)
N2 ⁱ —Zn1—N3—C1	11.2 (5)	C8—C7—C9—O1	-12.6 (5)
N4—Zn1—N3—C1	-78.0 (5)	C6—C7—C9—C10	-12.8 (5)
N4 ⁱ —Zn1—N3—C1	102.0 (5)	C8—C7—C9—C10	169.8 (3)
N2 ⁱ —Zn1—N4—C8	58.2 (2)	O1—C9—C10—C11	150.0 (4)
N3 ⁱ —Zn1—N4—C8	-34.0 (2)	C7—C9—C10—C11	-32.4 (5)
N3—Zn1—N4—C8	146.0 (2)	O1—C9—C10—C12	-26.4 (5)
N2—Zn1—N4—C4	65.9 (2)	C7—C9—C10—C12	151.2 (3)
N2 ⁱ —Zn1—N4—C4	-114.1 (2)	C12—C10—C11—C15	0.2 (5)
N3 ⁱ —Zn1—N4—C4	153.6 (2)	C9—C10—C11—C15	-176.1 (3)

supplementary materials

N3—Zn1—N4—C4	-26.4 (2)	C11—C10—C12—C13	0.1 (5)
C8—N4—C4—C5	-3.7 (4)	C9—C10—C12—C13	176.6 (3)
Zn1—N4—C4—C5	169.0 (2)	C10—C12—C13—C14	-0.5 (6)
N4—C4—C5—C6	1.7 (5)	C12—C13—C14—C15	0.6 (6)
C4—C5—C6—C7	1.8 (5)	C13—C14—C15—C11	-0.4 (6)
C5—C6—C7—C8	-3.2 (5)	C10—C11—C15—C14	0.0 (5)
C5—C6—C7—C9	179.4 (3)		

Symmetry codes: (i) $-x+1, -y-1, -z$; (ii) $x, y-1, z$; (iii) $x, y+1, z$.

Fig. 1




Fig. 2

