

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-Biimidazolium hexaqua-manganese(II) bis(sulfate)

Mukhtar A. Kurawa, Christopher J. Adams and A. Guy Orpen*

School of Chemistry, University of Bristol, Bristol BS8 1TS, England
Correspondence e-mail: guy.orpen@bris.ac.uk


Received 25 June 2008; accepted 2 July 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.023; wR factor = 0.066; data-to-parameter ratio = 13.8.

The title compound, $(\text{C}_6\text{H}_8\text{N}_4)[\text{Mn}(\text{H}_2\text{O})_6](\text{SO}_4)_2$, was obtained by cocrystallization of 2,2'-biimidazolium sulfate and bis(tetrabutylammonium) tetrachloridomanganate(II). The asymmetric unit contains one isolated $(\text{SO}_4)^{2-}$ anion, one half of an octahedral $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ dication and one half of a 2,2'-biimidazolium dication, each of which lies on an inversion centre. Molecules are connected by a three-dimensional $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen-bond network.

Related literature

For the syntheses, structural studies and thermal behaviour of related compounds, see: Rekik *et al.* (2006, 2007).



Experimental

Crystal data

 $(\text{C}_6\text{H}_8\text{N}_4)[\text{Mn}(\text{H}_2\text{O})_6](\text{SO}_4)_2$ $M_r = 491.34$ Monoclinic, $P2_1/c$ $a = 6.0625$ (7) Å $b = 11.606$ (2) Å $c = 12.218$ (2) Å $\beta = 91.65$ (1)° $V = 859.3$ (2) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 1.09$ mm⁻¹ $T = 100$ (2) K $0.4 \times 0.3 \times 0.2$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.686$, $T_{\max} = 0.800$

9389 measured reflections

1954 independent reflections

1897 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.018$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.023$ $wR(F^2) = 0.066$ $S = 1.06$

1954 reflections

142 parameters

6 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.65$ e Å⁻³

Table 1

Selected bond lengths (Å).

Mn1—O6	2.1335 (10)	Mn1—O5	2.2218 (10)
Mn1—O7	2.1856 (10)		

Symmetry code: (i) $-x, -y + 2, -z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots O3 ⁱⁱ	0.88	1.93	2.7699 (15)	159
N1—H1A \cdots O2 ⁱⁱ	0.88	2.45	3.0994 (15)	131
N2—H2A \cdots O3 ⁱⁱⁱ	0.88	1.92	2.7562 (16)	159
O5—H5A \cdots O2	0.843 (14)	1.917 (15)	2.7600 (15)	176.8 (18)
O5—H5B \cdots O3 ^{iv}	0.813 (14)	2.088 (15)	2.8638 (15)	159.7 (17)
O6—H6A \cdots O4 ^v	0.839 (14)	1.908 (15)	2.7402 (15)	171.2 (18)
O6—H6B \cdots O1 ^{vi}	0.846 (14)	1.847 (15)	2.6904 (14)	174.2 (18)
O7—H7A \cdots O4 ^{vii}	0.851 (14)	1.888 (15)	2.7298 (15)	169.5 (18)
O7—H7B \cdots O2 ^v	0.846 (14)	1.892 (14)	2.7266 (14)	169.0 (17)

Symmetry codes: (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - 1, -y + \frac{1}{2}, z - \frac{1}{2}$; (iv) $x - 1, y, z$; (v) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (vi) $-x + 1, -y + 1, -z$; (vii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

MAK thanks Bayero University, Kano, Nigeria, for funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2142).

References

- Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Rekik, W., Naili, H., Bataille, T., Roisnel, T. & Mhiri, T. (2006). *Inorg. Chim. Acta*, **359**, 3954–3962.
- Rekik, W., Naili, H., Mhiri, T. & Bataille, T. (2007). *J. Chem. Crystallogr.* **37**, 149–155.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1005 [doi:10.1107/S1600536808020291]

2,2'-Biimidazolium hexaaquamanganese(II) bis(sulfate)

M. A. Kurawa, C. J. Adams and A. G. Orpen

Comment

The syntheses, structural studies and thermal behaviour of similar complexes with piperazinium, $(C_4H_{12}N_2)^{2+}$, and 1,4-diaza-bicyclo[2.2.2]octandium, $(C_6H_{14}N_2)^{2+}$, cations have been reported (Rekik *et al.*, 2006, 2007).

In the crystal structure of the title compound (Fig. 1; Table 1), the $(C_6H_8N_4)^{2+}$, $[Mn(H_2O)_6]^{2+}$ and $(SO_4)^{2-}$ ions are connected by N—H \cdots O and O—H \cdots O hydrogen bonds (Table 2), with the 2,2'-biimidazolium dications in the supramolecular cavities formed by the metal–sulfate framework (Fig. 2). The corresponding structures of some first row transition metal M^{II} sulfates ($M = Mn, Ni, Fe$ and Cu) templated with piperazinium display similar three-dimensional hydrogen-bonded networks (Rekik, Naili, Bataille *et al.*, 2006). In particular, the structures of the $(C_4H_{12}N_2)^{2+}[M(H_2O)_6]^{2+}(SO_4)_2^{2-}$ ($M = Mn$ or Ni) compounds contain channels (running parallel to the c -axis in those cases), which are defined by a square arrangement of $[M(H_2O)_6]^{2+}$ cations and which contain the organic dications, mirroring the channels seen in the title compound (Fig. 2).

Experimental

The title compound was obtained unintentionally as the product of an attempted synthesis of a hydrogen-bonded salt of the tetrachloromanganate(II) anion and the biimidazolium cation, using slow evaporation of a water–acetonitrile solution (1:1 v/v) of equimolar amounts of bis(tetrabutylammonium) tetrachloromanganate(II) and 2,2'-biimidazolium sulfate at room temperature.

Refinement

H atoms bonded to O atoms were located in a difference map and refined with distance restraints of O—H = 0.84 (2) Å and with $U_{iso}(H) = 1.2U_{eq}(O)$. Other H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 and N—H = 0.88 Å, and with $U_{iso}(H) = 1.2U_{eq}(C, N)$.

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry codes: (i) $-x, 2-y, -z$; (vi) $1-x, 1-y, -z$.]




Fig. 2. Packing diagram for the title compound viewed along the a -axis.

2,2'-Biimidazolium hexaaquamanganese(II) bis(sulfate)

Crystal data

$(C_6H_8N_4)[Mn(H_2O)_6](SO_4)_2$

$M_r = 491.34$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 6.0625\ (7)\ \text{\AA}$

$b = 11.606\ (2)\ \text{\AA}$

$c = 12.218\ (2)\ \text{\AA}$

$\beta = 91.65\ (1)^\circ$

$V = 859.3\ (2)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 506$

$D_x = 1.899\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 7442 reflections

$\theta = 2.4\text{--}27.5^\circ$

$\mu = 1.09\ \text{mm}^{-1}$

$T = 100\ (2)\ \text{K}$

Block, colourless

$0.4 \times 0.3 \times 0.2\ \text{mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100\ (2)\ \text{K}$

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.686, T_{\max} = 0.800$

9389 measured reflections

1954 independent reflections

1897 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 2.4^\circ$

$h = -7 \rightarrow 7$

$k = -15 \rightarrow 15$

$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.023$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.066$	$w = 1/[\sigma^2(F_o^2) + (0.042P)^2 + 0.4497P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
1954 reflections	$(\Delta/\sigma)_{\max} < 0.001$
142 parameters	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
6 restraints	$\Delta\rho_{\min} = -0.65 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.5000	0.5000	0.0000	0.00892 (9)
S1	0.87954 (5)	0.76694 (3)	0.24181 (2)	0.00829 (10)
N1	0.20621 (19)	1.00486 (9)	0.11407 (9)	0.0107 (2)
H1A	0.1629	1.0584	0.1597	0.013*
N2	0.21706 (17)	0.89187 (9)	-0.02676 (9)	0.0105 (2)
H2A	0.1825	0.8585	-0.0896	0.013*
C1	0.3926 (2)	0.93818 (11)	0.12742 (11)	0.0125 (2)
H1B	0.4969	0.9413	0.1869	0.015*
C2	0.3988 (2)	0.86702 (11)	0.03924 (11)	0.0124 (2)
H2B	0.5082	0.8106	0.0257	0.015*
C3	0.1021 (2)	0.97533 (11)	0.02099 (10)	0.0098 (2)
O1	0.87996 (15)	0.78300 (8)	0.12285 (7)	0.0128 (2)
O2	0.65505 (15)	0.73605 (8)	0.27723 (8)	0.0126 (2)
O3	1.02939 (15)	0.66897 (8)	0.27336 (7)	0.01179 (19)
O4	0.95582 (15)	0.87250 (8)	0.29863 (8)	0.01210 (19)
O5	0.39485 (16)	0.59047 (8)	0.15004 (8)	0.0137 (2)
O6	0.19308 (16)	0.40995 (8)	-0.00527 (8)	0.0135 (2)
O7	0.63321 (15)	0.36233 (8)	0.10452 (8)	0.01286 (19)
H5A	0.474 (3)	0.6368 (14)	0.1868 (14)	0.015*
H6A	0.134 (3)	0.4002 (15)	0.0553 (13)	0.015*
H7A	0.753 (3)	0.3688 (15)	0.1418 (14)	0.015*
H5B	0.273 (2)	0.6050 (15)	0.1724 (14)	0.015*
H6B	0.162 (3)	0.3514 (14)	-0.0441 (14)	0.015*
H7B	0.548 (3)	0.3290 (15)	0.1482 (13)	0.015*

Atomic displacement parameters (\AA^2)

U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
----------	----------	----------	----------	----------	----------

supplementary materials

Mn1	0.00931 (15)	0.00906 (15)	0.00839 (15)	-0.00039 (9)	0.00042 (10)	0.00021 (9)
S1	0.00802 (16)	0.00845 (16)	0.00839 (17)	0.00010 (10)	0.00033 (11)	-0.00019 (10)
N1	0.0120 (5)	0.0100 (5)	0.0103 (5)	0.0002 (4)	0.0001 (4)	-0.0003 (4)
N2	0.0109 (5)	0.0105 (5)	0.0102 (5)	-0.0002 (4)	0.0007 (4)	-0.0012 (4)
C1	0.0128 (6)	0.0120 (6)	0.0127 (6)	0.0003 (4)	-0.0010 (4)	0.0025 (5)
C2	0.0112 (6)	0.0119 (6)	0.0141 (6)	0.0008 (4)	-0.0006 (4)	0.0019 (5)
C3	0.0103 (6)	0.0089 (5)	0.0104 (6)	-0.0016 (5)	0.0015 (4)	0.0009 (4)
O1	0.0167 (5)	0.0125 (4)	0.0090 (4)	-0.0011 (3)	-0.0005 (3)	0.0010 (3)
O2	0.0088 (4)	0.0139 (4)	0.0150 (5)	-0.0014 (3)	0.0021 (3)	-0.0016 (3)
O3	0.0116 (4)	0.0115 (4)	0.0123 (4)	0.0028 (3)	0.0006 (3)	0.0016 (3)
O4	0.0124 (4)	0.0110 (4)	0.0129 (4)	-0.0018 (3)	0.0009 (3)	-0.0027 (3)
O5	0.0106 (4)	0.0166 (5)	0.0141 (5)	-0.0010 (4)	0.0026 (3)	-0.0052 (4)
O6	0.0139 (4)	0.0152 (5)	0.0116 (5)	-0.0036 (4)	0.0027 (3)	-0.0025 (4)
O7	0.0106 (4)	0.0145 (5)	0.0134 (5)	-0.0004 (3)	0.0002 (3)	0.0035 (3)

Geometric parameters (\AA , $^\circ$)

Mn1—O6	2.1335 (10)	N2—C3	1.3371 (16)
Mn1—O6 ⁱ	2.1335 (10)	N2—C2	1.3771 (17)
Mn1—O7 ⁱ	2.1856 (10)	N2—H2A	0.8800
Mn1—O7	2.1856 (10)	C1—C2	1.3589 (19)
Mn1—O5	2.2218 (10)	C1—H1B	0.9500
Mn1—O5 ⁱ	2.2218 (10)	C2—H2B	0.9500
S1—O1	1.4653 (10)	C3—C3 ⁱⁱ	1.445 (2)
S1—O4	1.4759 (10)	O5—H5A	0.843 (14)
S1—O2	1.4839 (10)	O5—H5B	0.813 (14)
S1—O2	1.4839 (10)	O6—H6A	0.839 (14)
S1—O3	1.4989 (9)	O6—H6B	0.846 (14)
N1—C3	1.3296 (17)	O7—H7A	0.851 (14)
N1—C1	1.3754 (17)	O7—H7B	0.846 (14)
N1—H1A	0.8800		
O6—Mn1—O6 ⁱ	180.0	C3—N1—C1	108.94 (11)
O6—Mn1—O7 ⁱ	91.90 (4)	C3—N1—H1A	125.5
O6 ⁱ —Mn1—O7 ⁱ	88.10 (4)	C1—N1—H1A	125.5
O6—Mn1—O7	88.10 (4)	C3—N2—C2	108.33 (11)
O6 ⁱ —Mn1—O7	91.90 (4)	C3—N2—H2A	125.8
O7 ⁱ —Mn1—O7	180.0	C2—N2—H2A	125.8
O6—Mn1—O5	89.18 (4)	C2—C1—N1	106.82 (11)
O6 ⁱ —Mn1—O5	90.82 (4)	C2—C1—H1B	126.6
O7 ⁱ —Mn1—O5	91.53 (4)	N1—C1—H1B	126.6
O7—Mn1—O5	88.47 (4)	C1—C2—N2	107.28 (11)
O6—Mn1—O5 ⁱ	90.82 (4)	C1—C2—H2B	126.4
O6 ⁱ —Mn1—O5 ⁱ	89.18 (4)	N2—C2—H2B	126.4
O7 ⁱ —Mn1—O5 ⁱ	88.47 (4)	N1—C3—N2	108.63 (11)
O7—Mn1—O5 ⁱ	91.53 (4)	N1—C3—C3 ⁱⁱ	125.62 (15)
O5—Mn1—O5 ⁱ	180.0	N2—C3—C3 ⁱⁱ	125.75 (15)

O1—S1—O4	110.58 (6)	Mn1—O5—H5A	124.6 (12)
O1—S1—O2	110.35 (6)	Mn1—O5—H5B	131.4 (13)
O4—S1—O2	109.94 (6)	H5A—O5—H5B	101.5 (17)
O1—S1—O2	110.35 (6)	Mn1—O6—H6A	115.7 (12)
O4—S1—O2	109.94 (6)	Mn1—O6—H6B	126.2 (12)
O1—S1—O3	109.47 (6)	H6A—O6—H6B	107.0 (17)
O4—S1—O3	109.23 (6)	Mn1—O7—H7A	123.0 (12)
O2—S1—O3	107.21 (6)	Mn1—O7—H7B	118.8 (12)
O2—S1—O3	107.21 (6)	H7A—O7—H7B	103.3 (17)
C3—N1—C1—C2	0.07 (15)	C1—N1—C3—C3 ⁱⁱ	179.08 (15)
N1—C1—C2—N2	0.36 (14)	C2—N2—C3—N1	0.71 (14)
C3—N2—C2—C1	-0.66 (14)	C2—N2—C3—C3 ⁱⁱ	-178.85 (15)
C1—N1—C3—N2	-0.48 (14)		

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x, -y+2, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O3 ⁱⁱⁱ	0.88	1.93	2.7699 (15)	159
N1—H1A \cdots O2 ⁱⁱⁱ	0.88	2.45	3.0994 (15)	131
N2—H2A \cdots O3 ^{iv}	0.88	1.92	2.7562 (16)	159
O5—H5A \cdots O2	0.843 (14)	1.917 (15)	2.7600 (15)	176.8 (18)
O5—H5B \cdots O3 ^v	0.813 (14)	2.088 (15)	2.8638 (15)	159.7 (17)
O6—H6A \cdots O4 ^{vi}	0.839 (14)	1.908 (15)	2.7402 (15)	171.2 (18)
O6—H6B \cdots O1 ⁱ	0.846 (14)	1.847 (15)	2.6904 (14)	174.2 (18)
O7—H7A \cdots O4 ^{vii}	0.851 (14)	1.888 (15)	2.7298 (15)	169.5 (18)
O7—H7B \cdots O2 ^{vi}	0.846 (14)	1.892 (14)	2.7266 (14)	169.0 (17)

Symmetry codes: (iii) $-x+1, y+1/2, -z+1/2$; (iv) $x-1, -y+3/2, z-1/2$; (v) $x-1, y, z$; (vi) $-x+1, y-1/2, -z+1/2$; (i) $-x+1, -y+1, -z$; (vii) $-x+2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

