

Bis[μ -4-chloro-2-[(2-pyridylethyl)imino-methyl]phenolato]bis[chloridocopper(II)]

Jianlan Suo

Department of Chemistry, Baoji University of Arts and Sciences, Baoji, Shaanxi 721007, People's Republic of China

Correspondence e-mail: suojn@yahoo.com.cn


Received 23 June 2008; accepted 15 July 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.031; wR factor = 0.082; data-to-parameter ratio = 14.3.

The title compound, $[\text{Cu}_2(\text{C}_{14}\text{H}_{12}\text{ClN}_2\text{O})_2\text{Cl}_2]$, is a copper(II) dimer where the metal centres are bridged by O atoms from a 5-chlorosalicylaldehyde group. The coordination geometry of each copper(II) centre is distorted square-pyramidal, with two N atoms from a 2-ethylaminopyridine group and two O atoms from a 5-chlorosalicylaldehyde group occupying the basal positions, and with a Cl atom at the apical position. The dimer is centrosymmetric, with a crystallographic inversion centre midway between the two Cu atoms [$\text{Cu} \cdots \text{Cu} = 3.103$ (9) Å].

Related literature

For related literature, see: Du *et al.* (2003); Rojas *et al.* (2004); Yamada (1999).



Experimental

Crystal data

 $[\text{Cu}_2(\text{C}_{14}\text{H}_{12}\text{ClN}_2\text{O})_2\text{Cl}_2]$
 $M_r = 717.39$

 Monoclinic, $P2_1/c$
 $a = 9.9703$ (10) Å

 $b = 9.0119$ (11) Å

 $c = 16.5018$ (16) Å

 $\beta = 96.0390$ (10)°

 $V = 1474.5$ (3) Å³
 $Z = 2$

 Mo $K\alpha$ radiation

 $\mu = 1.84$ mm⁻¹
 $T = 298$ (2) K

 $0.50 \times 0.42 \times 0.02$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.460$, $T_{\max} = 0.957$

7139 measured reflections

2587 independent reflections

 2101 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.082$
 $S = 1.05$

2587 reflections

181 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.32$ e Å⁻³
Table 1

Selected bond lengths (Å).

Cu1—O1 ⁱ	1.9547 (18)	Cu1—Cl2	2.3187 (9)
Cu1—N2	1.958 (2)		

 Symmetry code: (i) $-x + 1, -y + 2, -z$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The author thanks the Science and Educational Fund of Shaanxi Province for a research grant (No. 06k16-G16).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GW2043).

References

- Du, M., Guo, Y.-M., Chen, S.-T., Bu, X.-H. & Ribas, J. (2003). *Inorg. Chim. Acta*, **346**, 207–214.
 Rojas, D., García, A. M., Vega, A., Moreno, Y., Venegas-Yazigi, D., Garland, M. T. & Manzur, J. (2004). *Inorg. Chem.* **43**, 6324–6330.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Yamada, S. (1999). *Coord. Chem. Rev.* **190–192**, 537–555.

supplementary materials

Acta Cryst. (2008). E64, m1046 [doi:10.1107/S1600536808022162]

Bis{ μ -4-chloro-2-[(2-pyridylethyl)iminomethyl]phenolato}bis[chloridocopper(II)]

J. Suo

Comment

Transition metal complexes containing Schiff base ligands have been of great interest for many years (Yamada, 1999). These complexes play an important role in the coordination chemistry related to catalysis and enzymatic reactions, magnetism and molecular architectures. The complexes of salicylaldehyde with polyamines and bis(phenoxy) bridged dinuclear copper(II) complexes are rare. As an extension of the work on the structural characterization of Schiff base complexes, the crystal structure of a mononuclear copper(II) compound, (I), is reported here.

The molecular structure of complex (I) is defined by two [CuLCl] units [4-chloro-2-pyridylethylamine-phenolato], which are bridged by two atoms from 5-Chlorosalicylaldehyde, in such a way as to define a central $N_2CuO_2CuN_2$ core. Additionally, there is an Cl atom from $CuCl_2 \cdot 2H_2O$ completing the pentacoordination of each Cu atom, thus defining a slightly distorted square-based pyramidal coordination for the metal centres. The basal square of the pyramid is defined by two N atoms (N1 and N2) from 2-ethylaminopyridine and two O atoms from 5-Chlorosalicylaldehyde [O1 and O1A; symmetry code: (A)-x + 1, y, -z].

The Cu—Cl2 distance is 2.3187 (9) Å, which is a rather long value for the normal length of this kind of bond (2.0512 Å). A similar value has been reported for $[Cu_2(m\text{-oxalato})(dipyridylamino)_2(CH_3-CN)_2](ClO_4)_2$ (Du *et al.*, 2003). The Cu—Cu distance of 3.103 (9) Å is close to this kind of complex (Rojas *et al.*, 2004). Consistently, the O—Cu—O1A angle is 78.44 (8)°. The atom sequence Cu—O1—Cu1A—O1A is a rather parallelogram. The Cu—O1 and Cu—O1A distances are 1.9547 (18) Å and 2.0500 (19) Å, respectively.

Experimental

5-Chlorosalicylaldehyde (0.1 mmol, 15.7 mg), $CuCl_2 \cdot 2H_2O$ (0.1 mmol, 17.05 mg) and 2-ethylaminopyridine (0.1 mmol, 122.2 mg) were dissolved in methanol (10 ml). The mixture was stirred for 30 min at room temperature to give a clear brown solution. After allowing the resulting solution to stand in air for 11 d, brown block-shaped crystals of (I) were formed on slow evaporation of the solvent. The crystals were collected, washed with methanol and dried in a vacuum desiccator using anhydrous $CaCl_2$ (yield 54%). Analysis found: C 46.84%, H 3.35%, N 7.80%. Calculated for $Cu_2(C_{14}H_{12}N_2OCl_2)_2Cl_2$: C 46.86%, H 3.35%, N 7.81%.

Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances in the range 0.93–0.97 Å and $U_{iso}(H) = 1.2U_{eq}$ or $1.5U_{eq}(C/O)$

Figures




Fig. 1. The structure of the title compound in 30% probability ellipsoids. H atoms are omitted for clarity. [Symmetry code: (A)- $x + 1, y, -z + 1$]




Fig. 2. The molecular packing of (I) viewed along the b axis.

Bis{ μ -4-chloro-2-[(2-pyridylethyl)iminomethyl]phenolato}bis[chloridocopper(II)]

Crystal data

[Cu₂(C₁₄H₁₂Cl₁N₂O)₂Cl₂]

$M_r = 717.39$

Monoclinic, $P2_1/c$

$a = 9.9703$ (10) Å

$b = 9.0119$ (11) Å

$c = 16.5018$ (16) Å

$\beta = 96.0390$ (10)°

$V = 1474.5$ (3) Å³

$Z = 2$

$F_{000} = 724$

$D_x = 1.616$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3383 reflections

$\theta = 2.3$ – 28.1 °

$\mu = 1.84$ mm⁻¹

$T = 298$ (2) K

Block, brown

$0.50 \times 0.42 \times 0.02$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ (2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.460$, $T_{\max} = 0.957$

7139 measured reflections

2587 independent reflections

2101 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 25.0$ °

$\theta_{\text{min}} = 2.1$ °

$h = -11 \rightarrow 11$

$k = -10 \rightarrow 10$

$l = -14 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.031$	H-atom parameters constrained
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0359P)^2 + 0.6685P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2587 reflections	$(\Delta/\sigma)_{\max} = 0.001$
181 parameters	$\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.40654 (3)	0.92113 (4)	0.05596 (2)	0.03066 (13)
Cl1	1.06351 (9)	0.91583 (13)	0.29917 (6)	0.0704 (3)
Cl2	0.29315 (9)	1.11255 (9)	0.11335 (5)	0.0489 (2)
N1	0.3451 (2)	0.7177 (2)	0.01064 (14)	0.0326 (5)
N2	0.4538 (2)	0.8209 (2)	0.16031 (13)	0.0328 (5)
O1	0.60250 (18)	0.9870 (2)	0.05178 (11)	0.0332 (5)
C1	0.3444 (3)	0.7399 (4)	0.19511 (18)	0.0451 (8)
H1A	0.3116	0.7990	0.2380	0.054*
H1B	0.3796	0.6477	0.2191	0.054*
C2	0.2278 (3)	0.7060 (3)	0.13033 (19)	0.0431 (8)
H2A	0.1642	0.6415	0.1538	0.052*
H2B	0.1814	0.7979	0.1147	0.052*
C3	0.2706 (3)	0.6337 (3)	0.05563 (18)	0.0378 (7)
C4	0.2375 (4)	0.4895 (4)	0.0332 (2)	0.0553 (9)
H4	0.1859	0.4316	0.0648	0.066*
C5	0.2823 (5)	0.4327 (4)	-0.0368 (3)	0.0684 (12)
H5	0.2613	0.3357	-0.0526	0.082*
C6	0.3568 (4)	0.5184 (4)	-0.0824 (2)	0.0557 (10)

supplementary materials

H6	0.3866	0.4812	-0.1300	0.067*
C7	0.3882 (3)	0.6617 (3)	-0.05733 (18)	0.0423 (8)
H7	0.4401	0.7205	-0.0882	0.051*
C8	0.5714 (3)	0.8170 (3)	0.19992 (16)	0.0346 (7)
H8	0.5789	0.7624	0.2480	0.042*
C9	0.6936 (3)	0.8864 (3)	0.17933 (16)	0.0304 (6)
C10	0.7074 (3)	0.9662 (3)	0.10664 (16)	0.0304 (6)
C11	0.8341 (3)	1.0216 (3)	0.09434 (18)	0.0392 (7)
H11	0.8453	1.0702	0.0458	0.047*
C12	0.9435 (3)	1.0062 (4)	0.15249 (19)	0.0429 (8)
H12	1.0267	1.0463	0.1439	0.051*
C13	0.9277 (3)	0.9303 (3)	0.22358 (19)	0.0418 (8)
C14	0.8076 (3)	0.8687 (3)	0.23679 (18)	0.0387 (7)
H14	0.8005	0.8144	0.2841	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0349 (2)	0.0311 (2)	0.02574 (19)	-0.00498 (15)	0.00216 (14)	0.00274 (14)
Cl1	0.0410 (5)	0.1083 (9)	0.0574 (6)	0.0051 (5)	-0.0164 (4)	0.0100 (5)
Cl2	0.0554 (5)	0.0391 (5)	0.0541 (5)	0.0034 (4)	0.0150 (4)	-0.0099 (4)
N1	0.0365 (14)	0.0285 (13)	0.0315 (13)	-0.0010 (11)	-0.0028 (10)	0.0010 (10)
N2	0.0392 (14)	0.0313 (13)	0.0285 (12)	-0.0062 (11)	0.0061 (10)	0.0022 (10)
O1	0.0299 (11)	0.0400 (11)	0.0286 (10)	-0.0046 (9)	-0.0018 (8)	0.0094 (9)
C1	0.048 (2)	0.053 (2)	0.0358 (17)	-0.0115 (16)	0.0077 (14)	0.0078 (15)
C2	0.0405 (18)	0.0408 (18)	0.0485 (19)	-0.0105 (15)	0.0071 (14)	0.0110 (15)
C3	0.0378 (18)	0.0288 (16)	0.0441 (18)	-0.0024 (13)	-0.0078 (14)	0.0052 (13)
C4	0.067 (2)	0.0361 (19)	0.059 (2)	-0.0146 (17)	-0.0094 (18)	0.0057 (17)
C5	0.090 (3)	0.032 (2)	0.077 (3)	-0.001 (2)	-0.022 (2)	-0.0097 (19)
C6	0.073 (3)	0.042 (2)	0.050 (2)	0.0119 (19)	-0.0068 (18)	-0.0130 (17)
C7	0.0445 (19)	0.0434 (19)	0.0376 (17)	0.0064 (15)	-0.0026 (14)	-0.0032 (14)
C8	0.0470 (19)	0.0328 (16)	0.0236 (14)	0.0004 (14)	0.0017 (12)	0.0027 (12)
C9	0.0344 (16)	0.0293 (15)	0.0270 (15)	-0.0006 (12)	0.0009 (12)	-0.0026 (11)
C10	0.0345 (16)	0.0268 (15)	0.0293 (15)	-0.0010 (12)	0.0006 (12)	-0.0015 (11)
C11	0.0358 (18)	0.0423 (18)	0.0393 (17)	-0.0025 (14)	0.0028 (13)	0.0081 (14)
C12	0.0275 (16)	0.0469 (19)	0.053 (2)	-0.0012 (14)	-0.0005 (14)	0.0035 (16)
C13	0.0336 (18)	0.0494 (19)	0.0402 (18)	0.0040 (15)	-0.0070 (13)	-0.0002 (15)
C14	0.0416 (19)	0.0407 (17)	0.0321 (16)	0.0058 (14)	-0.0031 (13)	0.0030 (13)

Geometric parameters (\AA , $^\circ$)

Cu1—O1 ⁱ	1.9547 (18)	C4—C5	1.381 (5)
Cu1—N2	1.958 (2)	C4—H4	0.9300
Cu1—N1	2.049 (2)	C5—C6	1.353 (6)
Cu1—O1	2.0500 (19)	C5—H5	0.9300
Cu1—Cl2	2.3187 (9)	C6—C7	1.382 (5)
Cl1—C13	1.747 (3)	C6—H6	0.9300
N1—C3	1.339 (4)	C7—H7	0.9300

N1—C7	1.341 (4)	C8—C9	1.441 (4)
N2—C8	1.282 (4)	C8—H8	0.9300
N2—C1	1.478 (4)	C9—C14	1.411 (4)
O1—C10	1.323 (3)	C9—C10	1.417 (4)
O1—Cu1 ⁱ	1.9547 (18)	C10—C11	1.392 (4)
C1—C2	1.525 (4)	C11—C12	1.382 (4)
C1—H1A	0.9700	C11—H11	0.9300
C1—H1B	0.9700	C12—C13	1.381 (4)
C2—C3	1.495 (4)	C12—H12	0.9300
C2—H2A	0.9700	C13—C14	1.358 (4)
C2—H2B	0.9700	C14—H14	0.9300
C3—C4	1.382 (4)		
O1 ⁱ —Cu1—N2	168.33 (9)	C5—C4—C3	118.9 (4)
O1 ⁱ —Cu1—N1	93.65 (9)	C5—C4—H4	120.5
N2—Cu1—N1	86.74 (9)	C3—C4—H4	120.5
O1 ⁱ —Cu1—O1	78.44 (8)	C6—C5—C4	119.9 (3)
N2—Cu1—O1	91.22 (8)	C6—C5—H5	120.0
N1—Cu1—O1	119.75 (9)	C4—C5—H5	120.0
O1 ⁱ —Cu1—C12	94.50 (6)	C5—C6—C7	119.1 (3)
N2—Cu1—C12	93.82 (7)	C5—C6—H6	120.4
N1—Cu1—C12	132.43 (7)	C7—C6—H6	120.4
O1—Cu1—C12	107.80 (6)	N1—C7—C6	121.3 (3)
C3—N1—C7	119.7 (3)	N1—C7—H7	119.3
C3—N1—Cu1	117.75 (19)	C6—C7—H7	119.3
C7—N1—Cu1	122.2 (2)	N2—C8—C9	128.2 (3)
C8—N2—C1	117.5 (2)	N2—C8—H8	115.9
C8—N2—Cu1	125.7 (2)	C9—C8—H8	115.9
C1—N2—Cu1	116.79 (18)	C14—C9—C10	118.9 (3)
C10—O1—Cu1 ⁱ	129.79 (18)	C14—C9—C8	115.7 (3)
C10—O1—Cu1	128.60 (17)	C10—C9—C8	125.4 (2)
Cu1 ⁱ —O1—Cu1	101.56 (8)	O1—C10—C11	120.9 (3)
N2—C1—C2	111.4 (2)	O1—C10—C9	120.7 (3)
N2—C1—H1A	109.3	C11—C10—C9	118.4 (3)
C2—C1—H1A	109.3	C12—C11—C10	121.6 (3)
N2—C1—H1B	109.3	C12—C11—H11	119.2
C2—C1—H1B	109.3	C10—C11—H11	119.2
H1A—C1—H1B	108.0	C13—C12—C11	119.2 (3)
C3—C2—C1	113.7 (3)	C13—C12—H12	120.4
C3—C2—H2A	108.8	C11—C12—H12	120.4
C1—C2—H2A	108.8	C14—C13—C12	121.3 (3)
C3—C2—H2B	108.8	C14—C13—C11	119.0 (2)
C1—C2—H2B	108.8	C12—C13—C11	119.6 (3)
H2A—C2—H2B	107.7	C13—C14—C9	120.5 (3)
N1—C3—C4	121.0 (3)	C13—C14—H14	119.7
N1—C3—C2	115.7 (3)	C9—C14—H14	119.7
C4—C3—C2	123.3 (3)		
O1 ⁱ —Cu1—N1—C3	-146.8 (2)	C1—C2—C3—N1	-66.8 (3)

supplementary materials

N2—Cu1—N1—C3	44.9 (2)	C1—C2—C3—C4	112.9 (3)
O1—Cu1—N1—C3	134.4 (2)	N1—C3—C4—C5	-0.2 (5)
Cl2—Cu1—N1—C3	-47.3 (2)	C2—C3—C4—C5	-179.8 (3)
O1 ⁱ —Cu1—N1—C7	40.0 (2)	C3—C4—C5—C6	-0.3 (6)
N2—Cu1—N1—C7	-128.3 (2)	C4—C5—C6—C7	0.7 (6)
O1—Cu1—N1—C7	-38.7 (2)	C3—N1—C7—C6	0.2 (4)
Cl2—Cu1—N1—C7	139.5 (2)	Cu1—N1—C7—C6	173.2 (2)
O1 ⁱ —Cu1—N2—C8	28.7 (6)	C5—C6—C7—N1	-0.7 (5)
N1—Cu1—N2—C8	121.0 (2)	C1—N2—C8—C9	-179.8 (3)
O1—Cu1—N2—C8	1.2 (2)	Cu1—N2—C8—C9	1.8 (4)
Cl2—Cu1—N2—C8	-106.7 (2)	N2—C8—C9—C14	177.1 (3)
O1 ⁱ —Cu1—N2—C1	-149.7 (4)	N2—C8—C9—C10	-4.4 (5)
N1—Cu1—N2—C1	-57.5 (2)	Cu1 ⁱ —O1—C10—C11	4.6 (4)
O1—Cu1—N2—C1	-177.2 (2)	Cu1—O1—C10—C11	-178.7 (2)
Cl2—Cu1—N2—C1	74.8 (2)	Cu1 ⁱ —O1—C10—C9	-175.26 (18)
O1 ⁱ —Cu1—O1—C10	-177.4 (3)	Cu1—O1—C10—C9	1.4 (4)
N2—Cu1—O1—C10	-2.9 (2)	C14—C9—C10—O1	-179.1 (3)
N1—Cu1—O1—C10	-89.8 (2)	C8—C9—C10—O1	2.5 (4)
Cl2—Cu1—O1—C10	91.5 (2)	C14—C9—C10—C11	1.1 (4)
O1 ⁱ —Cu1—O1—Cu1 ⁱ	0.0	C8—C9—C10—C11	-177.4 (3)
N2—Cu1—O1—Cu1 ⁱ	174.53 (10)	O1—C10—C11—C12	177.3 (3)
N1—Cu1—O1—Cu1 ⁱ	87.59 (11)	C9—C10—C11—C12	-2.9 (4)
Cl2—Cu1—O1—Cu1 ⁱ	-91.06 (8)	C10—C11—C12—C13	1.8 (5)
C8—N2—C1—C2	-159.3 (3)	C11—C12—C13—C14	1.1 (5)
Cu1—N2—C1—C2	19.3 (3)	C11—C12—C13—C11	-177.7 (2)
N2—C1—C2—C3	50.8 (4)	C12—C13—C14—C9	-2.9 (5)
C7—N1—C3—C4	0.2 (4)	C11—C13—C14—C9	175.9 (2)
Cu1—N1—C3—C4	-173.1 (2)	C10—C9—C14—C13	1.7 (4)
C7—N1—C3—C2	179.8 (3)	C8—C9—C14—C13	-179.7 (3)
Cu1—N1—C3—C2	6.5 (3)		

Symmetry codes: (i) $-x+1, -y+2, -z$.

Fig. 1




Fig. 2

