

## Tetraaqua(1,10-phenanthroline- $\kappa^2N,N'$ )-magnesium(II) bis[(2,4-dichlorophenyl)acetate]

Xiao-Min Hao,<sup>a\*</sup> Chang-Sheng Gu,<sup>a</sup> Weng-Dong Song<sup>a</sup> and Ji-Wei Liu<sup>b</sup>

<sup>a</sup>Department of Applied Chemistry, Guangdong Ocean University, Zhanjiang 524088, People's Republic of China, and <sup>b</sup>College of Chemistry and Chemical Technology, Daqing Petroleum Institute, Daqing 163318, People's Republic of China

Correspondence e-mail: liujiwei76@126.com


Received 22 May 2008; accepted 15 July 2008

Key indicators: single-crystal X-ray study;  $T = 273$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.043;  $wR$  factor = 0.118; data-to-parameter ratio = 18.0.

In the mononuclear title complex,  $[\text{Mg}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_4] \cdot (\text{C}_8\text{H}_5\text{Cl}_2\text{O}_2)_2$ , each  $\text{Mg}^{\text{II}}$  ion is hexacoordinated by two N atoms from a 1,10-phenanthroline ligand [ $\text{Mg}-\text{N} = 2.233$  (2) Å] and four water molecules [ $\text{Mg}-\text{OW} = 2.033$  (2) and 2.043 (1) Å] in a distorted octahedral geometry. A twofold rotation axis passes through the Mg atom. In the crystal structure, the cations and anions are linked by intermolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds and  $\pi-\pi$  stacking interactions [centroid-centroid distance = 3.804 (2) Å] into layers parallel to the  $ac$  plane.

### Related literature

For related literature, see: Castellari *et al.* (1999); Kopylovich *et al.* (2003); Sharma *et al.* (2007); Zhou *et al.* (2007).



### Experimental

#### Crystal data

$[\text{Mg}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_4] \cdot (\text{C}_8\text{H}_5\text{Cl}_2\text{O}_2)_2$

$M_r = 684.62$   
Monoclinic,  $C2/c$

$a = 28.926$  (1) Å  
 $b = 14.0447$  (6) Å  
 $c = 7.6074$  (3) Å  
 $\beta = 94.785$  (1)°  
 $V = 3079.8$  (2) Å<sup>3</sup>

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.46$  mm<sup>-1</sup>  
 $T = 273$  (2) K  
 $0.34 \times 0.26 \times 0.18$  mm

#### Data collection

Bruker P4 diffractometer  
Absorption correction: empirical  
[OR multi-scan](using intensity measurements)  
(SADABS; Sheldrick, 1996)  
 $T_{\text{min}} = 0.867$ ,  $T_{\text{max}} = 0.921$

10955 measured reflections  
3732 independent reflections  
2619 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.117$   
 $S = 1.02$   
3732 reflections  
207 parameters  
6 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.36$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.47$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2W}-\text{H2W1}\cdots\text{O1}^{\text{i}}$	0.849 (9)	1.876 (9)	2.725 (2)	178 (2)
$\text{O2W}-\text{H2W2}\cdots\text{O1}^{\text{ii}}$	0.841 (9)	1.96 (1)	2.772 (2)	161 (2)
$\text{O1W}-\text{H1W1}\cdots\text{O2}^{\text{iii}}$	0.851 (9)	1.91 (1)	2.728 (2)	162 (2)
$\text{O1W}-\text{H1W2}\cdots\text{O2}^{\text{i}}$	0.849 (9)	1.84 (1)	2.685 (2)	171 (2)

Symmetry codes: (i)  $x, y, z - 1$ ; (ii)  $-x + 1, y, -z + \frac{1}{2}$ ; (iii)  $x, -y + 1, z - \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

The authors thank the Guangdong Ocean University Project (No. 0612178 and No. 0612179), the Zhanjiang City Technology Tender Project (No. 0810014) and Guangdong Ocean University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2068).

### References

- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
Castellari, C., Comelli, F. & Ottani, S. (1999). *Acta Cryst.* **C55**, 1054–1056.  
Kopylovich, M. N., Pombeiro, A. J. L., Fischer, A., Kloo, L. & Kukushkin, V. Yu. (2003). *Inorg. Chem.* **42**, 7239–7248.  
Sharma, R., Sharma, R. P., Balaa, R. & Kariuki, B. M. (2007). *J. Mol. Struct.* **826**, 177–184.  
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Zhou, J., Sun, C. & Jin, L. (2007). *J. Mol. Struct.* **832**, 55–62.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1052 [ doi:10.1107/S1600536808022150 ]

## Tetraaqua(1,10-phenanthroline- $\kappa^2N,N'$ )magnesium(II) bis[(2,4-dichlorophenyl)acetate]

X.-M. Hao, C.-S. Gu, W.-D. Song and J.-W. Liu

### Comment

The rigid-type phenylacetic acid and its derivatives have versatile binding abilities and the potential capabilities of generating complicated supramolecular architectures. Nevertheless, main group or transition metal compounds in which the phenylacetate ion does not coordinate the metals are rare (Castellari *et al.* (1999), Kopylovich *et al.* (2003), Sharma *et al.* (2007), Zhou *et al.* (2007)). In the present study, we chose 2,4-dichlorophenylacetate as the anion to prepare a new mono-nuclear magnesium<sup>II</sup> complex, [Mg(C<sub>12</sub>H<sub>8</sub>N<sub>2</sub>)(H<sub>2</sub>O)<sub>4</sub>][(C<sub>8</sub>H<sub>5</sub>Cl<sub>2</sub>O<sub>2</sub>)<sub>2</sub>], (I), the crystal structure of which is reported here.

As illustrated in Fig. 1, the asymmetric unit of (I) consists of one half of a [Mg(1,10-phen)(H<sub>2</sub>O)<sub>4</sub>] cation and one 2,4-dichlorophenylacetate anion. The Mg<sup>II</sup> atom displays a distorted octahedral geometry defined by two N atoms from the 1,10-phenanthroline ligand [Mg—N 2.233 (2) Å] and four water molecules [Mg—O<sub>w</sub> 2.033 (2), 2.043 (1) Å], respectively. The characteristic C—O(carboxylate) bond lengths suggest electron delocalization in the carboxylate groups of the anionic moieties. In the crystal structure, the cations and anions are linked by intermolecular O—H...O hydrogen bonds between the carboxylate O atoms and the coordinated water molecules. Additional  $\pi$ - $\pi$  stacking interactions between 1,10-phen ligands with a distance of 3.804 (2) Å leads to the observation of layers parallel to the *ac*-plane (details see Table 1 and Fig. 2).

### Experimental

Benzyloxyacetic acid is commercially available and was used without further purification. The title complex was prepared by the addition of Mg(Cl)<sub>2</sub> × 6 H<sub>2</sub>O (4.06 g, 20 mmol) and 1,10-phenanthroline (3.98 g, 20 mmol) to a hot aqueous solution of 2,4-dichlorophenylacetic acid (4.10 g, 20 mmol); the pH was adjusted to 6 with 0.1M sodium hydroxide. The solution was allowed to evaporate at room temperature. Colorless crystals separated from the filtered solution after several days. CHN analysis: Calcd. for C<sub>28</sub>H<sub>26</sub>N<sub>2</sub>O<sub>8</sub>Cl<sub>4</sub>Mg: C 49.12, H 3.83, N 4.09%. Found: C 49.14, H 3.82, 4.08%.

### Refinement

The H atoms attached to C atoms were placed in calculated positions, with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The H atoms of water molecule were located in a difference Fourier map and refined with O—H distance restraint of 0.85 (1) Å, and  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ .

## Figures




Fig. 1. Molecular structure of (I) with 30% probability ellipsoids.




Fig. 2. Packing diagram of (I).

## Tetraaqua(1,10-phenanthroline- $\kappa^2N,N'$ )magnesium(II) bis[(2,4-dichlorophenyl)acetate]

### Crystal data

[Mg(C<sub>12</sub>H<sub>8</sub>N<sub>2</sub>)(H<sub>2</sub>O)<sub>4</sub>](C<sub>8</sub>H<sub>5</sub>Cl<sub>2</sub>O<sub>2</sub>)<sub>2</sub>

$M_r = 684.62$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 28.926 (1) \text{ \AA}$

$b = 14.0447 (6) \text{ \AA}$

$c = 7.6074 (3) \text{ \AA}$

$\beta = 94.785 (1)^\circ$

$V = 3079.8 (2) \text{ \AA}^3$

$Z = 4$

$F_{000} = 1408$

$D_x = 1.477 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 10366 reflections

$\theta = 2.8\text{--}28.2^\circ$

$\mu = 0.46 \text{ mm}^{-1}$

$T = 273 (2) \text{ K}$

Prism, colorless

$0.34 \times 0.26 \times 0.18 \text{ mm}$

### Data collection

Bruker P4  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution:  $10.000 \text{ pixels mm}^{-1}$

$T = 273(2) \text{ K}$

$\omega$  scans

Absorption correction: empirical (using intensity  
measurements)

(SADABS; Sheldrick, 1996)

3732 independent reflections

2619 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\text{max}} = 28.2^\circ$

$\theta_{\text{min}} = 2.8^\circ$

$h = -38 \rightarrow 33$

$k = -18 \rightarrow 18$

$T_{\min} = 0.867$ ,  $T_{\max} = 0.921$   
10955 measured reflections

$l = -9 \rightarrow 10$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.118$

$S = 1.03$

3732 reflections

207 parameters

6 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0514P)^2 + 1.7903P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$

Extinction correction: SHELXL97 (Sheldrick, 2008)

Extinction coefficient: 0.0001 (1)

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mg1	0.5000	0.70119 (6)	0.2500	0.0377 (2)
Cl1	0.30958 (2)	0.68124 (5)	0.95908 (8)	0.06695 (19)
Cl2	0.25982 (3)	0.96821 (5)	0.52259 (13)	0.0957 (3)
O1	0.42857 (5)	0.68303 (10)	0.76580 (19)	0.0501 (3)
O1W	0.44481 (6)	0.61206 (10)	0.20734 (19)	0.0613 (4)
O2W	0.50764 (5)	0.69659 (11)	-0.01445 (17)	0.0480 (3)
O2	0.41678 (5)	0.54277 (10)	0.88709 (19)	0.0544 (4)
N1	0.45496 (5)	0.82808 (11)	0.1896 (2)	0.0423 (4)
C1	0.41135 (7)	0.82785 (17)	0.1205 (3)	0.0577 (6)
H1A	0.3966	0.7697	0.0991	0.069*
C2	0.38664 (9)	0.9115 (2)	0.0787 (4)	0.0733 (7)
H2A	0.3562	0.9084	0.0293	0.088*
C3	0.40715 (9)	0.9969 (2)	0.1101 (3)	0.0705 (7)
H3A	0.3907	1.0527	0.0846	0.085*
C4	0.45322 (8)	1.00081 (15)	0.1812 (3)	0.0552 (5)
C5	0.47614 (6)	0.91362 (12)	0.2161 (2)	0.0392 (4)
C6	0.47783 (10)	1.08762 (15)	0.2176 (4)	0.0734 (8)
H6A	0.4627	1.1453	0.1957	0.088*
C7	0.33544 (7)	0.68867 (14)	0.6254 (3)	0.0457 (5)
C8	0.31108 (6)	0.73310 (14)	0.7514 (3)	0.0456 (4)
C9	0.28790 (7)	0.81845 (15)	0.7222 (3)	0.0544 (5)
H9A	0.2720	0.8466	0.8101	0.065*
C10	0.28898 (7)	0.86044 (16)	0.5596 (3)	0.0603 (6)
C11	0.31228 (9)	0.81981 (19)	0.4301 (3)	0.0698 (7)
H11A	0.3128	0.8491	0.3206	0.084*
C12	0.33515 (8)	0.73431 (19)	0.4637 (3)	0.0628 (6)

## supplementary materials

---

H12A	0.3508	0.7066	0.3750	0.075*
C13	0.36083 (7)	0.59691 (15)	0.6623 (3)	0.0532 (5)
H13A	0.3682	0.5690	0.5514	0.064*
H13B	0.3406	0.5530	0.7176	0.064*
C14	0.40558 (7)	0.60887 (14)	0.7815 (2)	0.0413 (4)
H2W1	0.4831 (4)	0.6910 (16)	-0.083 (2)	0.062*
H2W2	0.5312 (4)	0.6889 (16)	-0.071 (2)	0.062*
H1W1	0.4397 (8)	0.5689 (12)	0.2818 (19)	0.062*
H1W2	0.4390 (8)	0.5895 (14)	0.1044 (13)	0.062*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Mg1	0.0472 (5)	0.0291 (4)	0.0373 (4)	0.000	0.0060 (4)	0.000
Cl1	0.0679 (4)	0.0739 (4)	0.0613 (4)	0.0017 (3)	0.0193 (3)	0.0105 (3)
Cl2	0.0834 (5)	0.0587 (4)	0.1391 (7)	0.0047 (3)	-0.0259 (5)	0.0226 (4)
O1	0.0404 (7)	0.0532 (8)	0.0568 (9)	-0.0033 (6)	0.0054 (6)	0.0051 (6)
O1W	0.0958 (12)	0.0482 (8)	0.0394 (8)	-0.0322 (8)	0.0021 (8)	0.0015 (6)
O2W	0.0441 (8)	0.0631 (9)	0.0373 (7)	0.0031 (7)	0.0062 (6)	0.0027 (6)
O2	0.0743 (10)	0.0418 (7)	0.0463 (8)	0.0029 (7)	-0.0003 (7)	-0.0041 (6)
N1	0.0421 (9)	0.0402 (8)	0.0452 (9)	0.0014 (7)	0.0074 (7)	0.0005 (7)
C1	0.0458 (12)	0.0650 (14)	0.0623 (14)	0.0013 (10)	0.0035 (10)	0.0007 (11)
C2	0.0489 (13)	0.095 (2)	0.0760 (17)	0.0240 (14)	0.0029 (11)	0.0105 (15)
C3	0.0744 (17)	0.0695 (16)	0.0687 (16)	0.0356 (14)	0.0129 (13)	0.0151 (13)
C4	0.0762 (15)	0.0408 (10)	0.0513 (12)	0.0170 (10)	0.0204 (11)	0.0065 (9)
C5	0.0504 (10)	0.0320 (8)	0.0368 (9)	0.0040 (8)	0.0137 (7)	0.0022 (7)
C6	0.114 (2)	0.0323 (10)	0.0777 (18)	0.0144 (11)	0.0287 (16)	0.0067 (11)
C7	0.0366 (10)	0.0509 (11)	0.0486 (11)	-0.0043 (8)	-0.0024 (8)	-0.0053 (9)
C8	0.0377 (10)	0.0482 (11)	0.0509 (11)	-0.0053 (8)	0.0037 (8)	0.0014 (9)
C9	0.0407 (11)	0.0514 (12)	0.0710 (15)	-0.0013 (9)	0.0041 (10)	-0.0032 (10)
C10	0.0440 (12)	0.0514 (12)	0.0827 (17)	-0.0039 (10)	-0.0108 (11)	0.0088 (12)
C11	0.0623 (15)	0.0838 (18)	0.0612 (15)	-0.0099 (13)	-0.0061 (12)	0.0236 (13)
C12	0.0558 (13)	0.0827 (17)	0.0498 (12)	-0.0009 (12)	0.0034 (10)	-0.0016 (12)
C13	0.0486 (12)	0.0500 (12)	0.0602 (13)	0.0000 (9)	-0.0006 (9)	-0.0155 (10)
C14	0.0430 (10)	0.0424 (10)	0.0395 (10)	0.0058 (8)	0.0095 (8)	-0.0086 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Mg1—N1	2.2327 (16)	C3—C4	1.397 (3)
Mg1—O1W	2.0333 (15)	C3—H3A	0.9300
Mg1—O2W	2.0432 (13)	C4—C5	1.407 (3)
Mg1—N1 <sup>i</sup>	2.2327 (16)	C4—C6	1.427 (3)
Mg1—O1W <sup>i</sup>	2.0333 (15)	C5—C5 <sup>i</sup>	1.433 (4)
Mg1—O2W <sup>i</sup>	2.0432 (13)	C6—C6 <sup>i</sup>	1.335 (6)
Cl1—C8	1.744 (2)	C6—H6A	0.9300
Cl2—C10	1.744 (2)	C7—C8	1.385 (3)
N1—C1	1.325 (3)	C7—C12	1.386 (3)
N1—C5	1.356 (2)	C7—C13	1.498 (3)

O1—C14	1.247 (2)	C8—C9	1.382 (3)
O2—C14	1.252 (2)	C9—C10	1.373 (3)
O1W—H1W1	0.851 (9)	C9—H9A	0.9300
O1W—H1W2	0.849 (9)	C10—C11	1.365 (4)
O2W—H2W1	0.849 (9)	C11—C12	1.384 (3)
O2W—H2W2	0.841 (9)	C11—H11A	0.9300
C1—C2	1.399 (3)	C12—H12A	0.9300
C1—H1A	0.9300	C13—C14	1.527 (3)
C2—C3	1.350 (4)	C13—H13A	0.9700
C2—H2A	0.9300	C13—H13B	0.9700
N1 <sup>i</sup> —Mg1—N1	74.07 (8)	C3—C4—C6	123.6 (2)
O1W—Mg1—N1	91.24 (6)	C4—C3—H3A	120.2
O1W <sup>i</sup> —Mg1—N1	163.97 (7)	C4—C5—C5 <sup>i</sup>	119.50 (13)
O1W—Mg1—O1W <sup>i</sup>	104.01 (11)	C4—C6—H6A	119.3
O1W—Mg1—O2W	88.36 (6)	C5—N1—Mg1	115.34 (12)
O1W <sup>i</sup> —Mg1—O2W	89.41 (6)	C5—C4—C6	119.2 (2)
O2W <sup>i</sup> —Mg1—N1	96.82 (6)	C6 <sup>i</sup> —C6—C4	121.32 (14)
O2W—Mg1—N1	86.08 (6)	C6 <sup>i</sup> —C6—H6A	119.3
O2W <sup>i</sup> —Mg1—O2W	176.38 (9)	C7—C8—C11	119.45 (15)
Mg1—O1W—H1W1	120.6 (15)	C7—C12—H12A	118.9
Mg1—O1W—H1W2	118.1 (15)	C7—C13—C14	113.25 (16)
Mg1—O2W—H2W1	117.2 (14)	C7—C13—H13A	108.9
Mg1—O2W—H2W2	131.6 (14)	C7—C13—H13B	108.9
N1—C1—C2	122.7 (2)	C8—C7—C12	116.1 (2)
N1—C1—H1A	118.7	C8—C7—C13	121.84 (19)
N1—C5—C4	122.90 (18)	C8—C9—H9A	121.0
N1—C5—C5 <sup>i</sup>	117.60 (10)	C9—C8—C7	123.2 (2)
O1—C14—O2	124.69 (18)	C9—C8—C11	117.34 (17)
O1—C14—C13	117.88 (18)	C9—C10—C12	118.1 (2)
O2—C14—C13	117.42 (18)	C10—C9—C8	118.0 (2)
O1W—Mg1—N1 <sup>i</sup>	163.97 (7)	C10—C9—H9A	121.0
O1W <sup>i</sup> —Mg1—N1 <sup>i</sup>	91.24 (6)	C10—C11—C12	119.1 (2)
O1W—Mg1—O2W <sup>i</sup>	89.41 (6)	C10—C11—H11A	120.5
O1W <sup>i</sup> —Mg1—O2W <sup>i</sup>	88.36 (6)	C11—C10—C9	121.4 (2)
O2W <sup>i</sup> —Mg1—N1 <sup>i</sup>	86.08 (6)	C11—C10—C12	120.5 (2)
O2W—Mg1—N1 <sup>i</sup>	96.82 (6)	C11—C12—C7	122.2 (2)
C1—N1—C5	117.67 (17)	C11—C12—H12A	118.9
C1—N1—Mg1	126.82 (14)	C12—C7—C13	122.1 (2)
C1—C2—H2A	120.1	C12—C11—H11A	120.5
C2—C1—H1A	118.7	C14—C13—H13A	108.9
C2—C3—C4	119.7 (2)	C14—C13—H13B	108.9
C2—C3—H3A	120.2	H1W1—O1W—H1W2	108.5 (14)
C3—C2—C1	119.8 (2)	H2W1—O2W—H2W2	110.2 (14)
C3—C2—H2A	120.1	H13A—C13—H13B	107.7
C3—C4—C5	117.2 (2)		
Mg1—N1—C1—C2	-176.53 (18)	C5—N1—C1—C2	-1.5 (3)

## supplementary materials

Mg1—N1—C5—C4	178.64 (15)	C5—C4—C6—C6 <sup>i</sup>	0.6 (5)
Mg1—N1—C5—C5 <sup>i</sup>	-1.9 (3)	C6—C4—C5—N1	178.3 (2)
N1 <sup>i</sup> —Mg1—N1—C1	175.8 (2)	C6—C4—C5—C5 <sup>i</sup>	-1.2 (3)
N1 <sup>i</sup> —Mg1—N1—C5	0.66 (9)	C7—C8—C9—C10	0.5 (3)
N1—C1—C2—C3	-0.6 (4)	C7—C13—C14—O1	-35.1 (3)
O1W—Mg1—N1—C1	-10.68 (18)	C7—C13—C14—O2	145.54 (19)
O1W <sup>i</sup> —Mg1—N1—C1	151.6 (2)	C8—C7—C12—C11	0.6 (3)
O1W—Mg1—N1—C5	174.14 (13)	C8—C7—C13—C14	-74.1 (2)
O1W <sup>i</sup> —Mg1—N1—C5	-23.6 (3)	C8—C9—C10—C11	-0.2 (3)
O2W <sup>i</sup> —Mg1—N1—C1	-100.24 (18)	C8—C9—C10—C12	-179.92 (15)
O2W—Mg1—N1—C1	77.59 (18)	C9—C10—C11—C12	0.2 (4)
O2W <sup>i</sup> —Mg1—N1—C5	84.59 (13)	C10—C11—C12—C7	-0.4 (4)
O2W—Mg1—N1—C5	-97.59 (13)	C11—C8—C9—C10	179.64 (16)
C1—N1—C5—C4	3.0 (3)	C12—C7—C8—C9	-0.7 (3)
C1—N1—C5—C5 <sup>i</sup>	-177.5 (2)	C12—C10—C11—C12	179.89 (17)
C1—C2—C3—C4	1.2 (4)	C13—C7—C8—C9	179.01 (18)
C2—C3—C4—C5	0.2 (4)	C12—C7—C8—C11	-179.82 (15)
C2—C3—C4—C6	179.5 (2)	C12—C7—C13—C14	105.5 (2)
C3—C4—C5—N1	-2.4 (3)	C13—C7—C8—C11	-0.2 (3)
C3—C4—C6—C6 <sup>i</sup>	-178.7 (3)	C13—C7—C12—C11	-179.0 (2)
C3—C4—C5—C5 <sup>i</sup>	178.1 (2)		

Symmetry codes: (i)  $-x+1, y, -z+1/2$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2W—H2W1 $\cdots$ O1 <sup>ii</sup>	0.849 (9)	1.876 (9)	2.725 (2)	178 (2)
O2W—H2W2 $\cdots$ O1 <sup>i</sup>	0.841 (9)	1.96 (1)	2.772 (2)	161 (2)
O1W—H1W1 $\cdots$ O2 <sup>iii</sup>	0.851 (9)	1.91 (1)	2.728 (2)	162 (2)
O1W—H1W2 $\cdots$ O2 <sup>ii</sup>	0.849 (9)	1.84 (1)	2.685 (2)	171 (2)

Symmetry codes: (ii)  $x, y, z-1$ ; (i)  $-x+1, y, -z+1/2$ ; (iii)  $x, -y+1, z-1/2$ .



Fig. 1




Fig. 2

