

## N-[**(5-Chloro-3-methyl-1-phenyl-1H-pyrazol-4-yl)carbonyl]-N'-(4-hydroxyphenyl)thiourea**

Haitang Du,<sup>a\*</sup> Haijun Du,<sup>b</sup> Ying An<sup>c</sup> and Shengnan Li<sup>c</sup>

<sup>a</sup>Department of Biology and Environment Technology, Guiyang College, Guiyang 550005, People's Republic of China, <sup>b</sup>School of Chemistry and Environment Science, Guizhou University for Nationalities, Guiyang 550025, People's Republic of China, and <sup>c</sup>Department of Chemistry, College of Science, Tianjin University, Tianjin 300072, People's Republic of China

Correspondence e-mail: haitangdu@gz139.com.cn


Received 17 June 2008; accepted 26 June 2008

Key indicators: single-crystal X-ray study;  $T = 294\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.046;  $wR$  factor = 0.123; data-to-parameter ratio = 12.7.

In the title compound,  $\text{C}_{18}\text{H}_{15}\text{ClN}_4\text{O}_2\text{S}$ , the pyrazole ring makes dihedral angles of  $67.4(1)$  and  $12.5(1)^\circ$  with the phenyl and 4-hydroxyphenyl groups, respectively; the two benzene rings are twisted by  $60.1(1)^\circ$  with respect to each other. The thiourea NH groups are involved in  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{Cl}$  intramolecular hydrogen bonds. A hydrogen bond between the phenolic OH group and the pyrazole N atom connects molecules into a one-dimensional polymeric structure.

## Related literature

For related literature, see: Du *et al.* (2007); Saeed & Flörke (2007); Wang *et al.* (2007).



## Experimental

### Crystal data

$\text{C}_{18}\text{H}_{15}\text{ClN}_4\text{O}_2\text{S}$   
 $M_r = 386.85$   
Triclinic,  $P\bar{1}$   
 $a = 8.572(2)\text{ \AA}$   
 $b = 10.429(2)\text{ \AA}$   
 $c = 11.170(2)\text{ \AA}$   
 $\alpha = 99.936(4)^\circ$   
 $\beta = 105.817(4)^\circ$   
 $\gamma = 106.042(4)^\circ$   
 $V = 889.5(3)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.35\text{ mm}^{-1}$   
 $T = 294(2)\text{ K}$   
 $0.26 \times 0.24 \times 0.20\text{ mm}$

### Data collection

Bruker SMART 1K CCD  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $R_{\min} = 0.914$ ,  $T_{\max} = 0.933$   
4615 measured reflections  
3118 independent reflections  
2160 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.122$   
 $S = 1.04$   
3118 reflections  
245 parameters  
2 restraints  
H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2 $\cdots$ N2 <sup>i</sup>	0.82	2.15	2.938 (3)	162
N3—H3A $\cdots$ Cl1	0.891 (10)	2.422 (19)	3.168 (2)	141 (2)
N4—H4A $\cdots$ O1	0.901 (10)	1.92 (2)	2.661 (3)	139 (2)

Symmetry code: (i)  $x, y - 1, z - 1$ .

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Guiyang College (project No. 2008012) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2153).

## References

- Bruker (1997). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Du, H.-T., Lu, M., Zhou, W.-Y. & Sun, L.-L. (2007). *Acta Cryst. E63*, o4287.  
Saeed, A. & Flörke, U. (2007). *Acta Cryst. E63*, o3695.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.  
Wang, J., Tian, L. & Liu, S.-Y. (2007). *Acta Cryst. E63*, o3667.

## **supplementary materials**

Acta Cryst. (2008). E64, o1404 [doi:10.1107/S1600536808019417]

## N-[**(5-Chloro-3-methyl-1-phenyl-1H-pyrazol-4-yl)carbonyl]-N'-(4-hydroxyphenyl)thiourea**

**H. Du, H. Du, Y. An and S. Li**

### Comment

The title compound is similar to the previously reported N-(5-chloro-3-methyl-1-phenylpyrazole-4-ylcarbonyl)-N'-(4-methphenyl)thiourea (Du *et al.*, 2007). The molecular structure of the title compound and the atom-numbering scheme are shown in Fig.1. The pyrazole ring makes dihedral angles of 67.4 (1) and 12.5 (1) $^{\circ}$ , with the C1—C6 and C13—C18 rings, respectively. These two six-membered rings are twisted by 60.1 (1) $^{\circ}$  with respect to each other. This geometry is stabilized by intramolecular N4-H4A  $\cdots$  O1 and N3-H3A  $\cdots$  Cl hydrogen bonds (Fig.1, Table 1). In the crystal structure, molecules are linked by intermolecular N-H  $\cdots$  O hydrogen bonds to form a one-dimensional polymeric structure (Fig.2). All bond lengths and angles are in the normal range (Du *et al.*, 2007; Saeed & Flörke, 2007; Wang *et al.*, 2007).

### Experimental

Powdered ammonium thiocyanate (15 mmol), 5-chloro-3-methyl-1-phenyl-pyrazole-4-carbonyl chloride (10 mmol), PEG-400 (0.5 mL) and acetone (25 mL) were placed in a dried round-bottom flask and stirred at room temperature for 1 h, then 4-aminophenol (9.5 mmol) was added, and the mixture was stirred for 5 h. The mixture was poured into water (20 mL). The resulting solid was filtered, dried and recrystallized from *N,N*-dimethylformamide/ethanol to give the title compound. Single crystals were obtained by slow evaporation of a solution in *N,N*-dimethylformamide/ethanol (1:1, v/v).

### Refinement

H atoms bonded to N atoms were located in a difference Fourier map and refined with distance restraints (N—H = 0.89 Å) and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ . Other H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93–0.96 Å and O—H = 0.82 Å;  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{carrier atom})$  where  $x = 1.5$  for methyl groups and 1.2 for the remaining atoms.

### Figures



Fig. 1. The molecular structure of the title compound with the atom numbering scheme, showing displacement ellipsoids at the 50% probability level.



Fig. 2. The polymeric structure via O-H  $\cdots$  N hydrogen bonds. Hydrogen bonds are shown with dashed lines.

# supplementary materials

---

## *N*-(5-Chloro-3-methyl-1-phenyl-1*H*-pyrazol-4-yl)carbonyl]- *N'*-(4-hydroxyphenyl)thiourea

### Crystal data

C <sub>18</sub> H <sub>15</sub> ClN <sub>4</sub> O <sub>2</sub> S	Z = 2
$M_r = 386.85$	$F_{000} = 400$
Triclinic, $P\bar{1}$	$D_x = 1.444 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Melting point: 456 K
$a = 8.572 (2) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.429 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$c = 11.170 (2) \text{ \AA}$	Cell parameters from 1626 reflections
$\alpha = 99.936 (4)^\circ$	$\theta = 2.6\text{--}25.0^\circ$
$\beta = 105.817 (4)^\circ$	$\mu = 0.35 \text{ mm}^{-1}$
$\gamma = 106.042 (4)^\circ$	$T = 294 (2) \text{ K}$
$V = 889.5 (3) \text{ \AA}^3$	Prism, colorless
	0.26 × 0.24 × 0.20 mm

### Data collection

Bruker SMART 1K CCD diffractometer	3118 independent reflections
Radiation source: fine-focus sealed tube	2160 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 294(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10\text{--}7$
$T_{\text{min}} = 0.914$ , $T_{\text{max}} = 0.933$	$k = -12\text{--}12$
4615 measured reflections	$l = -10\text{--}13$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.123$	$w = 1/[\sigma^2(F_o^2) + (0.0601P)^2 + 0.0911P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3118 reflections	$\Delta\rho_{\text{max}} = 0.20 \text{ e \AA}^{-3}$
245 parameters	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$
2 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.12809 (11)	0.51284 (7)	0.87995 (7)	0.0638 (3)
S1	0.24176 (14)	0.24348 (8)	0.57180 (8)	0.0743 (3)
O1	0.2267 (3)	0.66532 (18)	0.52727 (17)	0.0588 (6)
O2	0.4029 (3)	0.19520 (19)	0.00036 (17)	0.0553 (5)
H2	0.3689	0.1104	-0.0168	0.083*
N1	0.1986 (3)	0.78380 (19)	0.93012 (18)	0.0391 (5)
N2	0.2336 (3)	0.8960 (2)	0.88065 (18)	0.0419 (5)
N3	0.2115 (3)	0.4873 (2)	0.6192 (2)	0.0462 (6)
N4	0.2649 (3)	0.4289 (2)	0.43005 (19)	0.0441 (5)
C1	0.1829 (3)	0.7990 (2)	1.0561 (2)	0.0380 (6)
C2	0.3043 (4)	0.7804 (3)	1.1543 (2)	0.0464 (7)
H2A	0.3950	0.7556	1.1394	0.056*
C3	0.2892 (4)	0.7994 (3)	1.2762 (3)	0.0533 (7)
H3	0.3708	0.7883	1.3442	0.064*
C4	0.1535 (4)	0.8345 (3)	1.2966 (3)	0.0582 (8)
H4	0.1431	0.8462	1.3782	0.070*
C5	0.0335 (4)	0.8525 (3)	1.1974 (3)	0.0622 (8)
H5	-0.0579	0.8763	1.2119	0.075*
C6	0.0477 (4)	0.8354 (3)	1.0758 (3)	0.0506 (7)
H6	-0.0329	0.8483	1.0084	0.061*
C7	0.2683 (4)	0.9426 (3)	0.6808 (2)	0.0564 (8)
H7A	0.2970	1.0362	0.7298	0.085*
H7B	0.1667	0.9190	0.6068	0.085*
H7C	0.3624	0.9343	0.6529	0.085*
C8	0.2354 (3)	0.8466 (2)	0.7635 (2)	0.0384 (6)
C9	0.2041 (3)	0.7020 (2)	0.7349 (2)	0.0362 (6)
C10	0.1803 (3)	0.6678 (2)	0.8446 (2)	0.0391 (6)
C11	0.2136 (3)	0.6184 (2)	0.6186 (2)	0.0388 (6)
C12	0.2401 (3)	0.3889 (3)	0.5323 (2)	0.0429 (6)
C13	0.3013 (3)	0.3615 (2)	0.3244 (2)	0.0395 (6)
C14	0.3412 (4)	0.4386 (3)	0.2413 (2)	0.0490 (7)
H14	0.3439	0.5298	0.2574	0.059*
C15	0.3771 (4)	0.3835 (3)	0.1354 (3)	0.0522 (7)

## supplementary materials

---

H15	0.4046	0.4376	0.0810	0.063*
C16	0.3726 (3)	0.2478 (3)	0.1094 (2)	0.0415 (6)
C17	0.3379 (4)	0.1723 (3)	0.1931 (2)	0.0504 (7)
H17	0.3389	0.0821	0.1780	0.061*
C18	0.3012 (4)	0.2268 (3)	0.2999 (3)	0.0535 (8)
H18	0.2765	0.1730	0.3551	0.064*
H3A	0.205 (4)	0.459 (3)	0.6890 (18)	0.066 (9)*
H4A	0.260 (4)	0.5134 (15)	0.426 (3)	0.064 (9)*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.1074 (7)	0.0351 (4)	0.0593 (5)	0.0189 (4)	0.0486 (4)	0.0147 (3)
S1	0.1377 (9)	0.0495 (5)	0.0739 (6)	0.0502 (5)	0.0681 (6)	0.0290 (4)
O1	0.1072 (17)	0.0502 (11)	0.0377 (10)	0.0421 (12)	0.0355 (11)	0.0157 (9)
O2	0.0702 (14)	0.0514 (11)	0.0490 (11)	0.0212 (11)	0.0335 (10)	0.0029 (9)
N1	0.0546 (14)	0.0326 (11)	0.0314 (11)	0.0138 (10)	0.0196 (10)	0.0054 (9)
N2	0.0624 (15)	0.0350 (11)	0.0338 (11)	0.0189 (10)	0.0223 (10)	0.0096 (9)
N3	0.0740 (17)	0.0357 (12)	0.0405 (13)	0.0245 (11)	0.0317 (12)	0.0098 (10)
N4	0.0695 (16)	0.0334 (12)	0.0375 (12)	0.0248 (11)	0.0240 (11)	0.0082 (10)
C1	0.0495 (17)	0.0319 (13)	0.0332 (13)	0.0127 (12)	0.0187 (12)	0.0046 (10)
C2	0.0603 (19)	0.0426 (15)	0.0458 (16)	0.0244 (14)	0.0247 (14)	0.0135 (12)
C3	0.076 (2)	0.0499 (16)	0.0380 (15)	0.0244 (15)	0.0209 (14)	0.0154 (12)
C4	0.085 (2)	0.0532 (17)	0.0442 (17)	0.0223 (17)	0.0366 (17)	0.0110 (14)
C5	0.068 (2)	0.075 (2)	0.0599 (19)	0.0313 (17)	0.0414 (17)	0.0162 (16)
C6	0.0510 (18)	0.0583 (18)	0.0478 (16)	0.0229 (14)	0.0205 (13)	0.0142 (13)
C7	0.095 (2)	0.0413 (15)	0.0418 (15)	0.0289 (16)	0.0298 (16)	0.0146 (12)
C8	0.0502 (16)	0.0359 (13)	0.0311 (13)	0.0182 (12)	0.0142 (11)	0.0080 (11)
C9	0.0457 (16)	0.0332 (13)	0.0305 (12)	0.0154 (11)	0.0135 (11)	0.0062 (10)
C10	0.0493 (16)	0.0330 (13)	0.0366 (14)	0.0150 (12)	0.0175 (12)	0.0067 (11)
C11	0.0460 (16)	0.0373 (14)	0.0324 (13)	0.0164 (12)	0.0124 (11)	0.0054 (11)
C12	0.0509 (17)	0.0373 (14)	0.0398 (14)	0.0156 (12)	0.0183 (12)	0.0032 (11)
C13	0.0491 (16)	0.0352 (13)	0.0339 (13)	0.0164 (12)	0.0154 (12)	0.0028 (11)
C14	0.071 (2)	0.0328 (13)	0.0509 (16)	0.0221 (14)	0.0291 (15)	0.0094 (12)
C15	0.071 (2)	0.0446 (16)	0.0491 (16)	0.0214 (14)	0.0310 (15)	0.0135 (13)
C16	0.0436 (16)	0.0398 (14)	0.0379 (14)	0.0141 (12)	0.0155 (12)	-0.0001 (11)
C17	0.076 (2)	0.0356 (14)	0.0448 (15)	0.0237 (14)	0.0276 (14)	0.0049 (12)
C18	0.088 (2)	0.0389 (15)	0.0445 (16)	0.0248 (15)	0.0351 (15)	0.0126 (12)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cl1—C10	1.698 (2)	C4—H4	0.9300
S1—C12	1.654 (3)	C5—C6	1.382 (4)
O1—C11	1.224 (3)	C5—H5	0.9300
O2—C16	1.369 (3)	C6—H6	0.9300
O2—H2	0.8200	C7—C8	1.496 (3)
N1—C10	1.347 (3)	C7—H7A	0.9600
N1—N2	1.373 (3)	C7—H7B	0.9600
N1—C1	1.436 (3)	C7—H7C	0.9600

N2—C8	1.327 (3)	C8—C9	1.419 (3)
N3—C11	1.363 (3)	C9—C10	1.384 (3)
N3—C12	1.407 (3)	C9—C11	1.470 (3)
N3—H3A	0.891 (10)	C13—C14	1.380 (3)
N4—C12	1.331 (3)	C13—C18	1.384 (3)
N4—C13	1.421 (3)	C14—C15	1.374 (3)
N4—H4A	0.901 (10)	C14—H14	0.9300
C1—C6	1.374 (4)	C15—C16	1.382 (3)
C1—C2	1.374 (4)	C15—H15	0.9300
C2—C3	1.389 (4)	C16—C17	1.363 (4)
C2—H2A	0.9300	C17—C18	1.384 (3)
C3—C4	1.375 (4)	C17—H17	0.9300
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.371 (4)		
C16—O2—H2	109.5	H7B—C7—H7C	109.5
C10—N1—N2	110.66 (18)	N2—C8—C9	111.7 (2)
C10—N1—C1	128.6 (2)	N2—C8—C7	119.6 (2)
N2—N1—C1	120.68 (17)	C9—C8—C7	128.7 (2)
C8—N2—N1	105.50 (18)	C10—C9—C8	103.6 (2)
C11—N3—C12	130.5 (2)	C10—C9—C11	130.9 (2)
C11—N3—H3A	118.6 (19)	C8—C9—C11	125.2 (2)
C12—N3—H3A	110.4 (19)	N1—C10—C9	108.6 (2)
C12—N4—C13	130.9 (2)	N1—C10—C11	120.04 (18)
C12—N4—H4A	115.9 (18)	C9—C10—C11	131.33 (19)
C13—N4—H4A	113.2 (18)	O1—C11—N3	121.8 (2)
C6—C1—C2	121.7 (2)	O1—C11—C9	121.6 (2)
C6—C1—N1	118.6 (2)	N3—C11—C9	116.6 (2)
C2—C1—N1	119.7 (2)	N4—C12—N3	114.1 (2)
C1—C2—C3	118.7 (3)	N4—C12—S1	129.42 (19)
C1—C2—H2A	120.6	N3—C12—S1	116.46 (19)
C3—C2—H2A	120.6	C14—C13—C18	118.3 (2)
C4—C3—C2	120.1 (3)	C14—C13—N4	116.5 (2)
C4—C3—H3	120.0	C18—C13—N4	125.3 (2)
C2—C3—H3	120.0	C15—C14—C13	121.4 (2)
C5—C4—C3	120.4 (3)	C15—C14—H14	119.3
C5—C4—H4	119.8	C13—C14—H14	119.3
C3—C4—H4	119.8	C14—C15—C16	120.1 (2)
C4—C5—C6	120.3 (3)	C14—C15—H15	119.9
C4—C5—H5	119.9	C16—C15—H15	119.9
C6—C5—H5	119.9	C17—C16—O2	122.9 (2)
C1—C6—C5	118.9 (3)	C17—C16—C15	118.7 (2)
C1—C6—H6	120.5	O2—C16—C15	118.3 (2)
C5—C6—H6	120.5	C16—C17—C18	121.6 (2)
C8—C7—H7A	109.5	C16—C17—H17	119.2
C8—C7—H7B	109.5	C18—C17—H17	119.2
H7A—C7—H7B	109.5	C13—C18—C17	119.9 (2)
C8—C7—H7C	109.5	C13—C18—H18	120.1
H7A—C7—H7C	109.5	C17—C18—H18	120.1

## supplementary materials

---


C10—N1—N2—C8	0.3 (3)	C11—C9—C10—N1	173.8 (3)
C1—N1—N2—C8	-178.5 (2)	C8—C9—C10—Cl1	176.7 (2)
C10—N1—C1—C6	-112.6 (3)	C11—C9—C10—Cl1	-8.8 (4)
N2—N1—C1—C6	66.0 (3)	C12—N3—C11—O1	7.4 (4)
C10—N1—C1—C2	69.0 (3)	C12—N3—C11—C9	-171.0 (3)
N2—N1—C1—C2	-112.5 (3)	C10—C9—C11—O1	176.2 (3)
C6—C1—C2—C3	-0.1 (4)	C8—C9—C11—O1	-10.4 (4)
N1—C1—C2—C3	178.2 (2)	C10—C9—C11—N3	-5.4 (4)
C1—C2—C3—C4	0.7 (4)	C8—C9—C11—N3	168.0 (2)
C2—C3—C4—C5	-0.7 (4)	C13—N4—C12—N3	178.3 (2)
C3—C4—C5—C6	0.0 (5)	C13—N4—C12—S1	-0.5 (4)
C2—C1—C6—C5	-0.5 (4)	C11—N3—C12—N4	-3.7 (4)
N1—C1—C6—C5	-178.9 (2)	C11—N3—C12—S1	175.3 (2)
C4—C5—C6—C1	0.6 (4)	C12—N4—C13—C14	-172.2 (3)
N1—N2—C8—C9	-0.8 (3)	C12—N4—C13—C18	7.1 (4)
N1—N2—C8—C7	179.3 (2)	C18—C13—C14—C15	1.2 (4)
N2—C8—C9—C10	1.0 (3)	N4—C13—C14—C15	-179.4 (3)
C7—C8—C9—C10	-179.2 (3)	C13—C14—C15—C16	0.5 (4)
N2—C8—C9—C11	-174.0 (2)	C14—C15—C16—C17	-2.2 (4)
C7—C8—C9—C11	5.9 (4)	C14—C15—C16—O2	177.6 (3)
N2—N1—C10—C9	0.3 (3)	O2—C16—C17—C18	-177.5 (3)
C1—N1—C10—C9	179.0 (2)	C15—C16—C17—C18	2.4 (4)
N2—N1—C10—Cl1	-177.47 (17)	C14—C13—C18—C17	-1.0 (4)
C1—N1—C10—Cl1	1.2 (4)	N4—C13—C18—C17	179.6 (3)
C8—C9—C10—N1	-0.7 (3)	C16—C17—C18—C13	-0.8 (4)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O2—H2 $\cdots$ N2 <sup>i</sup>	0.82	2.15	2.938 (3)	162
N3—H3A $\cdots$ Cl1	0.891 (10)	2.422 (19)	3.168 (2)	141 (2)
N4—H4A $\cdots$ O1	0.901 (10)	1.92 (2)	2.661 (3)	139 (2)

Symmetry codes: (i)  $x, y-1, z-1$ .

Fig. 1



## **supplementary materials**

---

**Fig. 2**

