

(3*R*,3*aS*,6*aR*)-2,5-Dimethyl-3-(5-phenyl-2-thienyl)perhydropyrrolo[3,4-*d*][1,2]-oxazole-4,6-dioneMustafa Odabaşoğlu,^a Hamdi Özkan,^b Yılmaz Yıldırım^b and Orhan Büyükgüngör^{c*}^aDepartment of Chemistry, Faculty of Arts and Sciences, Mehmet Akif Ersoy University, TR-15030 Burdur, Turkey, ^bDepartment of Chemistry, Faculty of Arts and Sciences, Gazi University, Ankara, Turkey, and ^cDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, TR-55139 Kurupelit Samsun, Turkey
Correspondence e-mail: orhanb@omu.edu.tr


Received 27 June 2008; accepted 1 July 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.032; wR factor = 0.088; data-to-parameter ratio = 8.5.

The crystal structure of the title compound, $\text{C}_{17}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$, exhibits intramolecular $\text{C}-\text{H}\cdots\text{S}$ and intermolecular $\text{C}-\text{H}\cdots\text{S}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, $\text{C}-\text{S}\cdots\text{N}$ [$\text{S}\cdots\text{N} = 3.033$ (2) Å and $\text{C}-\text{S}\cdots\text{N} = 142.76$ (9)°] interactions, and $\text{C}-\text{H}\cdots\pi$ interactions; these interactions generate $S(4)$, $S(6)$ and $R_2^2(14)$ ring motifs. The isoxazole ring adopts an envelope conformation, with the N atom displaced by 0.672 (2) Å from the plane of the other ring atoms. The thiophene ring is oriented with respect to the succinimide and phenyl rings at dihedral angles of 40.03 (12) and 5.21 (13)°, respectively. The dihedral angle between the succinimide and phenyl rings is 39.38 (12)°.

Related literature

For general background, see: Huisgen (1960); Black *et al.* (1975); Richman (2001); De Clercq (2002); Donadas *et al.* (2004); Merino *et al.* (2003); Chiacchio *et al.* (2003); Iannazzo *et al.* (2002). For related literature, see: Heaney *et al.* (2001). For ring motif details, see: Bernstein *et al.* (1995); Etter (1990).

**Experimental***Crystal data* $\text{C}_{17}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$
 $M_r = 328.38$
Orthorhombic, $Pna2_1$
 $a = 12.7768$ (7) Å
 $b = 10.9803$ (6) Å
 $c = 11.1069$ (9) Å
 $V = 1558.22$ (17) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.22$ mm⁻¹
 $T = 296$ K
 $0.65 \times 0.46 \times 0.27$ mm*Data collection*Stoe IPDSII diffractometer
Absorption correction: integration
(*X-RED32*; Stoe & Cie, 2002)
 $T_{\min} = 0.889$, $T_{\max} = 0.935$
6395 measured reflections
3332 independent reflections
2899 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$ *Refinement* $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.099$
 $S = 1.04$
3332 reflections
221 parameters
1 restraint
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.22$ e Å⁻³
 $\Delta\rho_{\min} = -0.22$ e Å⁻³
Absolute structure: Flack (1983), 1451 Friedel pairs
Flack parameter = -0.13 (8)**Table 1**

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the S1/C7–C10 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6–H6 \cdots S1	0.93	2.68	3.097 (3)	108
C8–H8 \cdots S1 ⁱ	0.93	3.00	3.887 (2)	160
C14–H14B \cdots O2 ⁱ	0.96	2.65	3.417 (4)	137
C13–H13 \cdots Cg1 ⁱⁱ	1.01 (3)	2.96 (3)	3.875 (2)	152 (2)

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z$; (ii) $-x, -y, z - \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDSII diffractometer (purchased under grant F.279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2484).

References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Black, D. C., Crozier, R. F. & Davis, V. C. (1975). *Synthesis*, pp. 205–221.
- Chiacchio, U., Corsaro, A., Iannazzo, D., Piperno, A., Pistara, V., Rescifina, A., Romeo, R., Sindona, G. & Romeo, G. (2003). *Tetrahedron Asymmetry*, **14**, 2717–2723.
- De Clercq, E. (2002). *Nat. Rev. Drug Discov.* **1**, 13–26.
- Donadas, H. A., Fishwick, C. W. G., Grigg, R. & Kilner, C. (2004). *Tetrahedron*, **60**, 3473–3485.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

- Heaney, F., Rooney, O., Cunningham, D. & McArdle, P. (2001). *J. Chem. Soc. Perkin Trans. 2*, pp. 373–378.
- Huisgen, R. (1960). *10 Jahre Fonds der Chemischen Industrie Düsseldorf*, p. 73; reprinted in *Naturwissenschaften* (1961), **4**, 63.
- Iannazzo, D., Piperno, A., Pistara, V., Rescifina, A. & Romeo, R. (2002). *Tetrahedron*, **58**, 581–587.
- Merino, P., Tejero, T., Laguna, M., Cerrada, E., Moreno, A. & Lopez, J. A. (2003). *Org. Biomol. Chem.* **1**, 2336–2342.
- Richman, D. D. (2001). *Nature (London)*, **410**, 995–1001.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2008). E64, o1423-o1424 [doi:10.1107/S1600536808020102]

(3*R*,3*aS*,6*aR*)-2,5-Dimethyl-3-(5-phenyl-2-thienyl)perhydropyrrolo[3,4-*d*][1,2]oxazole-4,6-dione

M. Odabasoglu, H. Özkan, Y. Yildirim and O. Büyükgüngör

Comment

A general principle for the synthesis of five-membered rings was introduced in 1960 as 1,3-dipolar cycloaddition and turned out to be remarkably widespread (Huisgen, 1960, 1961). Because of easy 1,3-dipolar cycloaddition reactions to alkenes, alkynes, isocyanates, isothiocyanates, phospharanes, sulphenes and sulphylnl compounds; nitrones are the important intermediates in synthetic organic chemistry (Black *et al.*, 1975). There has been an ever-increasing quest for modified nucleosides, due to their potential applications in antiviral and anticancer therapies (Richman, 2001; De Clercq, 2002; Donadas *et al.*, 2004). In a recent approach to modified nucleosides, the furanose ring has been replaced by other heterocyclic analogs (Merino *et al.*, 2003). Among these N and O containing heterocycles have emerged as important candidates, and have been shown to display useful anticancer and antiviral properties (Chiacchio *et al.*, 2003; Iannazzo *et al.*, 2002). The present work is part of a structural study of compounds of substituted 2,5-dimethyl-4-(thiophen-2-yl)-tetrahydropyrrolo[3,4-*c*]pyrrole-1,3(2*H*,3*aH*)-dione systems with hydrogen-bond donors, and we report herein the crystal structure of the title compound, (I).

The overall view and atom-labelling of the molecule of (I) are displayed in Fig. 1. The thiophene ring is oriented with respect to succinimide and phenyl rings at dihedral angles of 40.03 (12)° and 5.21 (13)°, respectively. The dihedral angle between succinimide and phenyl rings is 39.38 (12)°. The isoxazole ring has envelope conformation, with N1 atom displaced by -0.672 (2) Å from the plane of the other ring atoms.

The hydrogen-bonding parameters are given in Table 1 and the packing arrangements of the molecules are illustrated in Figs. 2 and 3. Compound is stabilized by intramolecular C—H···S hydrogen bond and S···N heteroatom interactions [in C1—S1···N; S···N = 3.033 (2) Å, C1—S1···N = 142.76 (9) °], which form S(4) and S(6) motifs, and intermolecular C—H···S and C—H···O hydrogen bonds and C—H···π interactions. As shown in Fig. 2 the structure of the compound is made up of C8—H8···S1 and C14—H14B···O2 H-bonded polymeric bands of [C₁₂H₁₄N₂O₃S] molecules, which are nearly elongated along [100]. These polymeric chains are linked to each other and generate R₂²(14) ring motifs (Bernstein *et al.*, 1995; Etter, 1990). The crystal packing is also stabilized by C13—H13···Cg1 interactions (Fig. 3, Table 1).

Experimental

N-Methyl-*C*-(5-Phenylthiophen) nitron, (II), was prepared from 5-Phenylthiophenecarbaldehyde, *N*-methyl-hydroxylamine hydrochloride and sodium carbonate in CH₂Cl₂ according to the literature method (Heaney *et al.*, 2001). For the preparation of the title compound, (II) (657 mg, 3 mmol) and *N*-methylmaleimide (370 mg, 3.3 mmol) were dissolved in benzene (50 ml). The reaction mixture was refluxed for 12 h, and monitored by TLC. After evaporation of the solvent, the reaction mixture was separated by column chromatography, using the mixture of petroleum ether/ethyl acetate (2:1) as the eluant. The *cis*-isomer, (I), was recrystallized from CHCl₃/n-hexane (mp: 452,9-454,4 K).

Refinement

1451 Friedel pairs were averaged before the final refinement, but the absolute configuration could not be determined unambiguously, although Si atom is present. The methine H atoms, H11, H12 and H13, were located in difference syntheses and refined isotropically [C-H = 0.95 (3)-1.04 (3) Å; $U_{\text{iso}}(\text{H}) = 0.035$ (6)-0.071 (10) Å²]. The remaining H atoms were positioned geometrically, with C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for C14 methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 20% probability level. Hydrogen bonds are shown as dashed lines.




Fig. 2. A partial packing diagram of (I), showing the formation of S(4), S(6) and $R_2^2(14)$ ring motifs along [100] [symmetry code: (i) $x - 1/2, y, z$]. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.




Fig. 3. A packing diagram of (I) [symmetry code: (i) $1 - x, 1 - y, -z$], where Cg1 is the centroid of ring (S1/C7-C10). Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

(3*R*,3*aS*,6*aR*)-2,5-Dimethyl-3-(5-phenyl-2-thienyl)perhydropyrrolo[3,4-*d*][1,2]oxazole-4,6-dione

Crystal data

$C_{17}H_{16}N_2O_3S$

$M_r = 328.38$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 12.7768$ (7) Å

$b = 10.9803$ (6) Å

$c = 11.1069$ (9) Å

$V = 1558.22$ (17) Å³

$Z = 4$

$F_{000} = 688$

$D_x = 1.400$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 6395 reflections

$\theta = 1.8$ –28.0°

$\mu = 0.22$ mm⁻¹

$T = 296$ K

Prism, colorless

$0.65 \times 0.46 \times 0.27$ mm

Data collection

Stoe IPDSII diffractometer	3332 independent reflections
Monochromator: plane graphite	2899 reflections with $I > 2\sigma(I)$
Detector resolution: 6.67 pixels mm^{-1}	$R_{\text{int}} = 0.051$
$T = 296$ K	$\theta_{\text{max}} = 27.5^\circ$
ω scan rotation method	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: integration (X-RED32; Stoe & Cie, 2002)	$h = -13 \rightarrow 16$
$T_{\text{min}} = 0.889$, $T_{\text{max}} = 0.935$	$k = -13 \rightarrow 14$
6395 measured reflections	$l = -12 \rightarrow 14$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.0659P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.099$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.22 \text{ e } \text{\AA}^{-3}$
3332 reflections	$\Delta\rho_{\text{min}} = -0.22 \text{ e } \text{\AA}^{-3}$
221 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
1 restraint	Extinction coefficient: 0.0122 (17)
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983)
Secondary atom site location: difference Fourier map	Flack parameter: -0.13 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.64659 (16)	0.19809 (17)	0.66560 (19)	0.0427 (4)

supplementary materials

C2	0.7443 (2)	0.1782 (2)	0.7188 (2)	0.0585 (6)
H2	0.8048	0.1938	0.6749	0.070*
C3	0.7523 (2)	0.1360 (3)	0.8346 (3)	0.0701 (7)
H3	0.8181	0.1227	0.8680	0.084*
C4	0.6641 (3)	0.1132 (3)	0.9022 (3)	0.0721 (8)
H4	0.6696	0.0857	0.9811	0.087*
C5	0.5678 (2)	0.1320 (3)	0.8504 (2)	0.0731 (8)
H5	0.5075	0.1163	0.8948	0.088*
C6	0.55911 (19)	0.1738 (2)	0.7341 (2)	0.0593 (6)
H6	0.4930	0.1857	0.7011	0.071*
C7	0.63920 (16)	0.24233 (18)	0.5415 (2)	0.0411 (4)
C8	0.71619 (15)	0.2782 (2)	0.4653 (3)	0.0530 (5)
H8	0.7866	0.2789	0.4864	0.064*
C9	0.67952 (15)	0.3144 (2)	0.3508 (2)	0.0511 (5)
H9	0.7236	0.3398	0.2891	0.061*
C10	0.57421 (14)	0.30882 (18)	0.33939 (19)	0.0409 (4)
C11	0.50963 (15)	0.3425 (2)	0.23306 (19)	0.0432 (4)
C12	0.46103 (16)	0.4708 (2)	0.22929 (19)	0.0458 (4)
C13	0.36143 (17)	0.4492 (2)	0.15649 (19)	0.0473 (5)
C14	0.4393 (2)	0.1435 (2)	0.1799 (2)	0.0589 (6)
H14A	0.3761	0.0963	0.1764	0.088*
H14B	0.4884	0.1044	0.2327	0.088*
H14C	0.4690	0.1498	0.1007	0.088*
C15	0.42400 (17)	0.51840 (19)	0.3496 (2)	0.0490 (5)
C16	0.27329 (17)	0.4855 (2)	0.2411 (2)	0.0506 (5)
C17	0.2543 (2)	0.5584 (3)	0.4516 (3)	0.0716 (8)
H17A	0.2708	0.6414	0.4711	0.086*
H17B	0.2700	0.5072	0.5192	0.086*
H17C	0.1812	0.5520	0.4326	0.086*
N1	0.41573 (13)	0.26461 (15)	0.22512 (15)	0.0419 (4)
N2	0.31576 (15)	0.52026 (17)	0.34890 (18)	0.0504 (4)
O1	0.47725 (15)	0.55117 (17)	0.43306 (17)	0.0690 (5)
O2	0.18093 (14)	0.48473 (19)	0.2184 (2)	0.0786 (6)
O3	0.35719 (12)	0.32333 (14)	0.12785 (14)	0.0506 (4)
S1	0.51891 (3)	0.25445 (5)	0.47066 (6)	0.04736 (14)
H11	0.5520 (18)	0.3385 (19)	0.161 (2)	0.038 (5)*
H12	0.511 (2)	0.535 (3)	0.191 (3)	0.068 (8)*
H13	0.356 (2)	0.494 (3)	0.079 (3)	0.064 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0437 (10)	0.0377 (9)	0.0468 (11)	0.0072 (8)	-0.0082 (8)	-0.0047 (8)
C2	0.0489 (11)	0.0677 (15)	0.0590 (14)	0.0064 (10)	-0.0144 (11)	0.0013 (11)
C3	0.0682 (16)	0.0798 (17)	0.0622 (16)	0.0142 (13)	-0.0284 (14)	-0.0017 (13)
C4	0.085 (2)	0.0832 (17)	0.0478 (14)	0.0131 (15)	-0.0162 (13)	0.0040 (12)
C5	0.0745 (18)	0.096 (2)	0.0489 (14)	0.0006 (14)	0.0033 (14)	0.0095 (13)
C6	0.0491 (13)	0.0780 (16)	0.0508 (13)	0.0091 (10)	-0.0068 (10)	0.0064 (11)

C7	0.0322 (9)	0.0409 (9)	0.0501 (12)	0.0039 (8)	-0.0056 (8)	-0.0030 (7)
C8	0.0282 (8)	0.0666 (13)	0.0643 (13)	0.0027 (8)	-0.0018 (11)	0.0054 (14)
C9	0.0299 (9)	0.0670 (13)	0.0564 (12)	0.0011 (9)	0.0055 (9)	0.0079 (10)
C10	0.0305 (8)	0.0497 (10)	0.0426 (10)	0.0005 (7)	0.0039 (8)	0.0012 (8)
C11	0.0339 (9)	0.0595 (11)	0.0361 (9)	-0.0023 (8)	0.0046 (8)	0.0052 (8)
C12	0.0392 (9)	0.0555 (11)	0.0428 (10)	-0.0083 (9)	-0.0024 (8)	0.0111 (9)
C13	0.0448 (10)	0.0572 (11)	0.0398 (11)	-0.0034 (9)	-0.0074 (8)	0.0106 (8)
C14	0.0554 (13)	0.0613 (13)	0.0600 (14)	0.0043 (10)	-0.0034 (11)	-0.0127 (10)
C15	0.0496 (11)	0.0466 (10)	0.0507 (11)	0.0009 (9)	-0.0084 (10)	0.0050 (8)
C16	0.0437 (12)	0.0522 (11)	0.0559 (14)	-0.0022 (8)	-0.0037 (9)	0.0045 (9)
C17	0.0770 (17)	0.0676 (15)	0.0704 (19)	0.0072 (12)	0.0215 (14)	-0.0051 (13)
N1	0.0374 (8)	0.0523 (9)	0.0361 (8)	-0.0025 (7)	-0.0026 (6)	-0.0002 (7)
N2	0.0497 (10)	0.0516 (9)	0.0499 (10)	-0.0011 (8)	0.0039 (9)	0.0046 (7)
O1	0.0711 (12)	0.0742 (11)	0.0618 (11)	0.0082 (9)	-0.0242 (9)	-0.0139 (9)
O2	0.0387 (9)	0.0955 (13)	0.1016 (16)	0.0011 (8)	-0.0111 (9)	-0.0080 (12)
O3	0.0506 (9)	0.0618 (9)	0.0394 (7)	-0.0024 (7)	-0.0121 (6)	0.0002 (6)
S1	0.0287 (2)	0.0711 (3)	0.0423 (2)	-0.0033 (2)	-0.0012 (2)	0.0108 (2)

Geometric parameters (Å, °)

C1—C6	1.378 (3)	C11—C12	1.540 (3)
C1—C2	1.398 (3)	C11—H11	0.97 (2)
C1—C7	1.464 (3)	C12—C15	1.511 (3)
C2—C3	1.371 (4)	C12—C13	1.526 (3)
C2—H2	0.9300	C12—H12	1.04 (3)
C3—C4	1.377 (5)	C13—O3	1.419 (3)
C3—H3	0.9300	C13—C16	1.520 (3)
C4—C5	1.374 (4)	C13—H13	1.00 (3)
C4—H4	0.9300	C14—N1	1.453 (3)
C5—C6	1.375 (4)	C14—H14A	0.9600
C5—H5	0.9300	C14—H14B	0.9600
C6—H6	0.9300	C14—H14C	0.9600
C7—C8	1.357 (3)	C15—O1	1.205 (3)
C7—S1	1.732 (2)	C15—N2	1.383 (3)
C8—C9	1.412 (4)	C16—O2	1.207 (3)
C8—H8	0.9300	C16—N2	1.369 (3)
C9—C10	1.353 (3)	C17—N2	1.447 (3)
C9—H9	0.9300	C17—H17A	0.9600
C10—C11	1.487 (3)	C17—H17B	0.9600
C10—S1	1.727 (2)	C17—H17C	0.9600
C11—N1	1.476 (3)	N1—O3	1.464 (2)
C6—C1—C2	117.4 (2)	C15—C12—C11	114.73 (17)
C6—C1—C7	122.11 (19)	C13—C12—C11	102.00 (17)
C2—C1—C7	120.5 (2)	C15—C12—H12	108.7 (15)
C3—C2—C1	121.1 (3)	C13—C12—H12	113.7 (15)
C3—C2—H2	119.5	C11—C12—H12	112.3 (16)
C1—C2—H2	119.5	O3—C13—C16	111.44 (18)
C2—C3—C4	120.8 (2)	O3—C13—C12	107.59 (17)
C2—C3—H3	119.6	C16—C13—C12	104.45 (18)

supplementary materials

C4—C3—H3	119.6	O3—C13—H13	106.8 (16)
C5—C4—C3	118.5 (3)	C16—C13—H13	110.9 (17)
C5—C4—H4	120.7	C12—C13—H13	115.7 (17)
C3—C4—H4	120.7	N1—C14—H14A	109.5
C4—C5—C6	121.0 (3)	N1—C14—H14B	109.5
C4—C5—H5	119.5	H14A—C14—H14B	109.5
C6—C5—H5	119.5	N1—C14—H14C	109.5
C5—C6—C1	121.2 (2)	H14A—C14—H14C	109.5
C5—C6—H6	119.4	H14B—C14—H14C	109.5
C1—C6—H6	119.4	O1—C15—N2	124.4 (2)
C8—C7—C1	129.6 (2)	O1—C15—C12	127.4 (2)
C8—C7—S1	109.71 (17)	N2—C15—C12	108.27 (18)
C1—C7—S1	120.70 (16)	O2—C16—N2	124.9 (2)
C7—C8—C9	113.79 (18)	O2—C16—C13	126.4 (2)
C7—C8—H8	123.1	N2—C16—C13	108.67 (18)
C9—C8—H8	123.1	N2—C17—H17A	109.5
C10—C9—C8	113.7 (2)	N2—C17—H17B	109.5
C10—C9—H9	123.2	H17A—C17—H17B	109.5
C8—C9—H9	123.2	N2—C17—H17C	109.5
C9—C10—C11	127.94 (19)	H17A—C17—H17C	109.5
C9—C10—S1	110.08 (16)	H17B—C17—H17C	109.5
C11—C10—S1	121.97 (13)	C14—N1—O3	104.71 (16)
N1—C11—C10	110.73 (16)	C14—N1—C11	112.49 (17)
N1—C11—C12	101.59 (15)	O3—N1—C11	101.78 (14)
C10—C11—C12	118.22 (18)	C16—N2—C15	113.39 (19)
N1—C11—H11	112.3 (13)	C16—N2—C17	123.7 (2)
C10—C11—H11	109.5 (14)	C15—N2—C17	122.9 (2)
C12—C11—H11	104.2 (13)	C13—O3—N1	104.13 (14)
C15—C12—C13	105.14 (18)	C10—S1—C7	92.71 (10)
C6—C1—C2—C3	0.0 (3)	C13—C12—C15—O1	-177.3 (2)
C7—C1—C2—C3	-179.6 (2)	C11—C12—C15—O1	71.5 (3)
C1—C2—C3—C4	-0.6 (4)	C13—C12—C15—N2	1.9 (2)
C2—C3—C4—C5	0.9 (4)	C11—C12—C15—N2	-109.3 (2)
C3—C4—C5—C6	-0.6 (5)	O3—C13—C16—O2	-65.7 (3)
C4—C5—C6—C1	0.0 (4)	C12—C13—C16—O2	178.4 (2)
C2—C1—C6—C5	0.3 (4)	O3—C13—C16—N2	114.50 (19)
C7—C1—C6—C5	179.9 (2)	C12—C13—C16—N2	-1.4 (2)
C6—C1—C7—C8	175.0 (2)	C10—C11—N1—C14	-76.4 (2)
C2—C1—C7—C8	-5.4 (3)	C12—C11—N1—C14	157.21 (18)
C6—C1—C7—S1	-5.3 (3)	C10—C11—N1—O3	172.10 (16)
C2—C1—C7—S1	174.28 (17)	C12—C11—N1—O3	45.68 (17)
C1—C7—C8—C9	179.4 (2)	O2—C16—N2—C15	-177.0 (2)
S1—C7—C8—C9	-0.4 (3)	C13—C16—N2—C15	2.8 (3)
C7—C8—C9—C10	1.2 (3)	O2—C16—N2—C17	1.5 (4)
C8—C9—C10—C11	178.6 (2)	C13—C16—N2—C17	-178.8 (2)
C8—C9—C10—S1	-1.4 (3)	O1—C15—N2—C16	176.2 (2)
C9—C10—C11—N1	147.6 (2)	C12—C15—N2—C16	-3.0 (3)
S1—C10—C11—N1	-32.4 (2)	O1—C15—N2—C17	-2.2 (4)
C9—C10—C11—C12	-95.9 (3)	C12—C15—N2—C17	178.5 (2)

S1—C10—C11—C12	84.2 (2)	C16—C13—O3—N1	-86.75 (19)
N1—C11—C12—C15	84.5 (2)	C12—C13—O3—N1	27.2 (2)
C10—C11—C12—C15	-36.9 (3)	C14—N1—O3—C13	-163.37 (17)
N1—C11—C12—C13	-28.59 (19)	C11—N1—O3—C13	-46.07 (19)
C10—C11—C12—C13	-149.94 (17)	C9—C10—S1—C7	1.06 (18)
C15—C12—C13—O3	-118.85 (18)	C11—C10—S1—C7	-178.96 (17)
C11—C12—C13—O3	1.2 (2)	C8—C7—S1—C10	-0.40 (17)
C15—C12—C13—C16	-0.3 (2)	C1—C7—S1—C10	179.86 (16)
C11—C12—C13—C16	119.74 (18)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C6—H6 \cdots S1	0.93	2.69	3.100 (2)	108
C8—H8 \cdots S1 ⁱ	0.93	3.00	3.887 (2)	160
C14—H14B \cdots O2 ⁱ	0.96	2.65	3.417 (4)	137
C13—H13 \cdots Cg1 ⁱⁱ	1.00 (3)	2.97 (3)	3.876 (2)	151 (2)

Symmetry codes: (i) $x+1/2, -y+1/2, z$; (ii) $-x, -y, z-1/2$.

Fig. 1




Fig. 2




Fig. 3

