

P,P-Diphenyl-N-(1,1,2,2-tetraphenyl-1λ⁵-diphosphanylidene)phosphinous amide

Diane A. Dickie and Richard A. Kemp*‡

Department of Chemistry and Chemical Biology, MSC03 2060, 1 University of New Mexico, Albuquerque, NM 87131, USA
Correspondence e-mail: rakemp@unm.edu


Received 23 June 2008; accepted 2 July 2008

Key indicators: single-crystal X-ray study; $T = 228$ K; mean $\sigma(C-C) = 0.002$ Å;
 R factor = 0.042; wR factor = 0.118; data-to-parameter ratio = 23.9.

The title compound, $C_{36}H_{30}NP_3$, a structural isomer of tris(diphenylphosphino)amine, was unexpectedly isolated as the sole phosphorus-containing product from the reaction of $Mg[N(PPh₂)₂]₂(THF)₂$ (THF is tetrahydrofuran) with CO_2 . Its identity was confirmed by ³¹P NMR spectroscopy and single-crystal X-ray diffraction. The geometry at the two P(III) atoms is trigonal pyramidal, while the P(V) atom adopts a distorted tetrahedral geometry.

Related literature

For the original synthesis and spectroscopic characterization of the title compound, see: Nöth & Meinel (1967); Meinel & Nöth (1970). For the crystallographic characterization of the structural isomer $N[P(C_6H_5)_2]_3$, see: Ellermann *et al.* (1987). For related literature, see: Bruno *et al.* (2004).



Experimental

Crystal data

$C_{36}H_{30}NP_3$	$V = 2960.0$ (7) Å ³
$M_r = 569.52$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 9.3026$ (13) Å	$\mu = 0.23$ mm ⁻¹
$b = 10.8167$ (15) Å	$T = 228$ (2) K
$c = 29.750$ (4) Å	$0.57 \times 0.51 \times 0.18$ mm
$\beta = 98.589$ (6)°	

Data collection

Bruker SMART CCD area-detector diffractometer	76797 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	11496 independent reflections
$T_{min} = 0.88$, $T_{max} = 0.96$	9167 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	481 parameters
$wR(F^2) = 0.118$	All H-atom parameters refined
$S = 1.05$	$\Delta\rho_{\max} = 0.41$ e Å ⁻³
11496 reflections	$\Delta\rho_{\min} = -0.29$ e Å ⁻³

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *publCIF* (Westrip, 2008).

The authors thank Eileen Duesler (UNM) for the X-ray data collection. Funding was provided by the Natural Sciences and Engineering Research Council of Canada (NSERC PDF to DAD), the National Science Foundation (grant Nos. CHE-0213165 and CHE-0443580) and the Sandia LDRD Program (grant Nos. 105932 and 113486). Sandia is a multiprogram laboratory operated by Sandia Corporation, a Lockheed Martin Company, for the United States Department of Energy under contract No. DE-AC04-94AL85000.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2581).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2003). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruno, I. J., Cole, J. C., Kessler, M., Luo, J., Motherwell, W. D. S., Purkis, L. H., Smith, B. R., Taylor, R., Cooper, R. I., Harris, S. E. & Orpen, A. G. (2004). *J. Chem. Inf. Comput. Sci.* **44**, 2133–2144.
- Ellermann, J., Köck, E., Zimmermann, H. & Gomm, M. (1987). *Acta Cryst. C* **43**, 1795–1798.
- Meinel, L. & Nöth, H. (1970). *Z. Anorg. Allg. Chem.* **373**, 36–47.
- Nöth, H. & Meinel, L. (1967). *Z. Anorg. Allg. Chem.* **349**, 225–240.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2008). *publCIF*. In preparation.

‡ Concurrent address: Advanced Materials Laboratory, Sandia National Laboratories, 1001 University Blvd SE, Albuquerque, NM 87106, USA.

supplementary materials

Acta Cryst. (2008). E64, o1449 [doi:10.1107/S1600536808020412]

P,P-Diphenyl-N-(1,1,2,2-tetraphenyl-1λ⁵-diphosphanylidene)phosphinous amide

D. A. Dickie and R. A. Kemp

Comment

The molecular structure of the title compound, (I), is shown in Fig. 1. It was originally prepared by Nöth and Meinel [Nöth & Meinel (1967); and Meinel & Nöth (1970)] but its crystal structure was not determined at that time. We report herein the isolation of (I) as an unexpected product of the reaction of carbon dioxide with Mg[N(PPh₂)₂](THF)₂. This compound was characterized by ³¹P NMR spectroscopy and single-crystal X-ray diffraction. The P=N double bond measures 1.5690 (10) Å, very close to the average value of similar bonds in a *Mogul* (Bruno *et al.*, 2004) search of the Cambridge structural database (mean P=N 1.573 Å). The P—N single bond of 1.6755 (11) Å is significantly shorter than those in the structural isomer N(PPh₂)₃ (Ellermann, *et al.* 1987) (mean P—N = 1.740 Å), which is not surprising when the different hybridization of nitrogen (*sp*² *versus* *sp*³) is considered. It is also, however, shorter than the average P—N(*sp*²) bond length of 1.706 Å. The geometry at each of the two P(III) atoms is trigonal pyramidal, due to the stereochemically active lone pair on each of these atoms. The P(V) atom adopts distorted tetrahedral geometry.

Experimental

Under an inert argon atmosphere, Mg[N(PPh₂)₂](THF)₂ (0.67 g, 0.72 mmol) was dissolved in 40 ml anhydrous THF. The solution was exposed to 2 eq. of carbon dioxide at 10 psig. After 16 h, the solution was purged with argon. Colourless crystals of the title compound crystallized from the solution over the course of two weeks. ³¹P{¹H} NMR (101.255 MHz, THF) δ 41.5 (d, ²J_{PP} = 97 Hz, Ph₂P-N), 17.8 (d of d, ²J_{PP} = 97 Hz, ¹J_{PP} = 249 Hz, N=PPh₂), -9.4 (d, ¹J_{PP} = 249 Hz, P-PPh₂) p.p.m..

Refinement

H atoms were located from a difference Fourier map and refined isotropically.

Figures




Fig. 1. View of the title compound showing full numbering scheme. Ellipsoids are shown at 50% probability and hydrogen atoms have been removed for clarity.

supplementary materials

P,P-Diphenyl-N-(1,1,2,2-tetraphenyl-1λ⁵-diphosphanylidene)phosphinous amide

Crystal data

C ₃₆ H ₃₀ NP ₃	$F_{000} = 1192$
$M_r = 569.52$	$D_x = 1.278 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 9.3026 (13) \text{ \AA}$	Cell parameters from 9384 reflections
$b = 10.8167 (15) \text{ \AA}$	$\theta = 2.3\text{--}33.2^\circ$
$c = 29.750 (4) \text{ \AA}$	$\mu = 0.23 \text{ mm}^{-1}$
$\beta = 98.589 (6)^\circ$	$T = 228 (2) \text{ K}$
$V = 2960.0 (7) \text{ \AA}^3$	Square, colourless
$Z = 4$	$0.58 \times 0.51 \times 0.18 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	11496 independent reflections
Radiation source: fine-focus sealed tube	9167 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.029$
$T = 228(2) \text{ K}$	$\theta_{\text{max}} = 33.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$h = -14\text{--}14$
$T_{\text{min}} = 0.88$, $T_{\text{max}} = 0.96$	$k = -16\text{--}16$
76797 measured reflections	$l = -45\text{--}45$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	All H-atom parameters refined
$wR(F^2) = 0.118$	$w = 1/[\sigma^2(F_o^2) + (0.0543P)^2 + 1.05P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} = 0.001$
11496 reflections	$\Delta\rho_{\text{max}} = 0.41 \text{ e \AA}^{-3}$
481 parameters	$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.89827 (12)	0.55872 (10)	0.13502 (4)	0.0285 (2)
P1	0.83490 (3)	0.52397 (3)	0.080793 (10)	0.02539 (7)
P2	0.98888 (3)	0.67406 (3)	0.154676 (10)	0.02169 (6)
P3	0.88674 (3)	0.85071 (3)	0.171889 (11)	0.02562 (7)
C1	0.92572 (12)	0.37613 (11)	0.07254 (4)	0.0265 (2)
C2	0.94010 (18)	0.34039 (15)	0.02865 (5)	0.0398 (3)
H2	0.915 (2)	0.396 (2)	0.0046 (7)	0.052 (5)*
C3	0.9911 (2)	0.22253 (18)	0.02040 (7)	0.0533 (4)
H3	0.998 (3)	0.200 (2)	-0.0104 (8)	0.067 (7)*
C4	1.03047 (18)	0.14200 (15)	0.05566 (7)	0.0494 (4)
H4	1.062 (2)	0.061 (2)	0.0493 (7)	0.064 (6)*
C5	1.02013 (17)	0.17767 (14)	0.09955 (7)	0.0433 (3)
H5	1.045 (2)	0.120 (2)	0.1238 (7)	0.059 (6)*
C6	0.96912 (15)	0.29430 (13)	0.10817 (5)	0.0344 (3)
H6	0.958 (2)	0.3171 (19)	0.1389 (7)	0.053 (6)*
C11	0.65810 (12)	0.45178 (11)	0.08595 (4)	0.0270 (2)
C12	0.60725 (15)	0.43744 (13)	0.12725 (5)	0.0339 (3)
H12	0.661 (2)	0.4670 (17)	0.1541 (6)	0.039 (5)*
C13	0.47230 (17)	0.38278 (15)	0.12874 (6)	0.0444 (3)
H13	0.437 (2)	0.380 (2)	0.1578 (7)	0.055 (6)*
C14	0.38678 (16)	0.34312 (14)	0.08938 (7)	0.0468 (4)
H14	0.297 (2)	0.306 (2)	0.0912 (7)	0.063 (6)*
C15	0.43625 (16)	0.35656 (14)	0.04816 (7)	0.0439 (4)
H15	0.378 (2)	0.332 (2)	0.0199 (7)	0.062 (6)*
C16	0.57122 (15)	0.41024 (13)	0.04646 (5)	0.0349 (3)
H16	0.6072 (19)	0.4179 (17)	0.0181 (6)	0.037 (4)*
C21	1.13511 (12)	0.72160 (10)	0.12433 (4)	0.02325 (19)
C22	1.20846 (14)	0.63055 (12)	0.10377 (5)	0.0318 (2)
H22	1.180 (2)	0.5479 (18)	0.1056 (6)	0.045 (5)*
C23	1.32338 (15)	0.66224 (14)	0.08106 (5)	0.0371 (3)
H23	1.372 (2)	0.597 (2)	0.0673 (7)	0.057 (6)*
C24	1.36496 (15)	0.78400 (14)	0.07826 (5)	0.0358 (3)
H24	1.448 (2)	0.8079 (19)	0.0628 (6)	0.049 (5)*

supplementary materials

C25	1.29177 (17)	0.87515 (14)	0.09808 (6)	0.0420 (3)
H25	1.318 (2)	0.957 (2)	0.0961 (7)	0.061 (6)*
C26	1.17755 (16)	0.84426 (12)	0.12116 (6)	0.0371 (3)
H26	1.132 (2)	0.909 (2)	0.1355 (7)	0.055 (6)*
C31	1.06971 (12)	0.63888 (11)	0.21231 (4)	0.0256 (2)
C32	1.16599 (14)	0.72246 (13)	0.23711 (5)	0.0326 (2)
H32	1.197 (2)	0.7950 (18)	0.2228 (6)	0.042 (5)*
C33	1.21154 (17)	0.70324 (17)	0.28318 (5)	0.0419 (3)
H33	1.277 (2)	0.7602 (18)	0.2992 (6)	0.045 (5)*
C34	1.16275 (19)	0.60144 (19)	0.30440 (5)	0.0488 (4)
H34	1.191 (2)	0.587 (2)	0.3358 (7)	0.056 (6)*
C35	1.0706 (2)	0.51676 (18)	0.27995 (6)	0.0491 (4)
H35	1.041 (2)	0.449 (2)	0.2946 (7)	0.061 (6)*
C36	1.02326 (16)	0.53572 (13)	0.23390 (5)	0.0362 (3)
H36	0.953 (2)	0.4794 (18)	0.2163 (6)	0.047 (5)*
C41	0.77800 (13)	0.90632 (11)	0.11987 (4)	0.0268 (2)
C42	0.78720 (15)	0.86532 (13)	0.07605 (5)	0.0334 (3)
H42	0.842 (2)	0.7942 (18)	0.0716 (6)	0.041 (5)*
C43	0.71842 (18)	0.93088 (16)	0.03862 (5)	0.0420 (3)
H43	0.727 (2)	0.9039 (19)	0.0095 (7)	0.052 (5)*
C44	0.64070 (18)	1.03665 (16)	0.04451 (6)	0.0470 (4)
H44	0.597 (2)	1.082 (2)	0.0194 (7)	0.058 (6)*
C45	0.6267 (2)	1.07566 (16)	0.08773 (7)	0.0506 (4)
H45	0.578 (3)	1.150 (2)	0.0926 (8)	0.070 (7)*
C46	0.69448 (17)	1.01113 (14)	0.12517 (6)	0.0406 (3)
H46	0.684 (3)	1.038 (2)	0.1553 (8)	0.069 (7)*
C51	0.75546 (14)	0.78245 (12)	0.20420 (4)	0.0296 (2)
C52	0.78649 (17)	0.79087 (16)	0.25158 (5)	0.0407 (3)
H52	0.878 (3)	0.839 (2)	0.2671 (8)	0.067 (7)*
C53	0.6955 (2)	0.73483 (19)	0.27841 (6)	0.0523 (4)
H53	0.719 (3)	0.744 (2)	0.3110 (8)	0.073 (7)*
C54	0.5732 (2)	0.67193 (19)	0.25838 (7)	0.0581 (5)
H54	0.512 (3)	0.633 (2)	0.2772 (8)	0.071 (7)*
C55	0.5397 (2)	0.66580 (18)	0.21142 (7)	0.0523 (4)
H55	0.460 (2)	0.623 (2)	0.1988 (7)	0.058 (6)*
C56	0.63044 (15)	0.72019 (14)	0.18414 (5)	0.0375 (3)
H56	0.608 (2)	0.7168 (18)	0.1505 (6)	0.044 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0314 (5)	0.0258 (4)	0.0285 (5)	-0.0062 (4)	0.0052 (4)	-0.0046 (4)
P1	0.02921 (14)	0.02131 (13)	0.02575 (14)	-0.00134 (10)	0.00446 (11)	0.00031 (10)
P2	0.02190 (12)	0.01984 (12)	0.02357 (13)	0.00031 (9)	0.00413 (9)	-0.00128 (9)
P3	0.02697 (13)	0.02346 (13)	0.02645 (14)	0.00363 (10)	0.00405 (11)	-0.00289 (10)
C1	0.0255 (5)	0.0250 (5)	0.0293 (5)	-0.0026 (4)	0.0054 (4)	-0.0024 (4)
C2	0.0473 (8)	0.0408 (7)	0.0325 (7)	0.0040 (6)	0.0102 (6)	-0.0065 (6)
C3	0.0592 (10)	0.0503 (10)	0.0526 (10)	0.0056 (8)	0.0156 (8)	-0.0225 (8)

C4	0.0397 (7)	0.0323 (7)	0.0766 (12)	0.0038 (6)	0.0104 (8)	-0.0161 (7)
C5	0.0372 (7)	0.0297 (6)	0.0628 (10)	0.0048 (5)	0.0063 (7)	0.0041 (6)
C6	0.0359 (6)	0.0307 (6)	0.0374 (7)	0.0035 (5)	0.0083 (5)	0.0025 (5)
C11	0.0249 (5)	0.0220 (5)	0.0331 (6)	0.0018 (4)	0.0009 (4)	-0.0004 (4)
C12	0.0308 (6)	0.0335 (6)	0.0378 (7)	-0.0013 (5)	0.0066 (5)	0.0010 (5)
C13	0.0349 (7)	0.0404 (8)	0.0605 (10)	-0.0025 (6)	0.0159 (7)	0.0065 (7)
C14	0.0275 (6)	0.0296 (6)	0.0824 (13)	-0.0023 (5)	0.0047 (7)	0.0017 (7)
C15	0.0327 (6)	0.0313 (7)	0.0621 (10)	0.0017 (5)	-0.0115 (6)	-0.0075 (6)
C16	0.0333 (6)	0.0302 (6)	0.0383 (7)	0.0035 (5)	-0.0038 (5)	-0.0041 (5)
C21	0.0229 (4)	0.0222 (5)	0.0249 (5)	0.0011 (4)	0.0042 (4)	-0.0002 (4)
C22	0.0330 (6)	0.0257 (5)	0.0389 (7)	0.0029 (4)	0.0133 (5)	-0.0014 (5)
C23	0.0362 (6)	0.0369 (7)	0.0418 (7)	0.0059 (5)	0.0179 (6)	-0.0011 (5)
C24	0.0294 (6)	0.0439 (7)	0.0362 (7)	0.0003 (5)	0.0112 (5)	0.0068 (5)
C25	0.0428 (7)	0.0304 (6)	0.0570 (9)	-0.0060 (6)	0.0216 (7)	0.0032 (6)
C26	0.0395 (6)	0.0231 (5)	0.0533 (8)	-0.0019 (5)	0.0214 (6)	-0.0032 (5)
C31	0.0246 (5)	0.0256 (5)	0.0265 (5)	0.0029 (4)	0.0035 (4)	0.0013 (4)
C32	0.0288 (5)	0.0358 (6)	0.0319 (6)	-0.0015 (5)	0.0006 (5)	-0.0008 (5)
C33	0.0358 (7)	0.0542 (9)	0.0329 (7)	0.0022 (6)	-0.0042 (5)	-0.0036 (6)
C34	0.0501 (9)	0.0644 (11)	0.0297 (7)	0.0090 (8)	-0.0009 (6)	0.0087 (7)
C35	0.0574 (9)	0.0501 (9)	0.0395 (8)	0.0004 (8)	0.0065 (7)	0.0178 (7)
C36	0.0414 (7)	0.0321 (6)	0.0347 (7)	-0.0024 (5)	0.0043 (5)	0.0063 (5)
C41	0.0270 (5)	0.0227 (5)	0.0306 (5)	0.0023 (4)	0.0044 (4)	0.0014 (4)
C42	0.0363 (6)	0.0321 (6)	0.0314 (6)	0.0050 (5)	0.0039 (5)	0.0009 (5)
C43	0.0453 (8)	0.0467 (8)	0.0322 (7)	-0.0005 (6)	-0.0002 (6)	0.0070 (6)
C44	0.0443 (8)	0.0412 (8)	0.0506 (9)	0.0016 (6)	-0.0088 (7)	0.0160 (7)
C45	0.0508 (9)	0.0344 (7)	0.0634 (11)	0.0167 (7)	-0.0021 (8)	0.0064 (7)
C46	0.0440 (7)	0.0319 (7)	0.0451 (8)	0.0143 (6)	0.0045 (6)	-0.0007 (6)
C51	0.0309 (5)	0.0318 (6)	0.0275 (5)	0.0088 (4)	0.0089 (4)	0.0001 (4)
C52	0.0431 (7)	0.0515 (9)	0.0290 (6)	0.0117 (6)	0.0105 (6)	0.0003 (6)
C53	0.0608 (10)	0.0637 (11)	0.0370 (8)	0.0184 (9)	0.0226 (7)	0.0118 (7)
C54	0.0637 (11)	0.0529 (10)	0.0678 (12)	0.0132 (8)	0.0425 (10)	0.0185 (9)
C55	0.0424 (8)	0.0508 (9)	0.0690 (12)	-0.0034 (7)	0.0252 (8)	0.0009 (8)
C56	0.0329 (6)	0.0409 (7)	0.0408 (7)	0.0017 (5)	0.0118 (5)	-0.0006 (6)

Geometric parameters (\AA , $^\circ$)

N1—P2	1.5690 (10)	C24—H24	0.99 (2)
N1—P1	1.6755 (11)	C25—C26	1.3892 (19)
P1—C1	1.8421 (12)	C25—H25	0.92 (2)
P1—C11	1.8476 (12)	C26—H26	0.95 (2)
P2—C31	1.8071 (12)	C31—C36	1.3882 (18)
P2—C21	1.8156 (11)	C31—C32	1.4023 (18)
P2—P3	2.2273 (5)	C32—C33	1.388 (2)
P3—C51	1.8196 (13)	C32—H32	0.958 (19)
P3—C41	1.8200 (13)	C33—C34	1.379 (3)
C1—C2	1.3873 (18)	C33—H33	0.94 (2)
C1—C6	1.3935 (19)	C34—C35	1.385 (3)
C2—C3	1.395 (2)	C34—H34	0.95 (2)
C2—H2	0.94 (2)	C35—C36	1.390 (2)

supplementary materials

C3—C4	1.371 (3)	C35—H35	0.92 (2)
C3—H3	0.96 (2)	C36—H36	0.99 (2)
C4—C5	1.379 (3)	C41—C42	1.3917 (18)
C4—H4	0.95 (2)	C41—C46	1.3965 (18)
C5—C6	1.385 (2)	C42—C43	1.393 (2)
C5—H5	0.96 (2)	C42—H42	0.943 (19)
C6—H6	0.97 (2)	C43—C44	1.378 (2)
C11—C12	1.3895 (18)	C43—H43	0.93 (2)
C11—C16	1.3972 (18)	C44—C45	1.378 (3)
C12—C13	1.394 (2)	C44—H44	0.93 (2)
C12—H12	0.933 (18)	C45—C46	1.384 (2)
C13—C14	1.382 (3)	C45—H45	0.95 (2)
C13—H13	0.97 (2)	C46—H46	0.96 (2)
C14—C15	1.380 (3)	C51—C56	1.398 (2)
C14—H14	0.93 (2)	C51—C52	1.3991 (19)
C15—C16	1.391 (2)	C52—C53	1.386 (2)
C15—H15	0.97 (2)	C52—H52	1.04 (2)
C16—H16	0.957 (17)	C53—C54	1.382 (3)
C21—C22	1.3909 (16)	C53—H53	0.97 (2)
C21—C26	1.3915 (17)	C54—C55	1.387 (3)
C22—C23	1.3901 (18)	C54—H54	0.96 (2)
C22—H22	0.94 (2)	C55—C56	1.386 (2)
C23—C24	1.379 (2)	C55—H55	0.91 (2)
C23—H23	0.97 (2)	C56—H56	0.990 (19)
C24—C25	1.379 (2)		
P2—N1—P1	129.01 (7)	C24—C25—C26	120.13 (13)
N1—P1—C1	102.74 (6)	C24—C25—H25	120.2 (14)
N1—P1—C11	101.80 (6)	C26—C25—H25	119.7 (14)
C1—P1—C11	94.35 (5)	C25—C26—C21	120.58 (12)
N1—P2—C31	108.08 (6)	C25—C26—H26	118.3 (13)
N1—P2—C21	116.14 (5)	C21—C26—H26	121.1 (13)
C31—P2—C21	107.15 (5)	C36—C31—C32	119.56 (12)
N1—P2—P3	122.96 (4)	C36—C31—P2	119.62 (10)
C31—P2—P3	95.39 (4)	C32—C31—P2	120.40 (9)
C21—P2—P3	104.38 (4)	C33—C32—C31	119.97 (13)
C51—P3—C41	104.57 (6)	C33—C32—H32	119.6 (11)
C51—P3—P2	96.68 (4)	C31—C32—H32	120.4 (11)
C41—P3—P2	106.86 (4)	C34—C33—C32	119.90 (15)
C2—C1—C6	118.77 (13)	C34—C33—H33	121.4 (12)
C2—C1—P1	118.55 (10)	C32—C33—H33	118.7 (12)
C6—C1—P1	122.45 (10)	C33—C34—C35	120.53 (15)
C1—C2—C3	120.23 (16)	C33—C34—H34	121.2 (13)
C1—C2—H2	119.5 (13)	C35—C34—H34	118.3 (13)
C3—C2—H2	120.2 (13)	C34—C35—C36	120.01 (15)
C4—C3—C2	120.34 (16)	C34—C35—H35	119.2 (14)
C4—C3—H3	121.6 (14)	C36—C35—H35	120.8 (14)
C2—C3—H3	118.0 (15)	C31—C36—C35	120.00 (14)
C3—C4—C5	119.92 (15)	C31—C36—H36	118.7 (11)
C3—C4—H4	119.2 (13)	C35—C36—H36	121.3 (11)

supplementary materials

C5—C4—H4	120.9 (13)	C42—C41—C46	118.52 (12)
C4—C5—C6	120.29 (16)	C42—C41—P3	125.94 (9)
C4—C5—H5	119.1 (13)	C46—C41—P3	114.86 (10)
C6—C5—H5	120.5 (13)	C41—C42—C43	120.11 (13)
C5—C6—C1	120.40 (14)	C41—C42—H42	120.0 (11)
C5—C6—H6	119.7 (12)	C43—C42—H42	119.8 (11)
C1—C6—H6	119.8 (12)	C44—C43—C42	120.54 (15)
C12—C11—C16	118.50 (12)	C44—C43—H43	119.9 (13)
C12—C11—P1	123.05 (10)	C42—C43—H43	119.6 (13)
C16—C11—P1	118.45 (10)	C45—C44—C43	119.81 (14)
C11—C12—C13	120.16 (14)	C45—C44—H44	119.6 (13)
C11—C12—H12	120.7 (11)	C43—C44—H44	120.6 (13)
C13—C12—H12	119.1 (11)	C44—C45—C46	120.09 (15)
C14—C13—C12	120.79 (16)	C44—C45—H45	121.3 (14)
C14—C13—H13	121.1 (12)	C46—C45—H45	118.4 (14)
C12—C13—H13	117.9 (13)	C45—C46—C41	120.86 (15)
C15—C14—C13	119.60 (14)	C45—C46—H46	120.0 (14)
C15—C14—H14	121.1 (13)	C41—C46—H46	119.1 (14)
C13—C14—H14	119.3 (13)	C56—C51—C52	119.65 (13)
C14—C15—C16	119.93 (15)	C56—C51—P3	123.44 (10)
C14—C15—H15	121.8 (13)	C52—C51—P3	116.88 (11)
C16—C15—H15	118.3 (13)	C53—C52—C51	119.98 (17)
C15—C16—C11	121.02 (15)	C53—C52—H52	119.4 (12)
C15—C16—H16	120.3 (11)	C51—C52—H52	120.6 (12)
C11—C16—H16	118.6 (11)	C54—C53—C52	120.06 (16)
C22—C21—C26	118.84 (11)	C54—C53—H53	122.1 (15)
C22—C21—P2	118.12 (9)	C52—C53—H53	117.8 (15)
C26—C21—P2	123.03 (9)	C53—C54—C55	120.31 (16)
C23—C22—C21	120.17 (12)	C53—C54—H54	119.3 (14)
C23—C22—H22	120.4 (12)	C55—C54—H54	120.4 (14)
C21—C22—H22	119.4 (12)	C56—C55—C54	120.30 (18)
C24—C23—C22	120.53 (12)	C56—C55—H55	120.5 (14)
C24—C23—H23	121.4 (13)	C54—C55—H55	119.2 (14)
C22—C23—H23	118.1 (13)	C55—C56—C51	119.66 (16)
C23—C24—C25	119.74 (12)	C55—C56—H56	121.4 (11)
C23—C24—H24	121.5 (12)	C51—C56—H56	118.9 (11)
C25—C24—H24	118.7 (12)		

supplementary materials

Fig. 1

