

(E)-1-(4-Bromophenyl)-3-(2-chlorophenyl)prop-2-en-1-one

Hoong-Kun Fun,^{a*} P. S. Patil,^b S. M. Dharmaprakash^b and Suchada Chantrapromma^{c‡}

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Studies in Physics, Mangalore University, Mangalagangothri, Mangalore 574 199, India, and ^cCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand
Correspondence e-mail: hkfun@usm.my


Received 3 July 2008; accepted 4 July 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.034; wR factor = 0.087; data-to-parameter ratio = 21.4.

The structure of the title compound, $C_{15}H_{10}BrClO$, comprises two substituted benzene rings bridged by a prop-2-en-1-one group and exists in an *E* configuration about the $C=N$ double bond. The dihedral angle formed between the 4-bromophenyl and 2-chlorophenyl rings is $23.77(18)^\circ$. In the crystal structure, the molecules are linked by weak $C-H \cdots O$ interactions, forming a supramolecular zigzag chain. Intramolecular $C-H \cdots Cl$ and $C-H \cdots O$ hydrogen bonds are also present.

Related literature

For related literature on hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: Patil *et al.* (2007); Moorthi *et al.* (2005). For applications of chalcones, see: Gu *et al.* (2008); Mishra *et al.* (2008); Nel *et al.* (1998); Patil & Dharmaprakash (2008); Wang *et al.* (2004).



Experimental

Crystal data

$C_{15}H_{10}BrClO$ $V = 1272.99(5) \text{ \AA}^3$
 $M_r = 321.59$ $Z = 4$
 Orthorhombic, Pna_21 Mo $K\alpha$ radiation
 $a = 27.8720(6) \text{ \AA}$ $\mu = 3.42 \text{ mm}^{-1}$
 $b = 3.9235(1) \text{ \AA}$ $T = 100.0(1) \text{ K}$
 $c = 11.6408(2) \text{ \AA}$ $0.33 \times 0.18 \times 0.09 \text{ mm}$

Data collection

Bruker SMART APEX2 CCD area-detector diffractometer 9658 measured reflections
 Absorption correction: multi-scan (SADABS; Bruker, 2005) 3495 independent reflections
 $T_{min} = 0.392, T_{max} = 0.736$ 2938 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$ H-atom parameters constrained
 $wR(F^2) = 0.086$ $\Delta\rho_{max} = 0.41 \text{ e \AA}^{-3}$
 $S = 1.03$ $\Delta\rho_{min} = -0.44 \text{ e \AA}^{-3}$
 3495 reflections Absolute structure: Flack (1983),
 163 parameters 1545 Friedel pairs
 1 restraint Flack parameter: 0.011 (12)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C1-H1A \cdots O1^i$	0.93	2.53	3.191 (4)	128
$C9-H9A \cdots Cl1$	0.93	2.61	3.064 (4)	111
$C9-H9A \cdots O1$	0.93	2.41	2.765 (5)	102

Symmetry code: (i) $-x, -y, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

This work is supported by the Department of Science and Technology (DST), Government of India, under grant No. SR/S2/LOP-17/2006. SC thanks Prince of Songkla University for generous support. The authors also thank Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2281).

References

Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Gu, B., Ji, W., Patil, P. S., Dharmaprakash, S. M. & Wang, H. T. (2008). *Appl. Phys. Lett.* 091118.
 Mishra, N., Arora, P., Kumar, B., Mishra, L. C., Bhattacharya, A., Awasthi, S. K. & Bhasin, V. K. (2008). *Eur. J. Med. Chem.* **43**, 1530–1535.
 Moorthi, S. S., Chinnakali, K., Nanjundan, S., Unnithan, C. S., Fun, H.-K. & Yu, X.-L. (2005). *Acta Cryst.* **E61**, o483–o485.
 Nel, R. J. J., Van Heerden, P. S., Van Rensburg, H. & Ferreira, D. (1998). *Tetrahedron Lett.* **39**, 5623–5626.
 Patil, P. S., Chantrapromma, S., Fun, H.-K., Dharmaprakash, S. M. & Babu, H. B. R. (2007). *Acta Cryst.* **E63**, o2612.
 Patil, P. S. & Dharmaprakash, S. M. (2008). *Mater. Lett.* **62**, 451–453.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Wang, L., Zhang, Y., Lu, C.-R. & Zhang, D.-C. (2004). *Acta Cryst.* **C60**, o696–o698.

‡ Additional correspondence author, e-mail: suchada.c@psu.ac.th.

supplementary materials

Acta Cryst. (2008). E64, o1464 [doi:10.1107/S1600536808020795]

(*E*)-1-(4-Bromophenyl)-3-(2-chlorophenyl)prop-2-en-1-one

H.-K. Fun, P. S. Patil, S. M. Dharmaprakash and S. Chantrapromma

Comment

Chalcone and its derivatives have a wide range of applications ranging from bioactivities (Mishra *et al.*, 2008; Nel *et al.*, 1998) to materials with non-linear optical (NLO) properties (Gu *et al.*, 2008 & Moorthi *et al.*, 2005). As part of our continuing interest in the latter application (Patil & Dharmaprakash, 2008), the synthesis and structure of the title compound (I, Fig. 1) is described herein. The non-centrosymmetric crystal of the title compound should exhibit 2nd-order NLO properties.

The structure of (I) comprises two six-membered rings bridged by a prop-2-en-1-one moiety. The molecule exists in the *E* conformation with respect to the C8=C9 double bond [1.328 (5) Å]. The molecule is not planar as seen in the dihedral angle of 23.77 (18)° formed between the 4-bromophenyl and 2-chlorophenyl rings. Further, the mean plane through the O1, C6, C7 & C8 atoms forms angles, respectively, of 13.2 (2)° and 11.0 (2)° with the planes of 4-bromophenyl and 2-chlorophenyl rings. Weak C9—H9A···O1 and C9—H9A···Cl1 intramolecular interactions (Fig. 1 & Table 1) generate S(5) ring motifs (Bernstein *et al.*, 1995). The derived bond distances and angles are comparable with those determined in the closely related structures (e.g. Patil *et al.*, 2007 & Sathiya Moorthi *et al.*, 2005).

In the crystal packing (Fig. 2), the molecules are linked into a supramolecular chain via C—H···O interactions aligned along the *c*-direction, Table 1.

Experimental

Compound (I) was synthesized by the condensation of 2-chlorobenzaldehyde (0.01 mol, 1.49 g) with 4-bromoacetophenone (0.01 mol, 1.99 g) in methanol (60 ml) in the presence of a catalytic amount of sodium hydroxide solution (5 ml, 20%). After stirring for 2 h, the contents of the flask were poured into ice-cold water (500 ml) and left to stand for 5 h. The resulting crude solid was filtered and dried. Single crystals were obtained by recrystallization from an acetone solution of (I).

Refinement

All H atoms were in the riding model approximation with C—H = 0.93 Å, and with $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom-numbering scheme. The dashed lines represent intramolecular C—H···O and C—H···Cl interactions.




Fig. 2. A view down the *b*-axis of the crystal packing in (I), highlighting a supramolecular chain aligned along the *c* axis. The C-H...O interactions are shown as dashed lines.

(E)-1-(4-Bromophenyl)-3-(2-chlorophenyl)prop-2-en-1-one

Crystal data

$C_{15}H_{10}BrClO$

$M_r = 321.59$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 27.8720$ (6) Å

$b = 3.9235$ (1) Å

$c = 11.6408$ (2) Å

$V = 1272.99$ (5) Å³

$Z = 4$

$F_{000} = 640$

$D_x = 1.678$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3495 reflections

$\theta = 1.5$ – 30.0°

$\mu = 3.42$ mm⁻¹

$T = 100.0$ (1) K

Block, colorless

$0.33 \times 0.18 \times 0.09$ mm

Data collection

Bruker SMART APEX2 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 8.33 pixels mm⁻¹

$T = 100.0$ (1) K

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.392$, $T_{\max} = 0.736$

9658 measured reflections

3495 independent reflections

2938 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\text{max}} = 30.0^\circ$

$\theta_{\text{min}} = 1.5^\circ$

$h = -36 \rightarrow 39$

$k = -5 \rightarrow 3$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.033$	$w = 1/[\sigma^2(F_o^2) + 1.3265P]$
$wR(F^2) = 0.086$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\max} = 0.001$
3495 reflections	$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
163 parameters	$\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1545 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.011 (12)

Special details

Experimental. The low-temperature data was collected with the Oxford Cryosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.165695 (11)	0.71499 (8)	0.27469 (5)	0.02346 (10)
Cl1	-0.18444 (4)	-0.5643 (3)	0.47411 (9)	0.0264 (2)
O1	-0.03953 (10)	0.0955 (8)	0.5096 (2)	0.0254 (6)
C1	0.03194 (13)	0.2560 (9)	0.2561 (3)	0.0187 (8)
H1A	0.0132	0.1502	0.2004	0.022*
C2	0.07645 (13)	0.3879 (11)	0.2266 (3)	0.0192 (8)
H2A	0.0878	0.3720	0.1517	0.023*
C3	0.10349 (14)	0.5426 (9)	0.3106 (3)	0.0192 (8)
C4	0.08725 (13)	0.5741 (10)	0.4230 (3)	0.0202 (8)
H4A	0.1059	0.6831	0.4782	0.024*
C5	0.04319 (13)	0.4411 (9)	0.4510 (3)	0.0177 (8)
H5A	0.0321	0.4585	0.5261	0.021*
C6	0.01502 (13)	0.2810 (9)	0.3686 (3)	0.0156 (7)
C7	-0.03108 (13)	0.1245 (10)	0.4073 (3)	0.0180 (8)
C8	-0.06613 (13)	0.0089 (10)	0.3202 (3)	0.0185 (8)
H8A	-0.0610	0.0543	0.2428	0.022*
C9	-0.10501 (14)	-0.1603 (10)	0.3534 (3)	0.0192 (8)
H9A	-0.1074	-0.2079	0.4315	0.023*
C10	-0.14477 (11)	-0.2809 (8)	0.2809 (5)	0.0180 (6)
C11	-0.18352 (14)	-0.4627 (10)	0.3288 (3)	0.0206 (8)

supplementary materials

C12	-0.22195 (13)	-0.5699 (9)	0.2625 (4)	0.0246 (8)
H12A	-0.2470	-0.6907	0.2960	0.030*
C13	-0.22293 (15)	-0.4970 (11)	0.1464 (4)	0.0276 (9)
H13A	-0.2488	-0.5662	0.1017	0.033*
C14	-0.18509 (15)	-0.3199 (11)	0.0968 (3)	0.0239 (8)
H14A	-0.1856	-0.2719	0.0185	0.029*
C15	-0.14652 (15)	-0.2140 (10)	0.1632 (3)	0.0213 (8)
H15A	-0.1214	-0.0963	0.1287	0.026*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.01631 (16)	0.02108 (18)	0.03298 (17)	-0.00286 (13)	0.0010 (2)	0.0002 (3)
Cl1	0.0257 (5)	0.0253 (5)	0.0282 (4)	-0.0021 (4)	0.0095 (4)	0.0033 (4)
O1	0.0260 (16)	0.0329 (18)	0.0171 (13)	-0.0047 (13)	0.0038 (11)	0.0008 (11)
C1	0.0159 (16)	0.0214 (19)	0.019 (2)	-0.0002 (13)	-0.0025 (13)	0.0006 (14)
C2	0.0177 (19)	0.021 (2)	0.0187 (16)	-0.0023 (15)	-0.0003 (14)	0.0018 (15)
C3	0.0170 (18)	0.0126 (19)	0.0278 (19)	0.0004 (14)	-0.0001 (14)	0.0023 (14)
C4	0.0186 (19)	0.017 (2)	0.0248 (19)	0.0025 (15)	-0.0068 (15)	-0.0046 (15)
C5	0.0180 (18)	0.017 (2)	0.0180 (17)	0.0023 (14)	0.0011 (13)	-0.0021 (14)
C6	0.0138 (17)	0.0161 (19)	0.0169 (16)	0.0025 (14)	-0.0010 (13)	0.0003 (13)
C7	0.0176 (18)	0.015 (2)	0.0218 (17)	0.0042 (14)	-0.0011 (14)	0.0011 (14)
C8	0.0153 (18)	0.022 (2)	0.0178 (16)	-0.0010 (15)	0.0015 (14)	0.0020 (14)
C9	0.018 (2)	0.020 (2)	0.0195 (16)	0.0021 (15)	-0.0003 (14)	0.0000 (14)
C10	0.0150 (14)	0.0143 (15)	0.0248 (15)	0.0030 (12)	0.001 (2)	0.0027 (19)
C11	0.0184 (19)	0.015 (2)	0.0280 (19)	0.0060 (15)	0.0042 (15)	0.0010 (15)
C12	0.0177 (17)	0.0158 (18)	0.040 (2)	0.0019 (13)	0.0016 (18)	-0.004 (2)
C13	0.020 (2)	0.022 (2)	0.041 (2)	0.0077 (17)	-0.0090 (18)	-0.0105 (18)
C14	0.025 (2)	0.025 (2)	0.0220 (19)	0.0065 (17)	-0.0067 (16)	-0.0013 (16)
C15	0.021 (2)	0.017 (2)	0.0257 (19)	-0.0003 (15)	0.0010 (15)	0.0019 (15)

Geometric parameters (\AA , $^\circ$)

Br1—C3	1.907 (4)	C8—C9	1.328 (5)
Cl1—C11	1.738 (4)	C8—H8A	0.9300
O1—C7	1.219 (4)	C9—C10	1.472 (6)
C1—C2	1.387 (5)	C9—H9A	0.9300
C1—C6	1.396 (5)	C10—C15	1.395 (7)
C1—H1A	0.9300	C10—C11	1.409 (5)
C2—C3	1.376 (5)	C11—C12	1.386 (6)
C2—H2A	0.9300	C12—C13	1.381 (7)
C3—C4	1.390 (5)	C12—H12A	0.9300
C4—C5	1.373 (5)	C13—C14	1.389 (6)
C4—H4A	0.9300	C13—H13A	0.9300
C5—C6	1.389 (5)	C14—C15	1.388 (6)
C5—H5A	0.9300	C14—H14A	0.9300
C6—C7	1.493 (5)	C15—H15A	0.9300
C7—C8	1.479 (5)		

C2—C1—C6	120.5 (3)	C7—C8—H8A	120.2
C2—C1—H1A	119.7	C8—C9—C10	127.4 (4)
C6—C1—H1A	119.7	C8—C9—H9A	116.3
C3—C2—C1	118.6 (3)	C10—C9—H9A	116.3
C3—C2—H2A	120.7	C15—C10—C11	117.2 (4)
C1—C2—H2A	120.7	C15—C10—C9	122.0 (3)
C2—C3—C4	122.0 (4)	C11—C10—C9	120.8 (5)
C2—C3—Br1	119.9 (3)	C12—C11—C10	121.7 (4)
C4—C3—Br1	118.1 (3)	C12—C11—Cl1	117.5 (3)
C5—C4—C3	118.7 (3)	C10—C11—Cl1	120.8 (3)
C5—C4—H4A	120.6	C13—C12—C11	119.8 (4)
C3—C4—H4A	120.6	C13—C12—H12A	120.1
C4—C5—C6	120.9 (3)	C11—C12—H12A	120.1
C4—C5—H5A	119.6	C12—C13—C14	119.7 (4)
C6—C5—H5A	119.6	C12—C13—H13A	120.1
C5—C6—C1	119.2 (3)	C14—C13—H13A	120.1
C5—C6—C7	117.7 (3)	C15—C14—C13	120.4 (4)
C1—C6—C7	123.0 (3)	C15—C14—H14A	119.8
O1—C7—C8	120.9 (4)	C13—C14—H14A	119.8
O1—C7—C6	119.9 (3)	C14—C15—C10	121.2 (4)
C8—C7—C6	119.2 (3)	C14—C15—H15A	119.4
C9—C8—C7	119.5 (3)	C10—C15—H15A	119.4
C9—C8—H8A	120.2		
C6—C1—C2—C3	0.1 (6)	C6—C7—C8—C9	173.5 (4)
C1—C2—C3—C4	-0.9 (6)	C7—C8—C9—C10	176.9 (3)
C1—C2—C3—Br1	178.1 (3)	C8—C9—C10—C15	-2.7 (6)
C2—C3—C4—C5	1.1 (6)	C8—C9—C10—C11	179.0 (4)
Br1—C3—C4—C5	-177.9 (3)	C15—C10—C11—C12	-0.3 (5)
C3—C4—C5—C6	-0.7 (6)	C9—C10—C11—C12	178.1 (3)
C4—C5—C6—C1	0.0 (6)	C15—C10—C11—Cl1	179.1 (3)
C4—C5—C6—C7	176.4 (3)	C9—C10—C11—Cl1	-2.5 (5)
C2—C1—C6—C5	0.3 (6)	C10—C11—C12—C13	-0.4 (6)
C2—C1—C6—C7	-175.9 (4)	Cl1—C11—C12—C13	-179.8 (3)
C5—C6—C7—O1	-10.8 (5)	C11—C12—C13—C14	0.7 (6)
C1—C6—C7—O1	165.5 (4)	C12—C13—C14—C15	-0.4 (6)
C5—C6—C7—C8	168.8 (3)	C13—C14—C15—C10	-0.2 (6)
C1—C6—C7—C8	-15.0 (5)	C11—C10—C15—C14	0.6 (5)
O1—C7—C8—C9	-7.0 (6)	C9—C10—C15—C14	-177.8 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C1—H1A \cdots O1 ⁱ	0.93	2.53	3.191 (4)	128
C9—H9A \cdots Cl1	0.93	2.61	3.064 (4)	111
C9—H9A \cdots O1	0.93	2.41	2.765 (5)	102

Symmetry codes: (i) $-x, -y, z-1/2$.

Fig. 1




Fig. 2

