

## 2'-Methyl-2'-nitro-1'-phenyl-2',3',5',6',7',7a'-hexahydrospiro[indoline-3,3'-1'H-pyrrolizin]-2-one

Yaghoub Sarrafi\* and Kamal Alimohammadi

Department of Chemistry, University of Mazandaran, 47415 Babolsar, Iran

Correspondence e-mail: ysarrafi@umz.ac.ir


Received 23 June 2008; accepted 5 July 2008

Key indicators: single-crystal X-ray study;  $T = 120$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.051;  $wR$  factor = 0.103; data-to-parameter ratio = 15.4.

The title compound,  $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$ , was synthesized by a multi-component 1,3-dipolar cycloaddition of azomethine ylide, derived from isatin and proline by a decarboxylative route, and (*E*)-1-phenyl-2-nitropropene. In the molecule, the spiro junction links a planar oxindole ring and a pyrrolidine ring in an envelope conformation. The molecular packing is stabilized by an intermolecular  $\text{N}-\text{H}\cdots\text{N}$  interaction of the oxindole and pyrrolizidine rings.

### Related literature

For related literature, see: Daly *et al.* (1986); Grigg & Sridharan (1993); Padwa (1984); Usha, Selvanayagam, Velmurugan, Ravikumar & Poornachandran (2005); Usha, Selvanayagam, Velmurugan, Ravikumar & Raghunathan (2005); Waldmann (1995).



### Experimental

#### Crystal data

$\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$   
 $M_r = 363.41$   
 Monoclinic,  $P2_1/n$   
 $a = 7.8524$  (16) Å  
 $b = 25.656$  (6) Å  
 $c = 9.1767$  (19) Å  
 $\beta = 110.489$  (4)°

$V = 1731.8$  (6) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.10$  mm<sup>-1</sup>  
 $T = 120$  (2) K  
 $0.21 \times 0.18 \times 0.15$  mm

#### Data collection

Bruker SMART 1000 CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.980$ ,  $T_{\max} = 0.989$

16064 measured reflections  
 3773 independent reflections  
 2183 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.064$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$   
 $wR(F^2) = 0.102$   
 $S = 1.01$   
 3773 reflections

245 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.25$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.28$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}'-\text{H1}'\cdots\text{N1}^1$	0.85	2.21	2.992 (3)	151

Symmetry code: (i)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We are grateful for the financial support of Mazandaran University of the Islamic Republic of Iran.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2246).

### References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS, Madison, Wisconsin, USA.  
 Daly, J. W., Spande, T. W., Whittaker, N., Highet, R. J., Feigl, D., Noshimori, N., Tokuyama, T. & Meyers, C. W. (1986). *J. Nat. Prod.* **46**, 210-???.  
 Grigg, R. & Sridharan, V. (1993). *Advances in Cycloaddition*, edited by D. P. Curran, Vol. 3, p. 161. London: Jai.  
 Padwa, A. (1984). *1,3-Dipolar Cycloaddition Chemistry*, Vols. 1 and 2. New York: Wiley.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.  
 Usha, G., Selvanayagam, S., Velmurugan, D., Ravikumar, K. & Poornachandran, M. (2005). *Acta Cryst.* **E61**, o3312-o3314.  
 Usha, G., Selvanayagam, S., Velmurugan, D., Ravikumar, K. & Raghunathan, R. (2005). *Acta Cryst.* **E61**, o3299-o3301.  
 Waldmann, H. (1995). *Synlett*, pp. 133-141.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1490 [ doi:10.1107/S1600536808020837 ]

## 2'-Methyl-2'-nitro-1'-phenyl-2',3',5',6',7',7a'-hexahydrospiro[indoline-3,3'-1'*H*-pyrrolizin]-2-one

Y. Sarrafi and K. Alimohammadi

### Comment

Multicomponent 1,3-dipolar cycloaddition reactions are considered to be one of the most useful processes for the construction of five-membered heterocyclic ring systems (Padwa, 1984; Grigg & Sridharan, 1993). These strategies offer significant advantages over more traditional approaches, allowing the construction of complex molecular architectures from easily available starting materials in a single synthetic operation without the need for isolation of intermediates. Particularly, the chemistry of the azomethine ylide has gained significance in recent years for the construction of nitrogen containing five-membered heterocycles, which are often the central ring systems of numerous natural products (Daly *et al.*, 1986; Waldmann, 1995). In contrast to similar compounds (Usha, Selvanayagam, Velmurugan, Ravikumar & Poornachandran, 2005; Usha, Selvanayagam, Velmurugan, Ravikumar & Raghunathan, 2005); Waldmann (1995)), in which the carbon atom bearing nitro group is bonded to the pyrrolidine ring, in the title compound it is bonded to the oxindole ring (Fig. 1). In the crystal structure, *N*—H···H hydrogen bonds link neighboring molecules. Molecules (Fig. 2) are also stacked in a side by side fashion along the *c* axis through  $\pi$ ··· $\pi$  interaction and are further linked by a few intermolecular C—H··· $\pi$  interactions,

### Refinement

The hydrogen atom of the NH group was found in difference Fourier synthesis. The H(C) atom positions were calculated. H(N) atom was refined in isotropic approximation in riding model, the H(C) atoms were refined in isotropic approximation in riding model with with the  $U_{\text{iso}}(\text{H})$  parameters equal to  $1.2 U_{\text{eq}}(\text{Ni})$ ,  $1.2 U_{\text{eq}}(\text{Ci})$  or  $1.5 U_{\text{eq}}(\text{Cii})$ , where  $U(\text{Ci})$  and  $U(\text{Cii})$  are respectively the equivalent thermal parameters of the (CH or CH<sub>2</sub>) and CH<sub>3</sub> carbon atoms to which the corresponding H atoms are bonded.

### Figures




Fig. 1. The molecular structure of the title compound with the numbering scheme for the atoms and 50% probability displacement ellipsoids.




Fig. 2. Packing diagram of the molecules, viewed down the *c* axis.

## 2'-Methyl-2'-nitro-1'-phenyl-2',3',5',6',7',7a'-hexahydrospiro[indoline- 3,3'-1'H-pyrrolizin]-2-one

### Crystal data

$C_{21}H_{21}N_3O_3$

$M_r = 363.41$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.8524$  (16) Å

$b = 25.656$  (6) Å

$c = 9.1767$  (19) Å

$\beta = 110.489$  (4)°

$V = 1731.8$  (6) Å<sup>3</sup>

$Z = 4$

$F_{000} = 768$

$D_x = 1.394$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 792 reflections

$\theta = 3$ – $23^\circ$

$\mu = 0.10$  mm<sup>-1</sup>

$T = 120$  (2) K

Prism, colorless

$0.21 \times 0.18 \times 0.15$  mm

### Data collection

Bruker SMART 1000 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 120$ (2) K

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.980$ ,  $T_{\max} = 0.989$

16064 measured reflections

3773 independent reflections

2183 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.064$

$\theta_{\max} = 27.0^\circ$

$\theta_{\min} = 1.6^\circ$

$h = -10 \rightarrow 10$

$k = -32 \rightarrow 32$

$l = -11 \rightarrow 11$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.103$

$S = 1.01$

3773 reflections

245 parameters

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.01P)^2 + 1.6P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.25 \text{ e } \text{Å}^{-3}$$

$$\Delta\rho_{\min} = -0.27 \text{ e } \text{Å}^{-3}$$

Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Experimental.** A mixture of isatin (0.147 g, 1 mmol), proline (0.115 g, 1 mmol), and (*E*)-1-phenyl-2-nitropropene (0.163 g, 1 mmol) in ethanol (10 ml) was stirred at reflux for 1 h. After completion of the reaction, as indicated by TLC, to the solution was added water (25 ml), and the precipitated solid was separated by filtration. The pure cycloadduct was obtained by recrystallization from ethanol.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.1573 (3)	0.19274 (7)	0.7966 (2)	0.0215 (4)
C2	0.1056 (3)	0.17387 (9)	0.6367 (3)	0.0210 (5)
C3	0.2848 (3)	0.14420 (9)	0.6478 (3)	0.0218 (5)
C4	0.3236 (3)	0.11281 (9)	0.8003 (3)	0.0209 (5)
H4A	0.2295	0.0859	0.7796	0.025*
C5	0.2810 (3)	0.15438 (9)	0.9046 (3)	0.0212 (5)
H5A	0.3944	0.1722	0.9645	0.025*
C6	0.1843 (3)	0.13826 (9)	1.0155 (3)	0.0250 (6)
H6A	0.2702	0.1284	1.1166	0.030*
H6B	0.1005	0.1097	0.9735	0.030*
C7	0.0836 (3)	0.18844 (9)	1.0247 (3)	0.0273 (6)
H7A	-0.0140	0.1817	1.0640	0.033*
H7B	0.1657	0.2141	1.0906	0.033*
C8	0.0098 (3)	0.20643 (10)	0.8562 (3)	0.0250 (6)
H8A	-0.1020	0.1883	0.7986	0.030*
H8B	-0.0129	0.2437	0.8495	0.030*
C9	0.4441 (3)	0.17830 (9)	0.6496 (3)	0.0245 (6)
H9A	0.5437	0.1566	0.6496	0.037*
H9B	0.4812	0.1996	0.7413	0.037*
H9C	0.4079	0.2002	0.5589	0.037*
C10	0.5061 (3)	0.08576 (9)	0.8734 (3)	0.0208 (5)
C11	0.5235 (3)	0.03378 (10)	0.8389 (3)	0.0285 (6)
H11A	0.4285	0.0173	0.7622	0.034*
C12	0.6799 (4)	0.00634 (10)	0.9170 (3)	0.0333 (6)
H12A	0.6905	-0.0282	0.8902	0.040*
C13	0.8199 (3)	0.02934 (10)	1.0338 (3)	0.0298 (6)
H13A	0.9226	0.0101	1.0894	0.036*
C14	0.8070 (3)	0.08119 (10)	1.0682 (3)	0.0280 (6)

## supplementary materials

---

H14A	0.9018	0.0972	1.1465	0.034*
C15	0.6529 (3)	0.10927 (10)	0.9861 (3)	0.0254 (6)
H15A	0.6475	0.1446	1.0068	0.030*
N2	0.2395 (3)	0.11023 (8)	0.5039 (2)	0.0241 (5)
O1	0.2495 (2)	0.06314 (7)	0.5158 (2)	0.0346 (4)
O2	0.1929 (2)	0.13377 (7)	0.37859 (19)	0.0310 (4)
N1'	-0.1068 (3)	0.21391 (8)	0.4167 (2)	0.0247 (5)
H1'	-0.1647	0.2377	0.3535	0.030*
C2'	0.0582 (3)	0.22165 (9)	0.5263 (3)	0.0245 (5)
O2'	0.1511 (2)	0.26105 (6)	0.54370 (19)	0.0298 (4)
C3A	-0.0662 (3)	0.14082 (9)	0.5689 (3)	0.0215 (5)
C4'	-0.1202 (3)	0.09354 (9)	0.6109 (3)	0.0248 (6)
H4D	-0.0455	0.0761	0.6989	0.030*
C5'	-0.2865 (3)	0.07228 (10)	0.5209 (3)	0.0271 (6)
H5D	-0.3224	0.0403	0.5482	0.032*
C6'	-0.3983 (3)	0.09824 (10)	0.3915 (3)	0.0266 (6)
H6D	-0.5086	0.0833	0.3316	0.032*
C7'	-0.3494 (3)	0.14616 (10)	0.3490 (3)	0.0252 (6)
H7D	-0.4259	0.1641	0.2628	0.030*
C7A	-0.1828 (3)	0.16639 (9)	0.4391 (3)	0.0231 (5)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0226 (11)	0.0248 (11)	0.0180 (10)	0.0022 (9)	0.0084 (9)	-0.0003 (8)
C2	0.0219 (13)	0.0213 (13)	0.0192 (12)	0.0010 (10)	0.0066 (10)	-0.0011 (10)
C3	0.0227 (13)	0.0227 (13)	0.0187 (12)	-0.0006 (10)	0.0058 (10)	-0.0016 (10)
C4	0.0227 (13)	0.0201 (12)	0.0196 (12)	-0.0022 (10)	0.0072 (10)	-0.0008 (10)
C5	0.0206 (12)	0.0239 (13)	0.0191 (12)	0.0012 (10)	0.0069 (10)	-0.0007 (10)
C6	0.0292 (14)	0.0265 (14)	0.0219 (13)	0.0027 (11)	0.0122 (11)	0.0021 (11)
C7	0.0309 (14)	0.0290 (14)	0.0252 (14)	0.0025 (11)	0.0136 (12)	0.0001 (11)
C8	0.0219 (13)	0.0284 (14)	0.0257 (13)	0.0032 (11)	0.0098 (11)	-0.0006 (11)
C9	0.0234 (13)	0.0292 (14)	0.0227 (13)	-0.0025 (11)	0.0101 (11)	-0.0005 (11)
C10	0.0244 (13)	0.0231 (13)	0.0178 (12)	0.0008 (10)	0.0110 (10)	0.0038 (10)
C11	0.0307 (15)	0.0270 (14)	0.0261 (14)	0.0025 (11)	0.0078 (12)	-0.0007 (11)
C12	0.0392 (16)	0.0280 (15)	0.0337 (15)	0.0072 (13)	0.0141 (13)	-0.0019 (12)
C13	0.0281 (14)	0.0364 (15)	0.0283 (14)	0.0096 (12)	0.0141 (12)	0.0089 (12)
C14	0.0236 (14)	0.0354 (15)	0.0255 (13)	0.0004 (11)	0.0093 (11)	0.0008 (12)
C15	0.0267 (14)	0.0265 (13)	0.0253 (13)	0.0019 (11)	0.0119 (11)	0.0012 (11)
N2	0.0232 (11)	0.0297 (12)	0.0201 (11)	0.0006 (9)	0.0084 (9)	-0.0007 (10)
O1	0.0474 (12)	0.0247 (10)	0.0324 (10)	0.0009 (9)	0.0149 (9)	-0.0043 (8)
O2	0.0330 (10)	0.0415 (11)	0.0188 (9)	0.0023 (8)	0.0093 (8)	0.0017 (8)
N1'	0.0249 (11)	0.0259 (11)	0.0214 (11)	0.0018 (9)	0.0057 (9)	0.0058 (9)
C2'	0.0284 (14)	0.0247 (13)	0.0220 (13)	0.0012 (11)	0.0107 (11)	-0.0031 (11)
O2'	0.0319 (10)	0.0242 (10)	0.0318 (10)	-0.0025 (8)	0.0093 (8)	0.0024 (8)
C3A	0.0217 (13)	0.0231 (13)	0.0204 (12)	0.0022 (10)	0.0083 (10)	-0.0025 (10)
C4'	0.0262 (14)	0.0244 (13)	0.0240 (13)	0.0048 (11)	0.0090 (11)	0.0017 (11)
C5'	0.0278 (14)	0.0259 (14)	0.0298 (14)	-0.0018 (11)	0.0131 (12)	-0.0023 (11)

C6'	0.0205 (13)	0.0325 (15)	0.0267 (14)	-0.0033 (11)	0.0082 (11)	-0.0093 (11)
C7'	0.0266 (14)	0.0293 (14)	0.0190 (12)	0.0033 (11)	0.0071 (11)	-0.0006 (11)
C7A	0.0276 (14)	0.0232 (13)	0.0200 (13)	0.0012 (11)	0.0102 (11)	-0.0019 (10)

*Geometric parameters (Å, °)*

N1—C2	1.461 (3)	C10—C11	1.388 (3)
N1—C8	1.486 (3)	C11—C12	1.379 (3)
N1—C5	1.490 (3)	C11—H11A	0.9300
C2—C3A	1.530 (3)	C12—C13	1.372 (3)
C2—C2'	1.550 (3)	C12—H12A	0.9300
C2—C3	1.571 (3)	C13—C14	1.379 (3)
C3—N2	1.517 (3)	C13—H13A	0.9300
C3—C9	1.522 (3)	C14—C15	1.383 (3)
C3—C4	1.550 (3)	C14—H14A	0.9300
C4—C10	1.521 (3)	C15—H15A	0.9300
C4—C5	1.546 (3)	N2—O1	1.213 (2)
C4—H4A	0.9800	N2—O2	1.235 (2)
C5—C6	1.525 (3)	N1'—C2'	1.348 (3)
C5—H5A	0.9800	N1'—C7A	1.404 (3)
C6—C7	1.528 (3)	N1'—H1'	0.8544
C6—H6A	0.9700	C2'—O2'	1.224 (3)
C6—H6B	0.9700	C3A—C4'	1.383 (3)
C7—C8	1.521 (3)	C3A—C7A	1.387 (3)
C7—H7A	0.9700	C4'—C5'	1.389 (3)
C7—H7B	0.9700	C4'—H4D	0.9300
C8—H8A	0.9700	C5'—C6'	1.377 (3)
C8—H8B	0.9700	C5'—H5D	0.9300
C9—H9A	0.9600	C6'—C7'	1.384 (3)
C9—H9B	0.9600	C6'—H6D	0.9300
C9—H9C	0.9600	C7'—C7A	1.381 (3)
C10—C15	1.388 (3)	C7'—H7D	0.9300
C2—N1—C8	118.04 (18)	C3—C9—H9C	109.5
C2—N1—C5	109.64 (18)	H9A—C9—H9C	109.5
C8—N1—C5	108.79 (17)	H9B—C9—H9C	109.5
N1—C2—C3A	119.15 (19)	C15—C10—C11	117.7 (2)
N1—C2—C2'	108.24 (18)	C15—C10—C4	122.6 (2)
C3A—C2—C2'	101.33 (18)	C11—C10—C4	119.4 (2)
N1—C2—C3	99.68 (17)	C12—C11—C10	120.8 (2)
C3A—C2—C3	113.59 (18)	C12—C11—H11A	119.6
C2'—C2—C3	115.5 (2)	C10—C11—H11A	119.6
N2—C3—C9	106.34 (19)	C13—C12—C11	120.7 (2)
N2—C3—C4	113.56 (19)	C13—C12—H12A	119.6
C9—C3—C4	112.82 (19)	C11—C12—H12A	119.6
N2—C3—C2	106.86 (17)	C12—C13—C14	119.4 (2)
C9—C3—C2	115.85 (19)	C12—C13—H13A	120.3
C4—C3—C2	101.46 (18)	C14—C13—H13A	120.3
C10—C4—C5	114.60 (18)	C15—C14—C13	119.9 (2)
C10—C4—C3	119.50 (19)	C15—C14—H14A	120.1

## supplementary materials

---

C5—C4—C3	100.59 (18)	C13—C14—H14A	120.1
C10—C4—H4A	107.1	C14—C15—C10	121.3 (2)
C5—C4—H4A	107.1	C14—C15—H15A	119.4
C3—C4—H4A	107.1	C10—C15—H15A	119.4
N1—C5—C6	105.17 (18)	O1—N2—O2	124.0 (2)
N1—C5—C4	106.01 (17)	O1—N2—C3	120.35 (19)
C6—C5—C4	119.5 (2)	O2—N2—C3	115.60 (19)
N1—C5—H5A	108.6	C2'—N1'—C7A	111.4 (2)
C6—C5—H5A	108.6	C2'—N1'—H1'	123.1
C4—C5—H5A	108.6	C7A—N1'—H1'	124.5
C5—C6—C7	101.31 (19)	O2'—C2'—N1'	126.3 (2)
C5—C6—H6A	111.5	O2'—C2'—C2	124.9 (2)
C7—C6—H6A	111.5	N1'—C2'—C2	108.6 (2)
C5—C6—H6B	111.5	C4'—C3A—C7A	118.7 (2)
C7—C6—H6B	111.5	C4'—C3A—C2	133.3 (2)
H6A—C6—H6B	109.3	C7A—C3A—C2	108.1 (2)
C8—C7—C6	102.72 (19)	C3A—C4'—C5'	119.6 (2)
C8—C7—H7A	111.2	C3A—C4'—H4D	120.2
C6—C7—H7A	111.2	C5'—C4'—H4D	120.2
C8—C7—H7B	111.2	C6'—C5'—C4'	120.4 (2)
C6—C7—H7B	111.2	C6'—C5'—H5D	119.8
H7A—C7—H7B	109.1	C4'—C5'—H5D	119.8
N1—C8—C7	103.52 (18)	C5'—C6'—C7'	121.1 (2)
N1—C8—H8A	111.1	C5'—C6'—H6D	119.4
C7—C8—H8A	111.1	C7'—C6'—H6D	119.4
N1—C8—H8B	111.1	C7A—C7'—C6'	117.6 (2)
C7—C8—H8B	111.1	C7A—C7'—H7D	121.2
H8A—C8—H8B	109.0	C6'—C7'—H7D	121.2
C3—C9—H9A	109.5	C7'—C7A—C3A	122.6 (2)
C3—C9—H9B	109.5	C7'—C7A—N1'	126.9 (2)
H9A—C9—H9B	109.5	C3A—C7A—N1'	110.5 (2)
C8—N1—C2—C3A	-34.1 (3)	C4—C10—C11—C12	-172.1 (2)
C5—N1—C2—C3A	91.2 (2)	C10—C11—C12—C13	1.9 (4)
C8—N1—C2—C2'	80.8 (2)	C11—C12—C13—C14	-3.0 (4)
C5—N1—C2—C2'	-153.89 (18)	C12—C13—C14—C15	0.7 (4)
C8—N1—C2—C3	-158.11 (19)	C13—C14—C15—C10	2.9 (4)
C5—N1—C2—C3	-32.8 (2)	C11—C10—C15—C14	-3.9 (3)
N1—C2—C3—N2	165.61 (17)	C4—C10—C15—C14	169.5 (2)
C3A—C2—C3—N2	37.8 (2)	C9—C3—N2—O1	119.1 (2)
C2'—C2—C3—N2	-78.7 (2)	C4—C3—N2—O1	-5.6 (3)
N1—C2—C3—C9	-76.1 (2)	C2—C3—N2—O1	-116.6 (2)
C3A—C2—C3—C9	156.02 (19)	C9—C3—N2—O2	-61.1 (2)
C2'—C2—C3—C9	39.5 (3)	C4—C3—N2—O2	174.21 (19)
N1—C2—C3—C4	46.4 (2)	C2—C3—N2—O2	63.2 (2)
C3A—C2—C3—C4	-81.4 (2)	C7A—N1'—C2'—O2'	-174.3 (2)
C2'—C2—C3—C4	162.10 (19)	C7A—N1'—C2'—C2	1.2 (3)
N2—C3—C4—C10	77.2 (3)	N1—C2—C2'—O2'	47.5 (3)
C9—C3—C4—C10	-43.9 (3)	C3A—C2—C2'—O2'	173.6 (2)
C2—C3—C4—C10	-168.54 (19)	C3—C2—C2'—O2'	-63.2 (3)



N2—C3—C4—C5	-156.50 (18)	N1—C2—C2'—N1'	-128.0 (2)
C9—C3—C4—C5	82.4 (2)	C3A—C2—C2'—N1'	-1.9 (2)
C2—C3—C4—C5	-42.2 (2)	C3—C2—C2'—N1'	121.3 (2)
C2—N1—C5—C6	-120.7 (2)	N1—C2—C3A—C4'	-59.9 (4)
C8—N1—C5—C6	9.7 (2)	C2'—C2—C3A—C4'	-178.4 (3)
C2—N1—C5—C4	6.8 (2)	C3—C2—C3A—C4'	57.1 (3)
C8—N1—C5—C4	137.19 (19)	N1—C2—C3A—C7A	120.6 (2)
C10—C4—C5—N1	152.38 (19)	C2'—C2—C3A—C7A	2.1 (2)
C3—C4—C5—N1	22.8 (2)	C3—C2—C3A—C7A	-122.5 (2)
C10—C4—C5—C6	-89.2 (3)	C7A—C3A—C4'—C5'	1.6 (3)
C3—C4—C5—C6	141.2 (2)	C2—C3A—C4'—C5'	-177.9 (2)
N1—C5—C6—C7	-32.2 (2)	C3A—C4'—C5'—C6'	-0.7 (4)
C4—C5—C6—C7	-151.0 (2)	C4'—C5'—C6'—C7'	-0.7 (4)
C5—C6—C7—C8	42.7 (2)	C5'—C6'—C7'—C7A	1.3 (4)
C2—N1—C8—C7	142.8 (2)	C6'—C7'—C7A—C3A	-0.4 (4)
C5—N1—C8—C7	17.1 (2)	C6'—C7'—C7A—N1'	179.8 (2)
C6—C7—C8—N1	-37.2 (2)	C4'—C3A—C7A—C7'	-1.0 (4)
C5—C4—C10—C15	-27.0 (3)	C2—C3A—C7A—C7'	178.6 (2)
C3—C4—C10—C15	92.4 (3)	C4'—C3A—C7A—N1'	178.8 (2)
C5—C4—C10—C11	146.3 (2)	C2—C3A—C7A—N1'	-1.5 (3)
C3—C4—C10—C11	-94.2 (3)	C2'—N1'—C7A—C7'	-179.9 (2)
C15—C10—C11—C12	1.5 (4)	C2'—N1'—C7A—C3A	0.2 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1'—H1'...N1 <sup>i</sup>	0.85	2.21	2.992 (3)	151

Symmetry codes: (i)  $x-1/2, -y+1/2, z-1/2$ .

Fig. 1




Fig. 2

