

1-(2,3-Di-O-acetyl-4-chloro-4-deoxy-6-O-tosyl- β -D-galactopyranosyl)propan-2-one methanol 0.25-solvate

Lin Yan,^{a*} Feng-Wu Liu^b and Hong-Min Liu^b

^aInstitute of Pharmacy, Henan University, Kaifeng, Henan 475001, People's Republic of China, and ^bNew Drug Research and Development Center, Zhengzhou University, Zhengzhou, Henan 450052, People's Republic of China
Correspondence e-mail: yanlin0378@163.com


Received 27 March 2008; accepted 16 July 2008

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; disorder in main residue; R factor = 0.062; wR factor = 0.154; data-to-parameter ratio = 13.6.

The asymmetric unit of the title solvate, $\text{C}_{20}\text{H}_{25}\text{ClO}_9\text{S} \cdot 0.25\text{CH}_3\text{OH}$, contains one galactopyranosyl derivative and one-quarter of a methanol solvent molecule. The galactopyranose ring is in the usual $^4\text{C}_1$ conformation, and the anomeric center of the sugar has a β configuration. The value of θ (3.44°) and the range of torsion angles [or 53.1 (5)– 63.0 (5) $^\circ$] reflect a slight distortion of the $^4\text{C}_1$ pyranose ring. A minor orientational disorder affects a carbonyl group, which was modeled with two sites for the O atom having occupancies of 0.79 (5) and 0.21 (5). The crystal studied exhibited inversion twinning.

Related literature

For related literature, see: Lewis *et al.* (1982); Nicolaou *et al.* (1995); Paterson & Mansuri (1985); Postema (1992); Tvaroška *et al.* (2002).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{25}\text{ClO}_9\text{S} \cdot 0.25\text{CH}_3\text{O}$	$V = 2390.9$ (8) \AA^3
$M_r = 484.92$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 7.3300$ (15) Å	$\mu = 0.29$ mm $^{-1}$
$b = 14.608$ (3) Å	$T = 291$ (2) K
$c = 22.329$ (5) Å	$0.20 \times 0.17 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer	7242 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	4129 independent reflections
$(SADABS$; Bruker, 2001)	3274 reflections with $I > 2\sigma(I)$
$R_{\text{int}} = 0.052$	
$T_{\min} = 0.944$, $T_{\max} = 0.954$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$	H-atom parameters constrained
$wR(F^2) = 0.154$	$\Delta\rho_{\max} = 0.19$ e \AA^{-3}
$S = 1.00$	$\Delta\rho_{\min} = -0.29$ e \AA^{-3}
4129 reflections	Absolute structure: Flack (1983), 1683 Friedel pairs
304 parameters	Flack parameter: 0.42 (11)
2 restraints	

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97*; software used to prepare material for publication: *SHELXL97*.

This work was financially supported by the NNSF of the People's Republic of China (No. 20572103).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2169).

References

- Bruker (2001). *SMART*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
Lewis, M. D., Cha, J. K. & Kishi, Y. (1982). *J. Am. Chem. Soc.* **104**, 4976–4978.
Nicolaou, K. C., Theodorakis, E. A., Rutjes, F. P. J. T., Tiebes, J., Sato, M., Untersteller, E. & Xiao, X.-Y. (1995). *J. Am. Chem. Soc.* **117**, 1171–1172.
Paterson, I. & Mansuri, M. M. (1985). *Tetrahedron*, **41**, 3569–3624.
Postema, M. H. D. (1992). *Tetrahedron*, **48**, 8545–8599.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Tvaroška, I., Taravel, F. R., Utile, J. P. & Carver, J. P. (2002). *Carbohydr. Res.* **337**, 353–367.

supplementary materials

Acta Cryst. (2008). E64, o1558 [doi:10.1107/S1600536808022253]

1-(2,3-Di-*O*-acetyl-4-chloro-4-deoxy-6-*O*-tosyl- β -D-galactopyranosyl)propan-2-one methanol 0.25-solvate

L. Yan, F.-W. Liu and H.-M. Liu

Comment

Due to their unique chemical and enzymatic hydrolysis stability, C-glycosides are becoming useful building blocks (Postema, 1992) for the total synthesis of various types of natural products such as palytoxin (Lewis *et al.*, 1982), brevetoxin (Nicolaou *et al.*, 1995) and polyether antibiotics (Paterson & Mansuri, 1985), and are used as a model in enzymatic and metabolic studies as well. Despite this attractive applications, structural investigations by using single-crystal X-ray analysis, which provides unambiguous structural data, are rare. Herein we report the design and synthesis of an acylated C-glycosidic analog, which would be of great interest in order to get additional crystallographic information about the substrate.

In the orthorhombic crystals of the title compound, the asymmetric unit contains one molecule and 0.25 methanol solvate. No significant hydrogen bonds exist in the crystal. The value of θ (3.44°) and the magnitude of the torsion angles in the ring ($52.9\text{--}63.6^\circ$) reveal that the $^4\text{C}_1$ pyranose ring presents a slight distortion. The primary hydroxyl group in the title compound is in the gt position [$\text{O}1\text{---C}5\text{---C}6\text{---O}7 = 74.9\text{ (5)}^\circ$], which is known to be the favored orientation for pyranose with the *galacto* configuration (Tvaroška *et al.*, 2002).

Experimental

All reagents were commercially available and of analytical grade. Sulfenylation of 1-(4-chloro-4-deoxy- β -D-galactopyranosyl)-propan-2-one with toluene-4-sulfonyl chloride in dry pyridine afforded the 6-toluenesulfonated intermediate. Further acetylation with acetyl anhydride in pyridine and subsequent purification by chromatography on silica gel furnished the title compound as a white solid. White crystals suitable for X-ray crystallographic analysis were obtained by recrystallization from methanol.

Refinement

All H atoms bonded to C atoms were positioned geometrically and refined as riding to their parent atoms, with $\text{C---H} = 0.93\text{--}0.98\text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. For the methanol molecule, the hydroxyl H atom was positioned geometrically and refined as riding with $\text{O---H} = 0.82\text{\AA}$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. Assuming that starting material is enantiomerically pure, and that anomalous dispersion effects from S and Cl atoms are significant, we suppose that the refined Flack parameter, 0.42 (11) based on 1683 measured Friedel pairs, reflects a partial twinning by merohedry for the sample used for data collection.

supplementary materials

Figures




Fig. 1. The molecular structure of (I), showing atom displacement ellipsoids drawn at the 50% probability level.

1-(2,3-Di-O-acetyl-4-chloro-4-deoxy-6-O-tosyl- β -D- galactopyranosyl)propan-2-one methanol 0.25-solvate

Crystal data

C ₂₀ H ₂₅ ClO ₉ S·0.25CH ₄ O	$F_{000} = 1018$
$M_r = 484.92$	$D_x = 1.347 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 7.3300 (15) \text{ \AA}$	Cell parameters from 399 reflections
$b = 14.608 (3) \text{ \AA}$	$\theta = 2\text{--}25.1^\circ$
$c = 22.329 (5) \text{ \AA}$	$\mu = 0.29 \text{ mm}^{-1}$
$V = 2390.9 (8) \text{ \AA}^3$	$T = 291 (2) \text{ K}$
$Z = 4$	Prism, colourless
	$0.20 \times 0.17 \times 0.16 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	4129 independent reflections
Radiation source: fine-focus sealed tube	3274 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.052$
$T = 291(2) \text{ K}$	$\theta_{\text{max}} = 25.3^\circ$
0.3° wide ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.944$, $T_{\text{max}} = 0.954$	$k = -17 \rightarrow 17$
7242 measured reflections	$l = -26 \rightarrow 26$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.062$	$w = 1/[\sigma^2(F_o^2) + (0.0873P)^2 + 0.6536P]$
$wR(F^2) = 0.154$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\text{max}} < 0.001$
4129 reflections	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$

304 parameters
 2 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Extinction correction: none
 Absolute structure: Flack (1983), 1683 Friedel pairs
 Flack parameter: 0.42 (11)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.32977 (15)	0.92085 (8)	0.03975 (5)	0.0411 (3)	
C11	0.65974 (18)	1.05079 (11)	0.24709 (5)	0.0662 (4)	
O1	0.7356 (4)	1.1010 (2)	0.11261 (14)	0.0425 (7)	
O2	0.8839 (8)	1.1761 (3)	-0.01365 (18)	0.0918 (16)	
O3	1.1793 (4)	1.1764 (2)	0.16930 (14)	0.0481 (8)	
O4	1.113 (5)	1.3083 (19)	0.2077 (18)	0.054 (11)	0.21 (5)
O4'	1.132 (2)	1.2846 (17)	0.2385 (15)	0.137 (9)	0.79 (5)
O5	1.0691 (4)	1.0491 (2)	0.25895 (13)	0.0512 (8)	
O6	1.2341 (6)	0.9227 (3)	0.2414 (2)	0.0779 (12)	
O7	0.5241 (4)	0.9533 (2)	0.06115 (13)	0.0453 (8)	
O8	0.3338 (5)	0.9324 (2)	-0.02346 (13)	0.0599 (9)	
O9	0.1958 (4)	0.9671 (2)	0.07483 (15)	0.0507 (8)	
O10	1.184 (7)	1.295 (3)	0.3532 (19)	0.205 (15)*	0.25
H10	1.1133	1.2916	0.3250	0.308*	0.25
C1	0.9128 (6)	1.1430 (3)	0.1106 (2)	0.0389 (10)	
H1A	0.9924	1.1071	0.0843	0.047*	
C2	0.9930 (6)	1.1440 (3)	0.17380 (19)	0.0384 (10)	
H2A	0.9211	1.1834	0.2003	0.046*	
C3	1.0018 (6)	1.0463 (3)	0.19808 (18)	0.0400 (10)	
H3A	1.0880	1.0112	0.1736	0.048*	
C4	0.8168 (7)	1.0005 (3)	0.19472 (19)	0.0425 (10)	
H4A	0.8316	0.9356	0.2047	0.051*	
C5	0.7433 (7)	1.0076 (3)	0.1312 (2)	0.0423 (11)	
H5A	0.8257	0.9745	0.1043	0.051*	
C6	0.5551 (7)	0.9680 (4)	0.1250 (2)	0.0528 (13)	
H6A	0.5464	0.9105	0.1466	0.063*	
H6B	0.4649	1.0100	0.1410	0.063*	
C7	0.8895 (7)	1.2381 (3)	0.0847 (2)	0.0525 (12)	
H7A	0.9911	1.2757	0.0979	0.063*	
H7B	0.7790	1.2647	0.1012	0.063*	
C8	0.8787 (7)	1.2427 (4)	0.0170 (2)	0.0544 (13)	
C9	0.8655 (10)	1.3370 (4)	-0.0087 (3)	0.0760 (18)	
H9A	0.8524	1.3331	-0.0514	0.114*	
H9B	0.9742	1.3706	0.0008	0.114*	
H9C	0.7615	1.3677	0.0080	0.114*	
C10	1.2286 (8)	1.2513 (4)	0.1997 (4)	0.0768 (19)	
C11	1.4220 (9)	1.2763 (5)	0.1873 (4)	0.101 (3)	
H11A	1.4464	1.3363	0.2029	0.152*	
H11B	1.4430	1.2759	0.1449	0.152*	
H11C	1.5014	1.2327	0.2063	0.152*	

supplementary materials

C12	1.1872 (7)	0.9807 (4)	0.2751 (2)	0.0527 (12)	
C13	1.2456 (10)	0.9929 (6)	0.3384 (3)	0.093 (2)	
H13A	1.3539	0.9577	0.3457	0.139*	
H13B	1.1502	0.9726	0.3647	0.139*	
H13C	1.2705	1.0565	0.3458	0.139*	
C14	0.3198 (6)	0.8038 (3)	0.05653 (18)	0.0417 (10)	
C15	0.2354 (7)	0.7735 (3)	0.1088 (2)	0.0492 (12)	
H15A	0.1912	0.8151	0.1368	0.059*	
C16	0.2191 (8)	0.6808 (3)	0.1182 (2)	0.0557 (13)	
H16A	0.1628	0.6605	0.1531	0.067*	
C17	0.2823 (7)	0.6174 (3)	0.0783 (2)	0.0524 (13)	
C18	0.3687 (7)	0.6498 (4)	0.0270 (3)	0.0593 (14)	
H18A	0.4158	0.6077	-0.0002	0.071*	
C19	0.3868 (7)	0.7414 (4)	0.0151 (2)	0.0547 (13)	
H19A	0.4429	0.7613	-0.0199	0.066*	
C20	0.2580 (10)	0.5161 (4)	0.0872 (3)	0.0769 (18)	
H20A	0.2353	0.5037	0.1288	0.115*	
H20B	0.3666	0.4847	0.0748	0.115*	
H20D	0.1565	0.4952	0.0637	0.115*	
C21	1.292 (6)	1.229 (3)	0.3514 (17)	0.135 (14)*	0.25
H21A	1.2769	1.1924	0.3866	0.202*	0.25
H21B	1.4149	1.2518	0.3496	0.202*	0.25
H21C	1.2672	1.1930	0.3164	0.202*	0.25

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0419 (6)	0.0444 (6)	0.0369 (5)	-0.0053 (5)	-0.0003 (5)	-0.0019 (5)
Cl1	0.0561 (7)	0.1008 (11)	0.0417 (6)	-0.0185 (8)	0.0082 (6)	-0.0121 (6)
O1	0.0387 (16)	0.0413 (18)	0.0476 (18)	-0.0039 (13)	-0.0022 (14)	0.0046 (14)
O2	0.167 (5)	0.053 (2)	0.055 (2)	-0.008 (3)	-0.020 (3)	0.002 (2)
O3	0.0376 (16)	0.0496 (18)	0.0570 (19)	-0.0041 (16)	0.0007 (15)	-0.0052 (15)
O4	0.062 (14)	0.026 (14)	0.07 (2)	-0.006 (9)	-0.013 (12)	-0.011 (12)
O4'	0.086 (6)	0.121 (12)	0.20 (2)	-0.026 (7)	0.017 (10)	-0.107 (14)
O5	0.0522 (18)	0.065 (2)	0.0363 (18)	0.0044 (17)	-0.0092 (14)	-0.0027 (16)
O6	0.078 (3)	0.083 (3)	0.073 (3)	0.026 (2)	-0.012 (2)	-0.003 (2)
O7	0.0475 (17)	0.057 (2)	0.0314 (16)	-0.0150 (16)	0.0000 (13)	-0.0044 (15)
O8	0.069 (2)	0.075 (2)	0.0353 (17)	-0.013 (2)	-0.0064 (16)	0.0054 (15)
O9	0.0485 (18)	0.0431 (18)	0.060 (2)	0.0050 (15)	0.0037 (16)	-0.0022 (15)
C1	0.041 (2)	0.032 (2)	0.044 (3)	-0.0023 (19)	0.0048 (19)	0.005 (2)
C2	0.035 (2)	0.042 (3)	0.039 (2)	0.0004 (19)	0.0023 (18)	-0.002 (2)
C3	0.044 (2)	0.046 (3)	0.030 (2)	0.000 (2)	0.0017 (18)	-0.002 (2)
C4	0.051 (3)	0.037 (2)	0.039 (2)	-0.006 (2)	0.002 (2)	-0.0006 (18)
C5	0.046 (3)	0.043 (3)	0.037 (2)	-0.006 (2)	0.0000 (19)	0.0007 (19)
C6	0.055 (3)	0.061 (3)	0.042 (3)	-0.024 (3)	0.002 (2)	-0.004 (2)
C7	0.058 (3)	0.043 (3)	0.056 (3)	-0.001 (2)	0.000 (2)	-0.002 (2)
C8	0.056 (3)	0.045 (3)	0.062 (3)	-0.008 (2)	-0.009 (2)	0.014 (3)
C9	0.089 (4)	0.054 (3)	0.085 (4)	0.001 (3)	-0.020 (4)	0.027 (3)

C10	0.051 (3)	0.054 (4)	0.125 (6)	-0.007 (3)	-0.004 (4)	-0.033 (4)
C11	0.060 (4)	0.079 (5)	0.166 (8)	-0.026 (3)	-0.006 (4)	-0.015 (5)
C12	0.040 (3)	0.069 (3)	0.049 (3)	0.004 (3)	-0.003 (2)	0.012 (3)
C13	0.089 (5)	0.139 (6)	0.050 (3)	0.041 (5)	-0.013 (3)	0.006 (4)
C14	0.037 (2)	0.048 (2)	0.040 (2)	-0.006 (2)	-0.001 (2)	-0.0071 (19)
C15	0.061 (3)	0.047 (3)	0.040 (3)	0.000 (2)	-0.001 (2)	-0.004 (2)
C16	0.077 (4)	0.044 (3)	0.046 (3)	-0.002 (2)	-0.004 (3)	0.000 (2)
C17	0.056 (3)	0.043 (3)	0.059 (3)	0.001 (2)	-0.021 (3)	-0.006 (2)
C18	0.048 (3)	0.052 (3)	0.077 (4)	0.008 (2)	0.000 (3)	-0.030 (3)
C19	0.051 (3)	0.055 (3)	0.059 (3)	-0.004 (2)	0.010 (2)	-0.019 (3)
C20	0.090 (4)	0.041 (3)	0.099 (5)	0.003 (3)	-0.032 (4)	-0.003 (3)

Geometric parameters (\AA , $^\circ$)

S1—O8	1.422 (3)	C7—H7A	0.9700
S1—O9	1.427 (3)	C7—H7B	0.9700
S1—O7	1.576 (3)	C8—C9	1.496 (7)
S1—C14	1.753 (5)	C9—H9A	0.9600
C11—C4	1.798 (5)	C9—H9B	0.9600
O1—C5	1.426 (5)	C9—H9C	0.9600
O1—C1	1.437 (5)	C10—C11	1.490 (8)
O2—C8	1.189 (7)	C11—H11A	0.9600
O3—C10	1.337 (6)	C11—H11B	0.9600
O3—C2	1.449 (5)	C11—H11C	0.9600
O4—C10	1.20 (4)	C12—C13	1.488 (8)
O4'—C10	1.222 (16)	C13—H13A	0.9600
O5—C12	1.370 (6)	C13—H13B	0.9600
O5—C3	1.447 (5)	C13—H13C	0.9600
O6—C12	1.184 (6)	C14—C19	1.388 (6)
O7—C6	1.459 (6)	C14—C15	1.393 (6)
O10—C21	1.24 (5)	C15—C16	1.375 (7)
O10—H10	0.8200	C15—H15A	0.9300
C1—C7	1.515 (6)	C16—C17	1.367 (7)
C1—C2	1.530 (6)	C16—H16A	0.9300
C1—H1A	0.9800	C17—C18	1.392 (7)
C2—C3	1.527 (6)	C17—C20	1.503 (8)
C2—H2A	0.9800	C18—C19	1.371 (8)
C3—C4	1.514 (7)	C18—H18A	0.9300
C3—H3A	0.9800	C19—H19A	0.9300
C4—C5	1.521 (6)	C20—H20A	0.9600
C4—H4A	0.9800	C20—H20B	0.9600
C5—C6	1.503 (6)	C20—H20D	0.9600
C5—H5A	0.9800	C21—H21A	0.9600
C6—H6A	0.9700	C21—H21B	0.9600
C6—H6B	0.9700	C21—H21C	0.9600
C7—C8	1.516 (7)		
O8—S1—O9	120.2 (2)	C8—C9—H9B	109.5
O8—S1—O7	104.3 (2)	H9A—C9—H9B	109.5
O9—S1—O7	108.28 (18)	C8—C9—H9C	109.5

supplementary materials

O8—S1—C14	109.2 (2)	H9A—C9—H9C	109.5
O9—S1—C14	108.4 (2)	H9B—C9—H9C	109.5
O7—S1—C14	105.4 (2)	O4—C10—O4'	37.7 (13)
C5—O1—C1	112.4 (3)	O4—C10—O3	116.9 (17)
C10—O3—C2	119.2 (4)	O4'—C10—O3	122.0 (8)
C12—O5—C3	116.3 (4)	O4—C10—C11	121.9 (16)
C6—O7—S1	118.8 (3)	O4'—C10—C11	125.9 (8)
C21—O10—H10	109.5	O3—C10—C11	111.3 (6)
O1—C1—C7	107.6 (4)	C10—C11—H11A	109.5
O1—C1—C2	108.8 (3)	C10—C11—H11B	109.5
C7—C1—C2	112.7 (4)	H11A—C11—H11B	109.5
O1—C1—H1A	109.2	C10—C11—H11C	109.5
C7—C1—H1A	109.2	H11A—C11—H11C	109.5
C2—C1—H1A	109.2	H11B—C11—H11C	109.5
O3—C2—C3	106.9 (3)	O6—C12—O5	122.5 (5)
O3—C2—C1	107.6 (3)	O6—C12—C13	127.3 (5)
C3—C2—C1	109.5 (3)	O5—C12—C13	110.1 (5)
O3—C2—H2A	110.9	C12—C13—H13A	109.5
C3—C2—H2A	110.9	C12—C13—H13B	109.5
C1—C2—H2A	110.9	H13A—C13—H13B	109.5
O5—C3—C4	111.4 (3)	C12—C13—H13C	109.5
O5—C3—C2	108.8 (3)	H13A—C13—H13C	109.5
C4—C3—C2	111.0 (4)	H13B—C13—H13C	109.5
O5—C3—H3A	108.6	C19—C14—C15	120.5 (4)
C4—C3—H3A	108.6	C19—C14—S1	118.9 (4)
C2—C3—H3A	108.6	C15—C14—S1	120.5 (3)
C3—C4—C5	109.5 (4)	C16—C15—C14	118.6 (5)
C3—C4—Cl1	111.1 (3)	C16—C15—H15A	120.7
C5—C4—Cl1	110.6 (3)	C14—C15—H15A	120.7
C3—C4—H4A	108.5	C17—C16—C15	122.6 (5)
C5—C4—H4A	108.5	C17—C16—H16A	118.7
Cl1—C4—H4A	108.5	C15—C16—H16A	118.7
O1—C5—C6	107.8 (4)	C16—C17—C18	117.5 (5)
O1—C5—C4	110.5 (4)	C16—C17—C20	122.7 (6)
C6—C5—C4	112.7 (4)	C18—C17—C20	119.8 (5)
O1—C5—H5A	108.6	C19—C18—C17	122.3 (4)
C6—C5—H5A	108.6	C19—C18—H18A	118.8
C4—C5—H5A	108.6	C17—C18—H18A	118.8
O7—C6—C5	106.9 (4)	C18—C19—C14	118.6 (5)
O7—C6—H6A	110.4	C18—C19—H19A	120.7
C5—C6—H6A	110.4	C14—C19—H19A	120.7
O7—C6—H6B	110.4	C17—C20—H20A	109.5
C5—C6—H6B	110.4	C17—C20—H20B	109.5
H6A—C6—H6B	108.6	H20A—C20—H20B	109.5
C1—C7—C8	115.2 (4)	C17—C20—H20D	109.5
C1—C7—H7A	108.5	H20A—C20—H20D	109.5
C8—C7—H7A	108.5	H20B—C20—H20D	109.5
C1—C7—H7B	108.5	O10—C21—H21A	109.5
C8—C7—H7B	108.5	O10—C21—H21B	109.5

H7A—C7—H7B	107.5	H21A—C21—H21B	109.5
O2—C8—C9	122.3 (5)	O10—C21—H21C	109.5
O2—C8—C7	122.5 (5)	H21A—C21—H21C	109.5
C9—C8—C7	115.2 (5)	H21B—C21—H21C	109.5
C8—C9—H9A	109.5		
O8—S1—O7—C6	167.7 (4)	O1—C5—C6—O7	74.9 (5)
O9—S1—O7—C6	38.6 (4)	C4—C5—C6—O7	-162.9 (4)
C14—S1—O7—C6	-77.2 (4)	O1—C1—C7—C8	-81.7 (5)
C5—O1—C1—C7	174.6 (4)	C2—C1—C7—C8	158.4 (4)
C5—O1—C1—C2	-63.0 (5)	C1—C7—C8—O2	1.7 (8)
C10—O3—C2—C3	-121.1 (5)	C1—C7—C8—C9	-177.0 (5)
C10—O3—C2—C1	121.3 (5)	C2—O3—C10—O4	-31 (2)
O1—C1—C2—O3	173.3 (3)	C2—O3—C10—O4'	12 (2)
C7—C1—C2—O3	-67.5 (5)	C2—O3—C10—C11	-177.5 (5)
O1—C1—C2—C3	57.5 (4)	C3—O5—C12—O6	0.8 (7)
C7—C1—C2—C3	176.7 (4)	C3—O5—C12—C13	179.5 (5)
C12—O5—C3—C4	96.5 (4)	O8—S1—C14—C19	25.0 (5)
C12—O5—C3—C2	-140.9 (4)	O9—S1—C14—C19	157.6 (4)
O3—C2—C3—O5	66.7 (4)	O7—S1—C14—C19	-86.6 (4)
C1—C2—C3—O5	-177.1 (3)	O8—S1—C14—C15	-151.0 (4)
O3—C2—C3—C4	-170.5 (3)	O9—S1—C14—C15	-18.4 (4)
C1—C2—C3—C4	-54.3 (5)	O7—S1—C14—C15	97.4 (4)
O5—C3—C4—C5	174.4 (4)	C19—C14—C15—C16	-0.5 (7)
C2—C3—C4—C5	53.1 (5)	S1—C14—C15—C16	175.4 (4)
O5—C3—C4—Cl1	52.0 (5)	C14—C15—C16—C17	0.0 (8)
C2—C3—C4—Cl1	-69.4 (4)	C15—C16—C17—C18	1.1 (8)
C1—O1—C5—C6	-173.7 (4)	C15—C16—C17—C20	-177.5 (5)
C1—O1—C5—C4	62.8 (5)	C16—C17—C18—C19	-1.8 (8)
C3—C4—C5—O1	-56.3 (5)	C20—C17—C18—C19	176.8 (5)
Cl1—C4—C5—O1	66.5 (4)	C17—C18—C19—C14	1.3 (8)
C3—C4—C5—C6	-176.9 (4)	C15—C14—C19—C18	-0.2 (7)
Cl1—C4—C5—C6	-54.2 (5)	S1—C14—C19—C18	-176.2 (4)
S1—O7—C6—C5	-174.1 (3)		

supplementary materials

Fig. 1

