

3-Chlorophenyl benzoate

B. Thimme Gowda,^{a*} Sabine Foro,^b K. S. Babitha^a and Hartmut Fuess^b

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 9 July 2008; accepted 19 July 2008

Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.004 \text{ \AA}$; R factor = 0.047; wR factor = 0.135; data-to-parameter ratio = 13.3.

The C=O group in the title compound, $C_{13}H_9ClO_2$, is *syn* to the chloro group. The two aromatic rings are twisted by $56.88(6)^\circ$. Adjacent molecules are linked via weak C—H···O hydrogen bonding into a linear chain.

Related literature

For previous studies, see: Gowda *et al.* (2007a,b,c); Nayak & Gowda (2008).



Experimental

Crystal data

$C_{13}H_9ClO_2$	$\gamma = 93.25(1)^\circ$
$M_r = 232.65$	$V = 549.89(13) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.0734(6) \text{ \AA}$	Cu $K\alpha$ radiation
$b = 8.389(1) \text{ \AA}$	$\mu = 2.92 \text{ mm}^{-1}$
$c = 11.747(2) \text{ \AA}$	$T = 299(2) \text{ K}$
$\alpha = 107.89(1)^\circ$	$0.60 \times 0.55 \times 0.50 \text{ mm}$
$\beta = 102.98(1)^\circ$	

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.197$, $T_{\max} = 0.233$
2143 measured reflections

1947 independent reflections
1872 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.086$
3 standard reflections
frequency: 120 min
intensity decay: 1.0%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.135$
 $S = 1.10$
1947 reflections

146 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C6—H6···O2 ⁱ	0.93	2.46	3.319 (3)	154

Symmetry code: (i) $x - 1, y, z$.

Data collection: *CAD-4-PC* (Enraf–Nonius, 1996); cell refinement: *CAD-4-PC*; data reduction: *REDU4* (Stoe & Cie, 1987); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97*.

BTG thanks the Alexander von Humboldt Foundation, Bonn, Germany, for extensions of his research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2472).

References

- Enraf–Nonius (1996). *CAD-4-PC*. Enraf–Nonius, Delft, The Netherlands.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2007a). *Acta Cryst. E63*, o3756.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2007b). *Acta Cryst. E63*, o3801.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2007c). *Acta Cryst. E63*, o4286.
- Nayak, R. & Gowda, B. T. (2008). *Z. Naturforsch. Teil A*, **63**. In the press.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A24*, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.
- Stoe & Cie (1987). *REDU4*. Stoe & Cie GmbH, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2008). E64, o1587 [doi:10.1107/S1600536808022721]

3-Chlorophenyl benzoate

B. T. Gowda, S. Foro, K. S. Babitha and H. Fuess

Comment

As part of a study of the substituent effects on the structures of aryl benzoates (Gowda *et al.*, 2007*a, b, c*), in the present work, the structure of 3-chlorophenyl benzoate (3CPBA) has been determined. The conformation of the C=O bond in 3CPBA is *syn* to the *meta*-chloro group in the phenolic benzene ring (Fig. 1), in contrast to the *anti* conformations of the C=O bond and the *meta*-methyl group in 3-methylphenyl benzoate (3MePBA) (Gowda *et al.*, 2007*a*). The bond parameters in 3CPBA are similar to those of 3MePBA (Gowda *et al.*, 2007*a*), 2,3-dichlorophenyl benzoate (23DCPBA) (Gowda *et al.*, 2007*c*), 3,4-dichlorophenyl benzoate (34DCPBA) (Gowda *et al.*, 2007*b*) and other aryl benzoates (Gowda *et al.*, 2007*a, b, c*). The dihedral angle between the benzene and benzoyl rings in 3CPBA is 56.88 (6) $^{\circ}$, compared to the values of 79.61 (6) $^{\circ}$ in 3MePBA, 50.16 (7) $^{\circ}$ in 23DCPBA and 53.77 (5) $^{\circ}$ in 34DCPBA. The packing diagram of the crystal structure in which the molecules are connected *via* intermolecular C—H—O hydrogen bonds (Table 1) is shown in Fig. 2.

Experimental

The title compound was prepared according to the method of Nayak & Gowda (2008). The purity of the compound was checked by determining its melting point. It was characterized by recording its infrared and NMR spectra (Nayak & Gowda, 2008). The single crystals used in X-ray diffraction studies were obtained by the slow evaporation of an ethanolic solution of the title compound at room temperature.

Refinement

All H atoms were included in the riding-model approximation with C—H = 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2 \times U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of the title compound, showing the atom labeling scheme. The displacement ellipsoids are drawn at the 50% probability level. The H atoms are represented as small spheres of arbitrary radii.

supplementary materials




Fig. 2. Molecular packing of the title compound.

3-Chlorophenyl benzoate

Crystal data

$C_{13}H_9ClO_2$	$Z = 2$
$M_r = 232.65$	$F_{000} = 240$
Triclinic, $P\bar{1}$	$D_x = 1.405 \text{ Mg m}^{-3}$
Hall symbol: -P 1	$\text{Cu } K\alpha \text{ radiation}$
$a = 6.0734 (6) \text{ \AA}$	$\lambda = 1.54180 \text{ \AA}$
$b = 8.389 (1) \text{ \AA}$	Cell parameters from 25 reflections
$c = 11.747 (2) \text{ \AA}$	$\theta = 5.6\text{--}31.7^\circ$
$\alpha = 107.89 (1)^\circ$	$\mu = 2.92 \text{ mm}^{-1}$
$\beta = 102.98 (1)^\circ$	$T = 299 (2) \text{ K}$
$\gamma = 93.25 (1)^\circ$	Prism, colourless
$V = 549.89 (13) \text{ \AA}^3$	$0.60 \times 0.55 \times 0.50 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.086$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 66.9^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 4.1^\circ$
$T = 299(2) \text{ K}$	$h = -7 \rightarrow 1$
$\omega/2\theta$ scans	$k = -9 \rightarrow 9$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -13 \rightarrow 14$
$T_{\text{min}} = 0.197$, $T_{\text{max}} = 0.233$	3 standard reflections
2143 measured reflections	every 120 min
1947 independent reflections	intensity decay: 1.0%
1872 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.046$	$w = 1/[\sigma^2(F_o^2) + (0.0771P)^2 + 0.1623P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.135$	$(\Delta/\sigma)_{\max} = 0.002$
$S = 1.10$	$\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
1947 reflections	$\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$
146 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.149 (8)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.29134 (10)	0.17688 (7)	-0.06457 (6)	0.0678 (3)
O1	0.2384 (2)	0.74790 (18)	0.23750 (15)	0.0565 (4)
O2	0.5777 (2)	0.70473 (19)	0.33852 (15)	0.0587 (4)
C1	0.1614 (3)	0.5746 (2)	0.18489 (18)	0.0474 (5)
C2	0.2647 (3)	0.4730 (2)	0.10218 (19)	0.0477 (5)
H2	0.3962	0.5149	0.0869	0.057*
C3	0.1670 (3)	0.3070 (3)	0.04263 (19)	0.0482 (5)
C4	-0.0281 (3)	0.2436 (3)	0.0636 (2)	0.0546 (5)
H4	-0.0937	0.1321	0.0212	0.065*
C5	-0.1247 (4)	0.3482 (3)	0.1485 (2)	0.0593 (6)
H5	-0.2549	0.3060	0.1647	0.071*
C6	-0.0314 (3)	0.5149 (3)	0.2099 (2)	0.0555 (5)
H6	-0.0977	0.5853	0.2669	0.067*
C7	0.4554 (3)	0.7994 (3)	0.30835 (17)	0.0458 (5)
C8	0.5176 (3)	0.9840 (3)	0.34208 (17)	0.0457 (5)
C9	0.3622 (4)	1.0902 (3)	0.3149 (2)	0.0568 (5)
H9	0.2109	1.0464	0.2740	0.068*
C10	0.4331 (5)	1.2610 (3)	0.3487 (2)	0.0680 (7)
H10	0.3287	1.3323	0.3307	0.082*
C11	0.6556 (5)	1.3268 (3)	0.4086 (2)	0.0692 (7)
H11	0.7018	1.4423	0.4315	0.083*
C12	0.8107 (5)	1.2214 (3)	0.4347 (2)	0.0717 (7)

supplementary materials

H12	0.9619	1.2658	0.4752	0.086*
C13	0.7428 (4)	1.0512 (3)	0.4013 (2)	0.0602 (6)
H13	0.8486	0.9804	0.4184	0.072*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0704 (4)	0.0528 (4)	0.0800 (5)	0.0084 (3)	0.0358 (3)	0.0095 (3)
O1	0.0472 (8)	0.0453 (8)	0.0692 (9)	0.0134 (6)	0.0057 (6)	0.0131 (7)
O2	0.0543 (8)	0.0553 (9)	0.0681 (10)	0.0198 (7)	0.0096 (7)	0.0247 (7)
C1	0.0436 (9)	0.0457 (11)	0.0534 (11)	0.0114 (8)	0.0092 (8)	0.0183 (9)
C2	0.0429 (10)	0.0475 (11)	0.0581 (11)	0.0066 (8)	0.0177 (8)	0.0215 (9)
C3	0.0471 (10)	0.0478 (11)	0.0560 (11)	0.0098 (8)	0.0179 (8)	0.0218 (9)
C4	0.0502 (11)	0.0492 (11)	0.0667 (13)	0.0021 (8)	0.0140 (9)	0.0241 (10)
C5	0.0447 (11)	0.0721 (15)	0.0711 (14)	0.0048 (9)	0.0210 (9)	0.0335 (12)
C6	0.0464 (10)	0.0671 (14)	0.0574 (12)	0.0162 (9)	0.0192 (9)	0.0211 (10)
C7	0.0452 (10)	0.0506 (11)	0.0456 (10)	0.0152 (8)	0.0148 (8)	0.0176 (8)
C8	0.0498 (10)	0.0493 (11)	0.0412 (9)	0.0133 (8)	0.0146 (8)	0.0161 (8)
C9	0.0574 (12)	0.0549 (13)	0.0593 (12)	0.0174 (9)	0.0124 (9)	0.0202 (10)
C10	0.0885 (17)	0.0525 (13)	0.0701 (15)	0.0255 (12)	0.0225 (12)	0.0254 (11)
C11	0.0935 (18)	0.0520 (13)	0.0634 (14)	0.0038 (12)	0.0272 (12)	0.0169 (11)
C12	0.0675 (15)	0.0653 (15)	0.0712 (15)	-0.0050 (11)	0.0129 (11)	0.0128 (12)
C13	0.0534 (12)	0.0594 (13)	0.0630 (13)	0.0109 (10)	0.0076 (9)	0.0177 (11)

Geometric parameters (\AA , $^\circ$)

Cl1—C3	1.739 (2)	C6—H6	0.9300
O1—C7	1.356 (2)	C7—C8	1.479 (3)
O1—C1	1.399 (2)	C8—C13	1.384 (3)
O2—C7	1.195 (2)	C8—C9	1.387 (3)
C1—C2	1.374 (3)	C9—C10	1.378 (3)
C1—C6	1.374 (3)	C9—H9	0.9300
C2—C3	1.380 (3)	C10—C11	1.370 (4)
C2—H2	0.9300	C10—H10	0.9300
C3—C4	1.374 (3)	C11—C12	1.376 (4)
C4—C5	1.376 (3)	C11—H11	0.9300
C4—H4	0.9300	C12—C13	1.371 (4)
C5—C6	1.379 (3)	C12—H12	0.9300
C5—H5	0.9300	C13—H13	0.9300
C7—O1—C1	118.81 (14)	O2—C7—C8	125.42 (19)
C2—C1—C6	122.04 (19)	O1—C7—C8	111.50 (15)
C2—C1—O1	120.12 (18)	C13—C8—C9	119.3 (2)
C6—C1—O1	117.55 (18)	C13—C8—C7	117.74 (18)
C1—C2—C3	117.83 (17)	C9—C8—C7	122.95 (18)
C1—C2—H2	121.1	C10—C9—C8	119.6 (2)
C3—C2—H2	121.1	C10—C9—H9	120.2
C4—C3—C2	121.76 (19)	C8—C9—H9	120.2
C4—C3—Cl1	119.48 (17)	C11—C10—C9	120.7 (2)

C2—C3—Cl1	118.73 (14)	C11—C10—H10	119.7
C3—C4—C5	118.8 (2)	C9—C10—H10	119.7
C3—C4—H4	120.6	C10—C11—C12	119.8 (2)
C5—C4—H4	120.6	C10—C11—H11	120.1
C4—C5—C6	120.99 (19)	C12—C11—H11	120.1
C4—C5—H5	119.5	C13—C12—C11	120.2 (2)
C6—C5—H5	119.5	C13—C12—H12	119.9
C1—C6—C5	118.6 (2)	C11—C12—H12	119.9
C1—C6—H6	120.7	C12—C13—C8	120.4 (2)
C5—C6—H6	120.7	C12—C13—H13	119.8
O2—C7—O1	123.08 (19)	C8—C13—H13	119.8
C7—O1—C1—C2	-61.9 (2)	C1—O1—C7—C8	172.24 (16)
C7—O1—C1—C6	124.2 (2)	O2—C7—C8—C13	7.9 (3)
C6—C1—C2—C3	0.7 (3)	O1—C7—C8—C13	-172.14 (18)
O1—C1—C2—C3	-172.97 (16)	O2—C7—C8—C9	-173.4 (2)
C1—C2—C3—C4	0.5 (3)	O1—C7—C8—C9	6.6 (3)
C1—C2—C3—Cl1	178.84 (14)	C13—C8—C9—C10	-1.0 (3)
C2—C3—C4—C5	-1.5 (3)	C7—C8—C9—C10	-179.7 (2)
Cl1—C3—C4—C5	-179.86 (16)	C8—C9—C10—C11	0.2 (4)
C3—C4—C5—C6	1.4 (3)	C9—C10—C11—C12	0.4 (4)
C2—C1—C6—C5	-0.8 (3)	C10—C11—C12—C13	-0.2 (4)
O1—C1—C6—C5	173.01 (18)	C11—C12—C13—C8	-0.6 (4)
C4—C5—C6—C1	-0.3 (3)	C9—C8—C13—C12	1.2 (3)
C1—O1—C7—O2	-7.8 (3)	C7—C8—C13—C12	180.0 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C6—H6···O2 ⁱ	0.93	2.46	3.319 (3)	154

Symmetry codes: (i) $x-1, y, z$.

supplementary materials

Fig. 1




Fig. 2

