

6,6'-Dihydroxy-2,2'-[(pentane-1,5-diyl-dioxy)bis(nitrilomethylidene)]diphenol

Wen-Kui Dong,* Xue-Ni He, Yong-Hong Guan, Li Xu and Zong-Li Ren

School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, People's Republic of China
Correspondence e-mail: dongwk@mail.lzjtu.cn


Received 18 July 2008; accepted 22 July 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.050; wR factor = 0.135; data-to-parameter ratio = 13.2.

The molecule of the title compound, $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_6$, assumes a W-shaped configuration with the dihedral angle between the two halves of the molecule being 82.48 (5)°. There is one half-molecule in the asymmetric unit with a crystallographic twofold rotation axis passing through the central C atom of the five methylene groups in the $[-\text{CH}=\text{N}-\text{O}-(\text{CH}_2)_5-\text{O}-\text{N}=\text{CH}-]$ bridge. The dihedral angle formed by the two benzene rings in each molecule of the title compound is 84.18 (4)°. There are strong intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds and weak intermolecular $\pi-\pi$ stacking interactions between neighbouring benzene rings, and the intermolecular plane-to-plane distances are 3.488 (2) and 3.841 (3) Å along the b and c axes, respectively. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link each molecule to two others, forming an infinite three-dimensional supramolecular structure.

Related literature

For related literature, see: Akine *et al.* (2001, 2005, 2006); Atwood (1997); Dong & Feng (2006); Dong, Zhao *et al.* (2008); Dong, He *et al.* (2008); Duan *et al.* (2007); Venkataramanan *et al.* (2005); Yu *et al.* (2008).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_6$
 $M_r = 374.39$
 Monoclinic, $C2/c$
 $a = 28.439$ (3) Å
 $b = 4.6997$ (6) Å
 $c = 14.0843$ (17) Å
 $\beta = 100.354$ (2)°

$V = 1851.8$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 298$ (2) K
 $0.46 \times 0.27 \times 0.25$ mm

Data collection

Siemens SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.955$, $T_{\max} = 0.975$

4246 measured reflections
 1621 independent reflections
 837 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.062$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.135$
 $S = 1.00$
 1621 reflections

123 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\cdots\text{N}1$	0.82	1.92	2.630 (3)	144
$\text{O}3-\text{H}3\cdots\text{O}2$	0.82	2.24	2.689 (3)	115
$\text{O}3-\text{H}3\cdots\text{O}1^i$	0.82	2.29	2.958 (3)	139

Symmetry code: (i) $x, -y, z - \frac{1}{2}$.

Data collection: SMART (Siemens, 1996); cell refinement: SMART; data reduction: SAINT (Siemens, 1996); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the Foundation of the Education Department of Gansu Province (grant No. 0604-01) and the 'Qing Lan' Talent Engineering Funds of Lanzhou Jiaotong University (grant No. QL-03-01A), which are gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2427).

References

- Akine, S., Dong, W. K. & Nabeshima, T. (2006). *Inorg. Chem.* **45**, 4677–4684.
 Akine, S., Takanori, T., Dong, W. K. & Nabeshima, T. (2005). *J. Org. Chem.* **70**, 1704–1711.
 Akine, S., Taniguchi, T. & Nabeshima, T. (2001). *Chem. Lett.* **30**, 682–683.
 Atwood, D. A. (1997). *Coord. Chem. Rev.* **165**, 267–296.
 Dong, W.-K. & Feng, J.-H. (2006). *Acta Cryst.* **E62**, o3577–o3578.
 Dong, W.-K., He, X.-N., Li, L., Lv, Z.-W. & Tong, J.-F. (2008). *Acta Cryst.* **E64**, o1405.
 Dong, W.-K., Zhao, C.-Y., Zhong, J.-K., Tang, X.-L. & Yu, T.-Z. (2008). *Acta Cryst.* **E64**, o1323.
 Duan, J.-G., Dong, C.-M., Shi, J.-Y., Wu, L. & Dong, W.-K. (2007). *Acta Cryst.* **E63**, o2704–o2705.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Venkataramanan, N. S., Kuppuraj, G. & Rajagopal, S. (2005). *Coord. Chem. Rev.* **249**, 1249–1268.

Yu, T. Z., Zhang, K., Zhao, Y. L., Yang, C. H., Zhang, H., Qian, L., Fan, D. W., Dong, W. K., Chen, L. L. & Qiu, Y. Q. (2008). *Inorg. Chim. Acta*, **361**, 233–240.

supplementary materials

Acta Cryst. (2008). E64, o1600-o1601 [doi:10.1107/S1600536808023064]

6,6'-Dihydroxy-2,2'-[(pentane-1,5-diylidioxy)bis(nitrilomethylidyne)]diphenol

W.-K. Dong, X.-N. He, Y.-H. Guan, L. Xu and Z.-L. Ren

Comment

Salen-type compounds are one of most versatile mixed-donor ligands in the field of coordination chemistry. There has been growing interest in salen-type ligands, mainly because of their wide application in the field of synthesis, biochemistry, photochemistry and catalysis (Akine *et al.*, 2006; Atwood, 1997; Yu *et al.*, 2008; Venkataramanan *et al.*, 2005). Many salen-type complexes have been structurally characterized (Akine *et al.*, 2006; Yu *et al.*, 2008), but only a relatively small number of free salen-type compounds have been characterized (Akine *et al.*, 2001). As an extension of our work (Dong & Feng, 2006; Duan *et al.*, 2007; Dong, Zhao *et al.*, 2008; Akine *et al.*, 2005) on the structural characterization of salen-type bisoxime compounds, the title compound, (Fig. 1), is reported here.

The molecule assumes a W shape with the dihedral angle between the two halves of the molecule 82.48 (5)°. There is 1/2 molecule per asymmetric unit with a crystallographic twofold rotation axis passing through the central carbon (symmetry code: $-x, y, 1/2 - z$) of the five carbon atoms in the ($-\text{CH}=\text{N}-\text{O}-(\text{CH}_2)_5-\text{O}-\text{N}=\text{CH}-$) bridge. This structure is similar to what was observed in our previously reported salen-type bisoxime compound (Duan *et al.*, 2007). The dihedral angle formed by the two benzene rings in each molecule of the title compound is 84.18 (4)°. There are strong intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds and weak intermolecular $\pi-\pi$ stacking interactions between the neighbouring benzene rings, and the inter-molecular plane-to-plane distances are 3.488 (2) and 3.841 (3) Å along *b* and *c* axis, respectively. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link each molecule to 2 others into infinite three-dimensional supramolecular structure, which is the crystal structure firstly reported of salen-type bisoxime compounds containing pentamethene bridge.

Experimental

6,6'-Dihydroxy-2,2'-[(pentane-1,5-diylidioxy)bis(nitrilomethylidyne)]diphenol was synthesized according to an analogous method reported earlier (Dong & Feng, 2006; Dong, He *et al.*, 2008). To an ethanol solution (3 ml) of 2,3-dihydroxybenzaldehyde (138.4 mg, 1.0 mmol) was added an ethanol solution (2 ml) of 1,5-bis(aminoxy)pentane (67.4 mg, 0.5 mmol). The reaction mixture was stirred at 328 K for 8 h. After cooling to room temperature, the formed precipitate was separated by filtration, and washed successively with ethanol and ethanol-hexane (1:4), respectively. The product was dried under vacuum to yield 109.3 mg of the title compound. Yield, 58.4%. mp. 408.5–409.5 K. Anal. Calc. for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_6$: C, 60.95; H, 5.92; N, 7.48. Found: C, 60.75; H, 5.99; N, 7.42.

Pale-brown needle-like single crystals suitable for X-ray diffraction studies were obtained after two weeks by slow evaporation from an ethanol-chloroform mixed solution of the title compound.

Refinement

Non-H atoms were refined anisotropically. H atoms were treated as riding atoms with distances $\text{C}-\text{H} = 0.97$ (CH_2), or 0.93 Å (CH), $\text{O}-\text{H} = 0.82$ Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the title compound with atom numbering scheme [Symmetry codes: $-x + 1, y, -z + 3/2$]. Displacement ellipsoids for non-hydrogen atoms are drawn at the 30% probability level.



Fig. 2. The W shape configuration of the title compound.




Fig. 3. Part of the supramolecular structure of the title compound along b axis. Intra- and intermolecular hydrogen bonds are shown as dashed lines.

6,6'-Dihydroxy-2,2'-[(pentane-1,5-dioldioxy)bis(nitrilomethylidene)]diphenol

Crystal data

$C_{19}H_{22}N_2O_6$

$M_r = 374.39$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 28.439 (3) \text{ \AA}$

$b = 4.6997 (6) \text{ \AA}$

$c = 14.0843 (17) \text{ \AA}$

$\beta = 100.354 (2)^\circ$

$V = 1851.8 (4) \text{ \AA}^3$

$Z = 4$

$F_{000} = 792$

$D_x = 1.343 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 875 reflections

$\theta = 2.9\text{--}22.4^\circ$

$\mu = 0.10 \text{ mm}^{-1}$

$T = 298 (2) \text{ K}$

Needle-like, pale-brown

$0.46 \times 0.27 \times 0.25 \text{ mm}$

Data collection

Siemens SMART 1000 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298(2) \text{ K}$

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.955, T_{\max} = 0.975$

4246 measured reflections

1621 independent reflections

837 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.062$

$\theta_{\max} = 25.0^\circ$

$\theta_{\min} = 1.5^\circ$

$h = -33 \rightarrow 30$

$k = -5 \rightarrow 5$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.049$	H-atom parameters constrained
$wR(F^2) = 0.135$	$w = 1/[\sigma^2(F_o^2) + (0.0407P)^2 + 1.5785P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
1621 reflections	$(\Delta/\sigma)_{\max} < 0.001$
123 parameters	$\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
N1	0.40208 (8)	-0.0702 (5)	0.51037 (15)	0.0479 (7)	
O1	0.41975 (7)	-0.2344 (4)	0.59236 (12)	0.0542 (6)	
O2	0.39611 (7)	0.1384 (5)	0.33535 (13)	0.0637 (7)	
H2	0.4076	0.0385	0.3815	0.095*	
O3	0.34987 (8)	0.4908 (5)	0.19941 (14)	0.0782 (8)	
H3	0.3730	0.3870	0.1991	0.117*	
C1	0.46067 (10)	-0.3889 (7)	0.57460 (19)	0.0535 (9)	
H1A	0.4849	-0.2576	0.5610	0.064*	
H1B	0.4518	-0.5126	0.5192	0.064*	
C2	0.47987 (11)	-0.5624 (7)	0.66243 (18)	0.0543 (9)	
H2A	0.5048	-0.6868	0.6478	0.065*	
H2B	0.4544	-0.6814	0.6780	0.065*	
C3	0.5000	-0.3833 (9)	0.7500	0.0500 (11)	
H3A	0.4749	-0.2617	0.7657	0.060*	0.50
H3B	0.5251	-0.2617	0.7343	0.060*	0.50
C4	0.36573 (10)	0.0763 (7)	0.52087 (19)	0.0470 (8)	
H4	0.3540	0.0644	0.5783	0.056*	
C5	0.34239 (9)	0.2611 (6)	0.44479 (18)	0.0418 (7)	

supplementary materials

C6	0.35833 (10)	0.2865 (6)	0.35701 (19)	0.0439 (7)
C7	0.33496 (11)	0.4666 (7)	0.28589 (19)	0.0509 (8)
C8	0.29684 (11)	0.6252 (7)	0.3023 (2)	0.0573 (9)
H8	0.2819	0.7498	0.2550	0.069*
C9	0.28046 (11)	0.6018 (7)	0.3883 (2)	0.0590 (9)
H9	0.2542	0.7075	0.3985	0.071*
C10	0.30300 (10)	0.4222 (7)	0.4586 (2)	0.0538 (9)
H10	0.2918	0.4075	0.5165	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0471 (15)	0.0517 (18)	0.0431 (13)	0.0017 (13)	0.0029 (11)	0.0029 (12)
O1	0.0510 (13)	0.0670 (16)	0.0443 (11)	0.0114 (12)	0.0076 (9)	0.0072 (10)
O2	0.0557 (14)	0.0800 (17)	0.0593 (13)	0.0191 (13)	0.0211 (10)	0.0113 (11)
O3	0.0804 (16)	0.099 (2)	0.0575 (13)	0.0188 (15)	0.0195 (12)	0.0206 (13)
C1	0.0520 (19)	0.056 (2)	0.0516 (17)	0.0052 (17)	0.0074 (15)	-0.0072 (15)
C2	0.058 (2)	0.049 (2)	0.0529 (17)	0.0074 (17)	0.0018 (15)	-0.0022 (15)
C3	0.044 (2)	0.049 (3)	0.055 (2)	0.000	0.0043 (19)	0.000
C4	0.0450 (18)	0.056 (2)	0.0408 (15)	-0.0017 (16)	0.0093 (13)	-0.0038 (15)
C5	0.0342 (16)	0.045 (2)	0.0447 (15)	-0.0038 (15)	0.0033 (13)	-0.0061 (13)
C6	0.0365 (16)	0.046 (2)	0.0498 (16)	-0.0021 (15)	0.0093 (13)	-0.0063 (14)
C7	0.053 (2)	0.055 (2)	0.0440 (16)	-0.0018 (17)	0.0059 (14)	0.0002 (15)
C8	0.054 (2)	0.055 (2)	0.0579 (19)	0.0044 (18)	-0.0037 (16)	0.0007 (16)
C9	0.0465 (19)	0.063 (2)	0.066 (2)	0.0112 (17)	0.0040 (16)	-0.0084 (18)
C10	0.0471 (19)	0.062 (2)	0.0521 (17)	0.0012 (17)	0.0099 (14)	-0.0109 (16)

Geometric parameters (\AA , $^\circ$)

N1—C4	1.273 (3)	C3—H3A	0.9700
N1—O1	1.405 (3)	C3—H3B	0.9700
O1—C1	1.431 (3)	C4—C5	1.444 (4)
O2—C6	1.360 (3)	C4—H4	0.9300
O2—H2	0.8200	C5—C10	1.394 (4)
O3—C7	1.364 (3)	C5—C6	1.396 (3)
O3—H3	0.8200	C6—C7	1.386 (4)
C1—C2	1.500 (4)	C7—C8	1.369 (4)
C1—H1A	0.9700	C8—C9	1.378 (4)
C1—H1B	0.9700	C8—H8	0.9300
C2—C3	1.518 (4)	C9—C10	1.370 (4)
C2—H2A	0.9700	C9—H9	0.9300
C2—H2B	0.9700	C10—H10	0.9300
C3—C2 ⁱ	1.518 (4)		
C4—N1—O1	112.3 (2)	N1—C4—C5	120.9 (3)
N1—O1—C1	108.47 (19)	N1—C4—H4	119.5
C6—O2—H2	109.5	C5—C4—H4	119.5
C7—O3—H3	109.5	C10—C5—C6	118.3 (3)
O1—C1—C2	108.6 (2)	C10—C5—C4	119.8 (3)

O1—C1—H1A	110.0	C6—C5—C4	122.0 (3)
C2—C1—H1A	110.0	O2—C6—C7	116.5 (3)
O1—C1—H1B	110.0	O2—C6—C5	123.4 (3)
C2—C1—H1B	110.0	C7—C6—C5	120.2 (3)
H1A—C1—H1B	108.4	O3—C7—C8	119.2 (3)
C1—C2—C3	113.4 (3)	O3—C7—C6	120.7 (3)
C1—C2—H2A	108.9	C8—C7—C6	120.1 (3)
C3—C2—H2A	108.9	C7—C8—C9	120.5 (3)
C1—C2—H2B	108.9	C7—C8—H8	119.8
C3—C2—H2B	108.9	C9—C8—H8	119.8
H2A—C2—H2B	107.7	C10—C9—C8	119.8 (3)
C2—C3—C2 ⁱ	112.6 (4)	C10—C9—H9	120.1
C2—C3—H3A	109.1	C8—C9—H9	120.1
C2 ⁱ —C3—H3A	109.1	C9—C10—C5	121.2 (3)
C2—C3—H3B	109.1	C9—C10—H10	119.4
C2 ⁱ —C3—H3B	109.1	C5—C10—H10	119.4
H3A—C3—H3B	107.8		
C4—N1—O1—C1	-179.4 (2)	O2—C6—C7—O3	-0.2 (4)
N1—O1—C1—C2	179.5 (2)	C5—C6—C7—O3	-179.2 (3)
O1—C1—C2—C3	-66.3 (3)	O2—C6—C7—C8	-179.4 (3)
C1—C2—C3—C2 ⁱ	-178.9 (3)	C5—C6—C7—C8	1.5 (4)
O1—N1—C4—C5	179.6 (2)	O3—C7—C8—C9	178.9 (3)
N1—C4—C5—C10	-179.2 (3)	C6—C7—C8—C9	-1.9 (5)
N1—C4—C5—C6	0.2 (4)	C7—C8—C9—C10	1.2 (5)
C10—C5—C6—O2	-179.5 (2)	C8—C9—C10—C5	-0.1 (5)
C4—C5—C6—O2	1.1 (4)	C6—C5—C10—C9	-0.2 (4)
C10—C5—C6—C7	-0.5 (4)	C4—C5—C10—C9	179.2 (3)
C4—C5—C6—C7	-179.9 (3)		

Symmetry codes: (i) $-x+1, y, -z+3/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots N1	0.82	1.92	2.630 (3)	144
O3—H3 \cdots O2	0.82	2.24	2.689 (3)	115
O3—H3 \cdots O1 ⁱⁱ	0.82	2.29	2.958 (3)	139

Symmetry codes: (ii) $x, -y, z-1/2$.

Fig. 1




Fig. 2




Fig. 3

