

## 3-Benzylsulfanyl-5-(4-phenyl-1*H*-1,2,3-triazol-1-ylmethyl)-4*H*-1,2,4-triazol-4-amine

Qing-Zhu Chu, Huan-Ran Zhou and Xiao-Ru Zhang\*

College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, 266042 Qingdao, Shandong, People's Republic of China  
Correspondence e-mail: qustchemistry@126.com


Received 14 July 2008; accepted 23 July 2008

Key indicators: single-crystal X-ray study;  $T = 187\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$ ;  $R$  factor = 0.079;  $wR$  factor = 0.183; data-to-parameter ratio = 12.1.

The molecule of the title compound,  $\text{C}_{18}\text{H}_{17}\text{N}_7\text{S}$ , is non-planar, with a dihedral angle of  $71.4(4)^\circ$  between the two triazole rings, and an angle of  $15.5(3)^\circ$  between the two phenyl rings. An intramolecular  $\text{N}-\text{H}\cdots\text{S}$  hydrogen bond forms a five-membered ring.

### Related literature

For related literature, see: Allen *et al.* (1987); Barchiesi *et al.* (2000); Colanceska-Ragenovic *et al.* (2001); Kaplancikli *et al.* (2008); Khanum *et al.* (2005); Rodriguez-Fernandez *et al.* (2005); Zhang *et al.* (2008).



### Experimental

#### Crystal data

$\text{C}_{18}\text{H}_{17}\text{N}_7\text{S}$

$M_r = 363.45$

Orthorhombic,  $Pna2_1$

$a = 8.0487(15)\text{ \AA}$

$b = 5.4689(10)\text{ \AA}$

$c = 38.721(7)\text{ \AA}$

$V = 1704.4(5)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.21\text{ mm}^{-1}$

$T = 186.5(2)\text{ K}$   
 $0.35 \times 0.25 \times 0.04\text{ mm}$

#### Data collection

Siemens SMART 1000 CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.931$ ,  $T_{\max} = 0.992$

7815 measured reflections  
2854 independent reflections  
2592 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.052$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.078$   
 $wR(F^2) = 0.183$   
 $S = 1.20$   
2854 reflections  
235 parameters  
1 restraint

H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.48\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.39\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
with 2854 Freidel pairs  
Flack parameter: 0.05 (19)

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N7—H7B $\cdots$ S1	0.86	2.77	3.116 (6)	105

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2003).

This project was supported by the Natural Science Foundation of Shandong Province (grant No. Y2006B07).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2597).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Barchiesi, F., Arzeni, D., Fothergill, A. W., Falconi di Francesco, L., Caselli, F., Rinaldi, M. G. & Scalise, G. (2000). *Antimicrob. Agents Chemother.* **44**, 226–229.
- Colanceska-Ragenovic, K., Dimova, V., Kakurinov, V., Gabor Molnar, D. & Buzarovska, A. (2001). *Molecules*, **6**, 815–824.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Kaplancikli, Z. A., Turan-Zitouni, G., Özdemir, A. & Revial, G. (2008). *Eur. J. Med. Chem.* **43**, 155–159.
- Khanum, S. A., Shashikanth, S., Umesha, S. & Kavitha, R. (2005). *Eur. J. Med. Chem.* **40**, 1156–1162.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Rodriguez-Fernandez, E., Manzano, J. L., Benito, J. J., Hermosa, R., Monte, E. & Criado, J. J. (2005). *J. Inorg. Biochem.* **99**, 1558–1572.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments, Inc., Madison, Wisconsin, USA.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Zhang, X. R., Xu, M. H. & Zhang, S. S. (2008). *J. Chin. Chem. Soc.* **26**, 745–750.

## **supplementary materials**

Acta Cryst. (2008). E64, o1611 [doi:10.1107/S1600536808023180]

### 3-Benzylsulfanyl-5-(4-phenyl-1H-1,2,3-triazol-1-ylmethyl)-4H-1,2,4-triazol-4-amine

Q.-Z. Chu, H.-R. Zhou and X.-R. Zhang

#### Comment

Triazole derivatives exhibit higher antifungal activity against filamentous fungi and yeasts (Barchiesi *et al.*, 2000), and their antibacterial properties have been reported as well (Colanceska-Ragenovic *et al.*, 2001). The biological activities of various 1,2,3-triazole and 1,2,4-triazole derivatives have been extensively studied (Rodriguez-Fernandez *et al.*, 2005; Kaplancikli *et al.*, 2008; Khanum *et al.*, 2005). As an extension of the work on structure characterization of triazole derivatives, the title compound (**I**) was synthesized and its structure was characterized by X-ray crystallographic analysis.

In (**I**) (Fig. 1), all the bond lengths and angles are within normal ranges (Allen *et al.*, 1987). A (N1–N3/C7/C8) triazole ring makes dihedral angles of 11.6 (3) and 25.6 (3)° with B (C1–C6) and C (C13–C18) phenyl rings, respectively, and of 71.4 (4)° with D (N4–N6/C10/C11) triazole ring. In addition, the dihedral angles between B and C, C and D, and B and D are 15.5 (3), 67.7 (3) and 73.7 (3)°, respectively. There exists one intramolecular hydrogen bond, N7—H7B···S1 (Table 1), forming a five-membered ring.

#### Experimental

3-{{[4-(4-phenyl)-1H-1,2,3-triazol-1-yl]methyl}-4-amino-5-thiol-1,2,4-triazole (Zhang *et al.*, 2008) (1.00 g, 3.66 mmol) was added to a mixture of 50% NaOH (*w/w*) 5 ml, water (30 ml) and benzyl bromide (0.87 ml, 7.32 mmol). The mixture was stirred for 5 h. The resulting precipitate was collected by filtration, washed by water and recrystallized from acetone to give compound **I** (1.09 g, 82.0%). Colourless single crystals suitable for an X-ray diffraction study were obtained by slow evaporation of an ethyl acetate solution.

#### Refinement

All H atoms were located in difference Fourier maps and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å and N—H distances of 0.86 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  H atoms.

#### Figures




Fig. 1. The structure of the compound (**I**) showing 50% probability displacement ellipsoids and the atom numbering scheme.




Fig. 2. A packing diagram of (**I**), viewed down the *c* axis.

# supplementary materials

---

## 3-Benzylsulfanyl-5-(4-phenyl-1*H*-1,2,3-triazol-1-ylmethyl)-4*H*-1,2,4-triazol-4-amine

### Crystal data

C <sub>18</sub> H <sub>17</sub> N <sub>7</sub> S	$F_{000} = 760$
$M_r = 363.45$	$D_x = 1.416 \text{ Mg m}^{-3}$
Orthorhombic, $Pna2_1$	Mo $K\alpha$ radiation
Hall symbol: P 2c -2n	$\lambda = 0.71073 \text{ \AA}$
$a = 8.0487 (15) \text{ \AA}$	Cell parameters from 1817 reflections
$b = 5.4689 (10) \text{ \AA}$	$\theta = 3.2\text{--}25.0^\circ$
$c = 38.721 (7) \text{ \AA}$	$\mu = 0.21 \text{ mm}^{-1}$
$V = 1704.4 (5) \text{ \AA}^3$	$T = 186.5 (2) \text{ K}$
$Z = 4$	Plate, colourless
	$0.35 \times 0.25 \times 0.04 \text{ mm}$

### Data collection

Siemens SMART 1000 CCD area-detector diffractometer	2854 independent reflections
Radiation source: fine-focus sealed tube	2592 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.052$
Detector resolution: 8.33 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 25.0^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 2.1^\circ$
$\omega$ scans	$h = -9 \rightarrow 9$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$k = -6 \rightarrow 5$
$T_{\text{min}} = 0.931$ , $T_{\text{max}} = 0.992$	$l = -40 \rightarrow 46$
7815 measured reflections	

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.078$	$w = 1/[\sigma^2(F_o^2) + (0.0579P)^2 + 3.955P]$
$wR(F^2) = 0.183$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.20$	$(\Delta/\sigma)_{\text{max}} = 0.002$
2854 reflections	$\Delta\rho_{\text{max}} = 0.48 \text{ e \AA}^{-3}$
235 parameters	$\Delta\rho_{\text{min}} = -0.39 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), with 2854 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.05 (19)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.0735 (2)	0.5751 (3)	0.34992 (5)	0.0348 (4)
N5	-0.0030 (7)	0.2276 (10)	0.39829 (15)	0.0349 (13)
C11	0.0723 (7)	0.4329 (11)	0.38974 (17)	0.0284 (14)
N6	0.1621 (6)	0.5157 (9)	0.41739 (14)	0.0299 (12)
C1	-0.0955 (7)	0.3110 (10)	0.60677 (16)	0.0266 (14)
H1A	-0.0314	0.1797	0.5993	0.032*
C7	-0.0485 (7)	0.5171 (11)	0.55000 (16)	0.0245 (14)
N4	0.0366 (7)	0.1806 (11)	0.43251 (16)	0.0398 (15)
C13	-0.0838 (8)	0.4412 (12)	0.29074 (17)	0.0344 (16)
N1	-0.0513 (8)	0.7223 (11)	0.53032 (15)	0.0428 (15)
C3	-0.2600 (7)	0.4969 (12)	0.65095 (18)	0.0305 (14)
H3A	-0.3061	0.4923	0.6730	0.037*
N2	0.0363 (8)	0.6895 (11)	0.50221 (17)	0.0464 (16)
C5	-0.2313 (8)	0.6963 (10)	0.59606 (18)	0.0297 (15)
H5B	-0.2576	0.8261	0.5815	0.036*
C14	-0.0064 (8)	0.2866 (11)	0.26757 (19)	0.0356 (17)
H14A	0.0496	0.1498	0.2758	0.043*
N3	0.0979 (6)	0.4597 (10)	0.50423 (14)	0.0322 (13)
C16	-0.0919 (9)	0.5334 (14)	0.2199 (2)	0.0419 (18)
H16A	-0.0967	0.5634	0.1963	0.050*
N7	0.2650 (7)	0.7185 (10)	0.41686 (15)	0.0423 (15)
H7A	0.3227	0.7555	0.4348	0.051*
H7B	0.2712	0.8075	0.3986	0.051*
C2	-0.1574 (9)	0.3117 (11)	0.63972 (17)	0.0354 (16)
H2B	-0.1299	0.1853	0.6547	0.042*
C4	-0.2935 (8)	0.6914 (11)	0.62878 (17)	0.0302 (15)
H4B	-0.3594	0.8201	0.6365	0.036*
C6	-0.1276 (7)	0.5067 (11)	0.58396 (16)	0.0258 (14)
C17	-0.1648 (8)	0.6881 (11)	0.24299 (18)	0.0334 (16)
H17A	-0.2206	0.8245	0.2346	0.040*
C10	0.1372 (7)	0.3584 (12)	0.44322 (16)	0.0286 (14)
C18	-0.1598 (8)	0.6523 (11)	0.27809 (18)	0.0321 (15)
H18A	-0.2062	0.7662	0.2931	0.038*

## supplementary materials

---

C8	0.0472 (7)	0.3493 (11)	0.53312 (16)	0.0271 (14)
H8A	0.0720	0.1910	0.5403	0.033*
C15	-0.0103 (8)	0.3297 (13)	0.2329 (2)	0.0386 (17)
H15A	0.0421	0.2220	0.2178	0.046*
C9	0.2124 (8)	0.3702 (15)	0.47810 (17)	0.0405 (17)
H9A	0.2502	0.2082	0.4846	0.049*
H9B	0.3088	0.4766	0.4774	0.049*
C12	-0.0816 (10)	0.3877 (15)	0.3288 (2)	0.053 (2)
H12A	-0.0565	0.2162	0.3325	0.063*
H12B	-0.1901	0.4213	0.3386	0.063*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0327 (8)	0.0356 (9)	0.0361 (9)	-0.0077 (7)	-0.0003 (8)	0.0057 (9)
N5	0.029 (3)	0.035 (3)	0.041 (3)	-0.009 (3)	0.000 (2)	0.007 (3)
C11	0.019 (3)	0.024 (3)	0.043 (4)	0.001 (3)	0.005 (3)	-0.001 (3)
N6	0.027 (3)	0.026 (3)	0.037 (3)	0.001 (2)	0.002 (2)	0.001 (3)
C1	0.018 (3)	0.019 (3)	0.042 (4)	0.004 (3)	-0.004 (3)	0.003 (3)
C7	0.019 (3)	0.017 (3)	0.037 (4)	0.003 (2)	-0.005 (2)	-0.002 (3)
N4	0.033 (3)	0.043 (4)	0.043 (4)	0.004 (3)	0.009 (3)	0.014 (3)
C13	0.032 (4)	0.029 (3)	0.042 (4)	-0.006 (3)	-0.006 (3)	0.008 (3)
N1	0.046 (3)	0.039 (3)	0.043 (4)	0.014 (3)	0.011 (3)	0.011 (3)
C3	0.020 (3)	0.032 (4)	0.040 (4)	-0.006 (3)	0.000 (3)	-0.001 (3)
N2	0.047 (4)	0.037 (4)	0.055 (4)	0.018 (3)	0.011 (3)	0.016 (3)
C5	0.032 (3)	0.010 (3)	0.048 (4)	0.004 (3)	-0.007 (3)	0.002 (2)
C14	0.021 (3)	0.015 (3)	0.071 (5)	0.007 (3)	-0.011 (3)	0.006 (3)
N3	0.016 (2)	0.041 (3)	0.039 (3)	0.011 (2)	0.000 (2)	0.002 (3)
C16	0.036 (4)	0.051 (5)	0.039 (4)	-0.019 (4)	-0.004 (3)	-0.001 (4)
N7	0.042 (3)	0.033 (3)	0.051 (4)	-0.011 (3)	-0.008 (3)	0.006 (3)
C2	0.042 (4)	0.025 (3)	0.040 (4)	-0.005 (3)	-0.002 (3)	0.005 (3)
C4	0.024 (3)	0.023 (3)	0.044 (4)	0.011 (3)	0.001 (3)	-0.004 (3)
C6	0.025 (3)	0.020 (3)	0.032 (4)	-0.008 (3)	-0.007 (3)	0.001 (2)
C17	0.026 (3)	0.018 (3)	0.056 (5)	-0.009 (3)	0.005 (3)	0.001 (3)
C10	0.024 (3)	0.030 (3)	0.031 (4)	0.005 (3)	0.003 (3)	0.004 (3)
C18	0.023 (3)	0.023 (3)	0.050 (4)	0.004 (3)	0.006 (3)	0.000 (3)
C8	0.029 (3)	0.015 (3)	0.037 (4)	0.011 (3)	-0.005 (3)	-0.001 (3)
C15	0.030 (4)	0.034 (4)	0.052 (5)	-0.001 (3)	-0.001 (3)	-0.015 (3)
C9	0.026 (3)	0.060 (5)	0.035 (4)	0.007 (3)	0.007 (3)	-0.001 (4)
C12	0.045 (4)	0.060 (5)	0.053 (5)	-0.034 (4)	-0.008 (4)	0.012 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

S1—C11	1.727 (6)	C5—C6	1.411 (9)
S1—C12	1.810 (7)	C5—H5B	0.9300
N5—C11	1.318 (8)	C14—C15	1.365 (10)
N5—N4	1.387 (8)	C14—H14A	0.9300
C11—N6	1.369 (8)	N3—C8	1.335 (8)
N6—C10	1.334 (8)	N3—C9	1.454 (8)

N6—N7	1.385 (7)	C16—C17	1.363 (10)
C1—C2	1.370 (9)	C16—C15	1.386 (10)
C1—C6	1.412 (8)	C16—H16A	0.9300
C1—H1A	0.9300	N7—H7A	0.8600
C7—N1	1.357 (8)	N7—H7B	0.8600
C7—C8	1.365 (8)	C2—H2B	0.9300
C7—C6	1.462 (8)	C4—H4B	0.9300
N4—C10	1.332 (9)	C17—C18	1.374 (9)
C13—C14	1.381 (10)	C17—H17A	0.9300
C13—C18	1.395 (9)	C10—C9	1.481 (9)
C13—C12	1.503 (10)	C18—H18A	0.9300
N1—N2	1.309 (8)	C8—H8A	0.9300
C3—C2	1.377 (9)	C15—H15A	0.9300
C3—C4	1.393 (9)	C9—H9A	0.9700
C3—H3A	0.9300	C9—H9B	0.9700
N2—N3	1.353 (8)	C12—H12A	0.9700
C5—C4	1.362 (9)	C12—H12B	0.9700
C11—S1—C12	98.3 (3)	H7A—N7—H7B	120.0
C11—N5—N4	107.0 (5)	C1—C2—C3	121.0 (6)
N5—C11—N6	109.2 (5)	C1—C2—H2B	119.5
N5—C11—S1	127.6 (5)	C3—C2—H2B	119.5
N6—C11—S1	123.1 (4)	C5—C4—C3	121.2 (6)
C10—N6—C11	107.1 (5)	C5—C4—H4B	119.4
C10—N6—N7	128.1 (6)	C3—C4—H4B	119.4
C11—N6—N7	124.7 (5)	C5—C6—C1	117.2 (6)
C2—C1—C6	120.9 (6)	C5—C6—C7	121.9 (6)
C2—C1—H1A	119.5	C1—C6—C7	120.8 (6)
C6—C1—H1A	119.5	C16—C17—C18	123.2 (7)
N1—C7—C8	107.3 (6)	C16—C17—H17A	118.4
N1—C7—C6	122.0 (5)	C18—C17—H17A	118.4
C8—C7—C6	130.6 (6)	N4—C10—N6	109.2 (6)
C10—N4—N5	107.6 (5)	N4—C10—C9	124.3 (6)
C14—C13—C18	118.4 (6)	N6—C10—C9	126.5 (6)
C14—C13—C12	120.8 (6)	C17—C18—C13	118.6 (6)
C18—C13—C12	120.7 (7)	C17—C18—H18A	120.7
N2—N1—C7	110.1 (5)	C13—C18—H18A	120.7
C2—C3—C4	118.9 (6)	N3—C8—C7	105.6 (5)
C2—C3—H3A	120.6	N3—C8—H8A	127.2
C4—C3—H3A	120.6	C7—C8—H8A	127.2
N1—N2—N3	106.0 (5)	C14—C15—C16	120.3 (7)
C4—C5—C6	120.7 (6)	C14—C15—H15A	119.8
C4—C5—H5B	119.6	C16—C15—H15A	119.8
C6—C5—H5B	119.6	N3—C9—C10	113.0 (5)
C15—C14—C13	121.6 (6)	N3—C9—H9A	109.0
C15—C14—H14A	119.2	C10—C9—H9A	109.0
C13—C14—H14A	119.2	N3—C9—H9B	109.0
C8—N3—N2	110.9 (5)	C10—C9—H9B	109.0
C8—N3—C9	128.7 (5)	H9A—C9—H9B	107.8
N2—N3—C9	120.3 (6)	C13—C12—S1	109.9 (5)

## supplementary materials


---

C17—C16—C15	117.8 (7)	C13—C12—H12A	109.7
C17—C16—H16A	121.1	S1—C12—H12A	109.7
C15—C16—H16A	121.1	C13—C12—H12B	109.7
N6—N7—H7A	120.0	S1—C12—H12B	109.7
N6—N7—H7B	120.0	H12A—C12—H12B	108.2
N4—N5—C11—N6	1.3 (7)	N1—C7—C6—C1	-167.2 (6)
N4—N5—C11—S1	177.6 (5)	C8—C7—C6—C1	7.4 (10)
C12—S1—C11—N5	9.0 (7)	C15—C16—C17—C18	0.5 (10)
C12—S1—C11—N6	-175.1 (6)	N5—N4—C10—N6	0.3 (7)
N5—C11—N6—C10	-1.1 (7)	N5—N4—C10—C9	-177.8 (6)
S1—C11—N6—C10	-177.7 (5)	C11—N6—C10—N4	0.5 (7)
N5—C11—N6—N7	175.7 (6)	N7—N6—C10—N4	-176.2 (6)
S1—C11—N6—N7	-0.8 (8)	C11—N6—C10—C9	178.5 (6)
C11—N5—N4—C10	-1.0 (7)	N7—N6—C10—C9	1.8 (10)
C8—C7—N1—N2	0.4 (8)	C16—C17—C18—C13	-3.3 (10)
C6—C7—N1—N2	176.2 (6)	C14—C13—C18—C17	4.6 (9)
C7—N1—N2—N3	-0.7 (8)	C12—C13—C18—C17	-177.3 (6)
C18—C13—C14—C15	-3.3 (10)	N2—N3—C8—C7	-0.6 (7)
C12—C13—C14—C15	178.6 (6)	C9—N3—C8—C7	175.1 (6)
N1—N2—N3—C8	0.8 (7)	N1—C7—C8—N3	0.1 (7)
N1—N2—N3—C9	-175.3 (6)	C6—C7—C8—N3	-175.1 (6)
C6—C1—C2—C3	3.4 (9)	C13—C14—C15—C16	0.5 (10)
C4—C3—C2—C1	-3.2 (9)	C17—C16—C15—C14	0.9 (10)
C6—C5—C4—C3	-1.6 (10)	C8—N3—C9—C10	121.5 (7)
C2—C3—C4—C5	2.3 (10)	N2—N3—C9—C10	-63.1 (8)
C4—C5—C6—C1	1.7 (9)	N4—C10—C9—N3	-79.2 (8)
C4—C5—C6—C7	-175.0 (6)	N6—C10—C9—N3	103.1 (8)
C2—C1—C6—C5	-2.6 (8)	C14—C13—C12—S1	101.0 (7)
C2—C1—C6—C7	174.1 (6)	C18—C13—C12—S1	-77.1 (8)
N1—C7—C6—C5	9.3 (9)	C11—S1—C12—C13	-169.3 (6)
C8—C7—C6—C5	-176.0 (6)		

*Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )*

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N7—H7B…S1	0.86	2.78	3.116 (6)	105

Fig. 1



## **supplementary materials**

---

**Fig. 2**

