

3,3'-Bis(3,4,5-trimethoxybenzoyl)-1,1'-(*o*-phenylene)dithiourea ethanol solvate

Hai-Tang Du^{a*} and Hai-Jun Du^b

^aInstitute of Natural Products, Research Center for Eco-Environmental Sciences, Guiyang College, Guiyang 550005, People's Republic of China, and ^bSchool of Chemistry and Environmental Science, Guizhou University for Nationalities, Guiyang 550025, People's Republic of China

Correspondence e-mail: haitangdu@gz139.com.cn


Received 24 July 2008; accepted 25 July 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.050; wR factor = 0.116; data-to-parameter ratio = 15.8.

In the molecule of the title compound, $\text{C}_{28}\text{H}_{30}\text{N}_4\text{O}_8\text{S}_2 \cdot \text{C}_2\text{H}_6\text{O}$, the benzene ring is oriented at dihedral angles of 38.50 (6) and 5.68 (5)° with respect to the trimethoxyphenyl rings, while the two trimethoxyphenyl rings are oriented at a dihedral angle of 44.18 (5)°. Intramolecular $\text{N}-\text{H} \cdots \text{O}$ and $\text{N}-\text{H} \cdots \text{S}$ hydrogen bonds result in the formation of non-planar six-, seven- and eight-membered rings. The twisting modes of the two side arms are different [$\text{C}-\text{N}-\text{C}-\text{O}$ and $\text{C}-\text{N}-\text{C}-\text{N}$ torsion angles = 0.1 (3) and 11.8 (3)°, respectively, in one arm, and 4.6 (3) and -11.5 (3)° in the other]. In the crystal structure, intermolecular $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link the molecules.

Related literature

For a related structure, see: Thiam *et al.* (2008). For ring conformation puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{30}\text{N}_4\text{O}_8\text{S}_2 \cdot \text{C}_2\text{H}_6\text{O}$

$M_r = 660.75$

Triclinic, $P\bar{1}$

$a = 7.7619$ (15) Å

$b = 14.473$ (3) Å

$c = 15.810$ (3) Å

$\alpha = 67.113$ (10)°

$\beta = 73.069$ (9)°

$\gamma = 78.210$ (12)°

$V = 1556.9$ (5) Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 0.23$ mm⁻¹

$T = 113$ (2) K

$0.14 \times 0.12 \times 0.10$ mm

Data collection

Rigaku Saturn CCD area-detector diffractometer

Absorption correction: multi-scan

(*CrystalClear*; Rigaku/MSC, 2005)

$T_{\min} = 0.968$, $T_{\max} = 0.977$

18692 measured reflections

6824 independent reflections

5694 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.115$

$S = 1.07$

6824 reflections

433 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.26$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N4}-\text{H4A} \cdots \text{O1}^i$	0.90 (2)	2.51 (2)	3.403 (2)	173 (2)
$\text{N2}-\text{H2} \cdots \text{S1}$	0.88 (2)	2.69 (2)	3.3527 (19)	133.0 (18)
$\text{N2}-\text{H2} \cdots \text{O2}$	0.88 (2)	1.97 (2)	2.677 (2)	136 (2)
$\text{N1}-\text{H1} \cdots \text{O1}$	0.89 (2)	1.90 (2)	2.621 (2)	137 (2)
$\text{N3}-\text{H3A} \cdots \text{O9}$	0.84 (2)	2.23 (2)	2.949 (2)	145 (2)
$\text{O9}-\text{H9} \cdots \text{O2}$	0.80 (3)	2.13 (3)	2.905 (2)	163 (3)

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku/MSC, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Guiyang College for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2504).

References

- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Rigaku/MS. (2005). *CrystalClear* and *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Thiam, E. I., Diop, M., Gaye, M., Sall, A. S. & Barry, A. H. (2008). *Acta Cryst.* **E64**, o776.

supplementary materials

Acta Cryst. (2008). E64, o1632-o1633 [doi:10.1107/S1600536808023556]

3,3'-Bis(3,4,5-trimethoxybenzoyl)-1,1'-(*o*-phenylene)dithiourea ethanol solvate

H.-T. Du and H.-J. Du

Comment

In the molecule of the title compound (Fig. 1) the bond lengths and angles are within normal ranges. Rings A (C1-C6), B (C9-C14) and C (C17-C22) are, of course, planar, and the dihedral angles between them are A/B = 38.50 (6)°, A/C = 5.68 (5)° and B/C = 44.18 (5)°.

The intramolecular N-H...O and N-H...S hydrogen bonds (Table 1) result in the formation of non-planar six-, seven- and eight-membered rings: D (O1/N1/N3/C7/C8/H1), E (O2/N2/N4/C15/C16/H2), F (S1/N1/N2/C1/C2/C7/H2) and G (S1/O2/O9/N3/C7/H2/H3A/H9). Rings D and E adopt flattened-boat [$\phi = 171.38$ (2)°, $\theta = 109.10$ (3)° (for ring D) and $\phi = -20.28$ (3)°, $\theta = 96.87$ (3)° (for ring E)] conformations, while rings F and G adopt highly twisted conformations having total puckering amplitudes, Q_T , of 0.160 (3), 0.109 (3), 2.486 (4) and 2.064 (4) Å, respectively (Cremer & Pople, 1975). The two side arms are not twisted in the same way, as evidenced by the torsion angles: C7-N3-C8-O1 [0.1 (3)°], C8-N3-C7-N1 [11.8 (3)°] and C15-N4-C16-O2 [4.6 (3)°], C16-N4-C15-N2 [-11.5 (3)°], as in 1,2-bis(*N*'-benzoylthioureido)- benzene (Thiam *et al.*, 2008).

In the crystal structure, intermolecular N-H...O hydrogen bonds link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure.

Experimental

For the preparation of the title compound, ammonium thiocyanate (30 mmol), 3,4,5-trimethoxybenzoyl chloride (20 mmol), PEG-400 (0.2 mmol) and acetone (50 mL) were placed in a dried round-bottomed flask containing a magnetic stirrer bar and stirred at room temperature for 1 h, then benzene-1,2-diamine (9.5 mmol) was added, and the mixture was stirred for 2 h. The mixture was poured into water (20 ml). The resulting solid was filtered, washed with water, and then dried. Crystals suitable for X-ray analysis were obtained by the recrystallization of the solid residue from a mixture of *N,N*-dimethyl-formamide/ethanol (1:1) by slow evaporation at room temperature.

Refinement

H1, H2, H3A, H4A (for NH) and H9 (for OH) atoms were located in difference syntheses and refined isotropically [N-H = 0.84 (2)-0.90 (2) Å and $U_{\text{iso}}(\text{H}) = 0.026$ (6)-0.036 (7) Å²; O-H = 0.80 (3) Å and $U_{\text{iso}}(\text{H}) = 0.043$ Å²]. The remaining H atoms were positioned geometrically, with C-H = 0.95, 0.98 and 0.99 Å for aromatic, methyl and methylene H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

3,3'-Bis(3,4,5-trimethoxybenzoyl)-1,1'-(o-phenylene)dithiourea ethanol solvate

Crystal data

$C_{28}H_{30}N_4O_8S_2 \cdot C_2H_6O_1$

$M_r = 660.75$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.7619$ (15) Å

$b = 14.473$ (3) Å

$c = 15.810$ (3) Å

$\alpha = 67.113$ (10)°

$\beta = 73.069$ (9)°

$\gamma = 78.210$ (12)°

$V = 1556.9$ (5) Å³

$Z = 2$

$F_{000} = 696$

$D_x = 1.409$ Mg m⁻³

Melting point: 475 K

Mo $K\alpha$ radiation

$\lambda = 0.71070$ Å

Cell parameters from 4542 reflections

$\theta = 2.4\text{--}27.2^\circ$

$\mu = 0.23$ mm⁻¹

$T = 113$ (2) K

Block, colorless

$0.14 \times 0.12 \times 0.10$ mm

Data collection

Rigaku Saturn CCD area-detector diffractometer

Radiation source: rotating anode

Monochromator: confocal

Detector resolution: 14.63 pixels mm⁻¹

$T = 113$ (2) K

ω scans

Absorption correction: multi-scan (CrystalClear; Rigaku/MS, 2005)

$T_{\min} = 0.968$, $T_{\max} = 0.977$

18692 measured reflections

6824 independent reflections

5694 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\text{max}} = 27.2^\circ$

$\theta_{\text{min}} = 2.5^\circ$

$h = -9 \rightarrow 9$

$k = -18 \rightarrow 18$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.050$	$w = 1/[\sigma^2(F_o^2) + (0.0441P)^2 + 0.4672P]$
$wR(F^2) = 0.115$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} < 0.001$
6824 reflections	$\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$
433 parameters	$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001 \times Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0140 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.47072 (7)	0.74462 (4)	0.51329 (3)	0.02065 (14)
S2	0.93034 (8)	0.44117 (4)	0.34131 (4)	0.02578 (15)
O1	1.02467 (19)	0.81993 (11)	0.48592 (9)	0.0250 (3)
O2	0.6648 (2)	0.51201 (10)	0.60921 (10)	0.0274 (4)
O3	1.2473 (2)	0.92569 (11)	0.70183 (10)	0.0286 (4)
O4	1.0727 (2)	0.82041 (11)	0.87598 (10)	0.0278 (4)
O5	0.7886 (2)	0.72356 (11)	0.90656 (10)	0.0287 (4)
O6	0.7859 (2)	0.27652 (10)	0.93150 (10)	0.0243 (3)
O7	0.7307 (2)	0.09337 (10)	0.94217 (9)	0.0244 (3)
O8	0.6871 (2)	0.06352 (10)	0.79539 (9)	0.0229 (3)
O9	0.5525 (2)	0.62076 (12)	0.74010 (11)	0.0285 (4)
H9	0.568 (4)	0.582 (2)	0.7125 (19)	0.043*
N1	0.8159 (2)	0.76073 (12)	0.41701 (11)	0.0177 (4)
H1	0.923 (3)	0.7772 (17)	0.4133 (16)	0.030 (7)*
N2	0.7346 (2)	0.57673 (12)	0.42050 (12)	0.0200 (4)

supplementary materials

H2	0.685 (3)	0.5885 (17)	0.4737 (17)	0.028 (6)*
N3	0.7451 (2)	0.76939 (12)	0.56722 (12)	0.0184 (4)
H3A	0.673 (3)	0.7516 (17)	0.6201 (17)	0.026 (6)*
N4	0.7865 (2)	0.41196 (12)	0.52015 (11)	0.0198 (4)
H4A	0.830 (3)	0.3486 (19)	0.5237 (17)	0.036 (7)*
C1	0.7839 (3)	0.75352 (14)	0.33508 (13)	0.0175 (4)
C2	0.7479 (3)	0.66254 (14)	0.33554 (13)	0.0176 (4)
C3	0.7222 (3)	0.65850 (15)	0.25351 (14)	0.0207 (4)
H3	0.6911	0.5981	0.2539	0.025*
C4	0.7419 (3)	0.74278 (16)	0.17098 (14)	0.0238 (5)
H4	0.7272	0.7392	0.1148	0.029*
C5	0.7829 (3)	0.83206 (16)	0.17021 (14)	0.0249 (5)
H5	0.7987	0.8890	0.1134	0.030*
C6	0.8006 (3)	0.83775 (15)	0.25285 (14)	0.0213 (4)
H6	0.8242	0.8995	0.2532	0.026*
C7	0.6875 (3)	0.75825 (13)	0.49576 (13)	0.0167 (4)
C8	0.9074 (3)	0.79901 (14)	0.56059 (14)	0.0193 (4)
C9	0.9324 (3)	0.80961 (14)	0.64638 (14)	0.0191 (4)
C10	1.0706 (3)	0.86638 (14)	0.63139 (14)	0.0199 (4)
H10	1.1358	0.9001	0.5690	0.024*
C11	1.1123 (3)	0.87340 (15)	0.70806 (14)	0.0207 (4)
C12	1.0173 (3)	0.82339 (15)	0.80012 (14)	0.0217 (4)
C13	0.8750 (3)	0.76923 (15)	0.81379 (13)	0.0215 (4)
C14	0.8326 (3)	0.76151 (15)	0.73745 (14)	0.0219 (4)
H14	0.7370	0.7240	0.7471	0.026*
C15	0.8105 (3)	0.48263 (14)	0.42749 (14)	0.0189 (4)
C16	0.7225 (3)	0.42760 (15)	0.60517 (14)	0.0197 (4)
C17	0.7314 (3)	0.33652 (14)	0.69148 (13)	0.0180 (4)
C18	0.7536 (3)	0.35253 (15)	0.76900 (14)	0.0200 (4)
H18	0.7662	0.4182	0.7647	0.024*
C19	0.7570 (3)	0.27097 (15)	0.85280 (13)	0.0193 (4)
C20	0.7301 (3)	0.17510 (14)	0.86015 (13)	0.0187 (4)
C21	0.7114 (3)	0.16009 (14)	0.78097 (14)	0.0187 (4)
C22	0.7134 (3)	0.24057 (14)	0.69593 (13)	0.0172 (4)
H22	0.7026	0.2304	0.6419	0.021*
C23	1.3530 (3)	0.97237 (17)	0.60921 (16)	0.0301 (5)
H23A	1.4118	0.9209	0.5812	0.045*
H23B	1.4455	1.0069	0.6127	0.045*
H23C	1.2744	1.0214	0.5701	0.045*
C24	0.9843 (3)	0.89960 (16)	0.91201 (15)	0.0274 (5)
H24A	1.0110	0.9651	0.8628	0.041*
H24B	1.0285	0.8918	0.9667	0.041*
H24C	0.8532	0.8958	0.9311	0.041*
C25	0.6375 (3)	0.67215 (18)	0.92291 (16)	0.0349 (6)
H25A	0.5474	0.7192	0.8903	0.052*
H25B	0.5834	0.6455	0.9909	0.052*
H25C	0.6777	0.6164	0.8988	0.052*
C26	0.8306 (3)	0.37088 (16)	0.92470 (15)	0.0277 (5)
H26A	0.9337	0.3926	0.8709	0.042*

H26B	0.8624	0.3629	0.9829	0.042*
H26C	0.7262	0.4217	0.9156	0.042*
C27	0.5853 (3)	0.09950 (17)	1.02149 (16)	0.0348 (6)
H27A	0.6219	0.1337	1.0552	0.052*
H27B	0.5573	0.0314	1.0643	0.052*
H27C	0.4777	0.1376	0.9989	0.052*
C28	0.6740 (3)	0.04397 (15)	0.71577 (14)	0.0249 (5)
H28A	0.5684	0.0849	0.6929	0.037*
H28B	0.6608	-0.0276	0.7343	0.037*
H28C	0.7839	0.0613	0.6654	0.037*
C29	0.3609 (3)	0.63770 (18)	0.77554 (16)	0.0312 (5)
H29A	0.2987	0.6509	0.7249	0.037*
H29B	0.3343	0.6983	0.7937	0.037*
C30	0.2880 (4)	0.5489 (2)	0.85986 (17)	0.0428 (7)
H30A	0.3103	0.4893	0.8416	0.064*
H30B	0.1575	0.5639	0.8824	0.064*
H30C	0.3489	0.5358	0.9103	0.064*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0181 (3)	0.0241 (3)	0.0211 (3)	-0.00133 (19)	-0.0054 (2)	-0.0093 (2)
S2	0.0320 (3)	0.0201 (3)	0.0198 (3)	0.0011 (2)	-0.0011 (2)	-0.0068 (2)
O1	0.0240 (8)	0.0343 (8)	0.0179 (7)	-0.0081 (6)	-0.0015 (6)	-0.0103 (6)
O2	0.0436 (10)	0.0171 (7)	0.0200 (8)	0.0031 (6)	-0.0090 (7)	-0.0070 (6)
O3	0.0308 (9)	0.0363 (9)	0.0256 (8)	-0.0101 (7)	-0.0086 (7)	-0.0137 (7)
O4	0.0350 (9)	0.0317 (8)	0.0237 (8)	0.0064 (6)	-0.0154 (7)	-0.0160 (7)
O5	0.0367 (9)	0.0332 (8)	0.0156 (7)	-0.0084 (7)	-0.0045 (6)	-0.0069 (6)
O6	0.0353 (9)	0.0211 (7)	0.0203 (7)	-0.0044 (6)	-0.0106 (6)	-0.0076 (6)
O7	0.0329 (9)	0.0187 (7)	0.0160 (7)	0.0021 (6)	-0.0059 (6)	-0.0024 (6)
O8	0.0356 (9)	0.0149 (7)	0.0202 (7)	-0.0050 (6)	-0.0100 (6)	-0.0046 (6)
O9	0.0278 (9)	0.0318 (9)	0.0286 (9)	-0.0028 (7)	-0.0046 (7)	-0.0149 (7)
N1	0.0187 (9)	0.0202 (8)	0.0159 (8)	-0.0019 (6)	-0.0051 (7)	-0.0073 (7)
N2	0.0280 (10)	0.0165 (8)	0.0142 (8)	-0.0005 (7)	-0.0056 (7)	-0.0045 (7)
N3	0.0199 (9)	0.0227 (9)	0.0134 (8)	-0.0030 (7)	-0.0032 (7)	-0.0074 (7)
N4	0.0268 (10)	0.0137 (8)	0.0165 (8)	-0.0013 (7)	-0.0036 (7)	-0.0041 (7)
C1	0.0152 (10)	0.0219 (10)	0.0155 (9)	0.0005 (7)	-0.0032 (7)	-0.0084 (8)
C2	0.0171 (10)	0.0183 (9)	0.0158 (9)	0.0013 (7)	-0.0035 (8)	-0.0061 (8)
C3	0.0218 (11)	0.0214 (10)	0.0202 (10)	0.0017 (8)	-0.0066 (8)	-0.0094 (8)
C4	0.0252 (11)	0.0293 (11)	0.0184 (10)	0.0045 (8)	-0.0084 (8)	-0.0112 (9)
C5	0.0285 (12)	0.0248 (11)	0.0155 (10)	0.0012 (8)	-0.0048 (9)	-0.0031 (8)
C6	0.0229 (11)	0.0174 (10)	0.0197 (10)	-0.0010 (8)	-0.0031 (8)	-0.0043 (8)
C7	0.0219 (10)	0.0131 (9)	0.0151 (9)	0.0000 (7)	-0.0069 (8)	-0.0039 (7)
C8	0.0206 (10)	0.0176 (9)	0.0205 (10)	-0.0003 (7)	-0.0067 (8)	-0.0071 (8)
C9	0.0209 (10)	0.0194 (10)	0.0195 (10)	0.0019 (8)	-0.0069 (8)	-0.0098 (8)
C10	0.0201 (10)	0.0198 (10)	0.0190 (10)	0.0009 (8)	-0.0057 (8)	-0.0066 (8)
C11	0.0210 (11)	0.0215 (10)	0.0234 (10)	0.0007 (8)	-0.0072 (8)	-0.0117 (8)
C12	0.0263 (11)	0.0229 (10)	0.0208 (10)	0.0051 (8)	-0.0121 (9)	-0.0119 (8)

supplementary materials

C13	0.0254 (11)	0.0217 (10)	0.0151 (10)	0.0019 (8)	-0.0055 (8)	-0.0056 (8)
C14	0.0241 (11)	0.0213 (10)	0.0212 (10)	-0.0013 (8)	-0.0063 (8)	-0.0083 (8)
C15	0.0197 (10)	0.0178 (9)	0.0201 (10)	-0.0032 (7)	-0.0058 (8)	-0.0061 (8)
C16	0.0222 (10)	0.0197 (10)	0.0180 (10)	-0.0037 (8)	-0.0055 (8)	-0.0063 (8)
C17	0.0176 (10)	0.0189 (10)	0.0158 (9)	0.0005 (7)	-0.0034 (8)	-0.0057 (8)
C18	0.0217 (10)	0.0182 (10)	0.0211 (10)	-0.0023 (7)	-0.0051 (8)	-0.0077 (8)
C19	0.0195 (10)	0.0240 (10)	0.0150 (9)	-0.0009 (8)	-0.0047 (8)	-0.0077 (8)
C20	0.0199 (10)	0.0173 (9)	0.0159 (9)	-0.0013 (7)	-0.0047 (8)	-0.0027 (8)
C21	0.0176 (10)	0.0161 (9)	0.0233 (10)	-0.0024 (7)	-0.0054 (8)	-0.0071 (8)
C22	0.0175 (10)	0.0187 (9)	0.0149 (9)	-0.0020 (7)	-0.0037 (7)	-0.0051 (8)
C23	0.0306 (13)	0.0312 (12)	0.0327 (12)	-0.0105 (9)	-0.0052 (10)	-0.0137 (10)
C24	0.0357 (13)	0.0284 (11)	0.0226 (11)	-0.0013 (9)	-0.0092 (9)	-0.0129 (9)
C25	0.0463 (15)	0.0343 (13)	0.0222 (11)	-0.0170 (11)	0.0012 (10)	-0.0083 (10)
C26	0.0365 (13)	0.0264 (11)	0.0263 (11)	-0.0053 (9)	-0.0096 (10)	-0.0129 (9)
C27	0.0348 (13)	0.0300 (12)	0.0250 (12)	-0.0045 (10)	0.0006 (10)	0.0008 (10)
C28	0.0355 (13)	0.0191 (10)	0.0235 (11)	-0.0022 (8)	-0.0110 (9)	-0.0086 (9)
C29	0.0277 (12)	0.0373 (13)	0.0277 (12)	-0.0028 (9)	-0.0076 (10)	-0.0100 (10)
C30	0.0557 (17)	0.0482 (16)	0.0263 (13)	-0.0224 (13)	0.0021 (12)	-0.0150 (12)

Geometric parameters (Å, °)

S1—C7	1.664 (2)	C9—C10	1.393 (3)
S2—C15	1.659 (2)	C9—C14	1.396 (3)
O1—C8	1.234 (2)	C10—C11	1.386 (3)
O2—C16	1.232 (2)	C10—H10	0.9500
O3—C11	1.373 (2)	C11—C12	1.400 (3)
O3—C23	1.425 (3)	C12—C13	1.402 (3)
O4—C12	1.370 (2)	C13—C14	1.389 (3)
O4—C24	1.438 (2)	C14—H14	0.9500
O5—C13	1.375 (2)	C16—C17	1.491 (3)
O5—C25	1.423 (3)	C17—C18	1.395 (3)
O6—C19	1.361 (2)	C17—C22	1.397 (3)
O6—C26	1.433 (2)	C18—C19	1.393 (3)
O7—C20	1.374 (2)	C18—H18	0.9500
O7—C27	1.441 (3)	C19—C20	1.400 (3)
O8—C21	1.370 (2)	C20—C21	1.402 (3)
O8—C28	1.428 (2)	C21—C22	1.393 (3)
O9—C29	1.434 (3)	C22—H22	0.9500
O9—H9	0.80 (3)	C23—H23A	0.9800
N1—C7	1.342 (2)	C23—H23B	0.9800
N1—C1	1.433 (3)	C23—H23C	0.9800
N1—H1	0.89 (2)	C24—H24A	0.9800
N2—C15	1.344 (3)	C24—H24B	0.9800
N2—C2	1.425 (2)	C24—H24C	0.9800
N2—H2	0.88 (2)	C25—H25A	0.9800
N3—C8	1.375 (3)	C25—H25B	0.9800
N3—C7	1.403 (3)	C25—H25C	0.9800
N3—H3A	0.84 (2)	C26—H26A	0.9800
N4—C16	1.381 (3)	C26—H26B	0.9800

N4—C15	1.407 (2)	C26—H26C	0.9800
N4—H4A	0.90 (2)	C27—H27A	0.9800
C1—C6	1.388 (3)	C27—H27B	0.9800
C1—C2	1.398 (3)	C27—H27C	0.9800
C2—C3	1.393 (3)	C28—H28A	0.9800
C3—C4	1.391 (3)	C28—H28B	0.9800
C3—H3	0.9500	C28—H28C	0.9800
C4—C5	1.387 (3)	C29—C30	1.508 (3)
C4—H4	0.9500	C29—H29A	0.9900
C5—C6	1.388 (3)	C29—H29B	0.9900
C5—H5	0.9500	C30—H30A	0.9800
C6—H6	0.9500	C30—H30B	0.9800
C8—C9	1.493 (3)	C30—H30C	0.9800
C11—O3—C23	116.05 (16)	C18—C17—C22	121.48 (17)
C12—O4—C24	114.05 (15)	C18—C17—C16	116.36 (18)
C13—O5—C25	116.56 (17)	C22—C17—C16	122.13 (18)
C19—O6—C26	117.41 (16)	C19—C18—C17	119.18 (18)
C20—O7—C27	114.85 (15)	C19—C18—H18	120.4
C21—O8—C28	117.05 (15)	C17—C18—H18	120.4
C29—O9—H9	107.2 (19)	O6—C19—C18	124.44 (18)
C7—N1—C1	124.35 (17)	O6—C19—C20	115.37 (17)
C7—N1—H1	116.3 (15)	C18—C19—C20	120.19 (18)
C1—N1—H1	118.6 (15)	O7—C20—C19	121.59 (18)
C15—N2—C2	125.79 (17)	O7—C20—C21	118.60 (17)
C15—N2—H2	117.3 (15)	C19—C20—C21	119.73 (17)
C2—N2—H2	116.3 (15)	O8—C21—C22	124.37 (18)
C8—N3—C7	127.62 (17)	O8—C21—C20	115.14 (17)
C8—N3—H3A	118.5 (16)	C22—C21—C20	120.47 (18)
C7—N3—H3A	113.8 (16)	C21—C22—C17	118.83 (18)
C16—N4—C15	129.46 (17)	C21—C22—H22	120.6
C16—N4—H4A	116.0 (15)	C17—C22—H22	120.6
C15—N4—H4A	114.3 (15)	O3—C23—H23A	109.5
C6—C1—C2	120.34 (18)	O3—C23—H23B	109.5
C6—C1—N1	118.33 (18)	H23A—C23—H23B	109.5
C2—C1—N1	121.21 (17)	O3—C23—H23C	109.5
C3—C2—C1	119.20 (18)	H23A—C23—H23C	109.5
C3—C2—N2	121.58 (18)	H23B—C23—H23C	109.5
C1—C2—N2	119.21 (17)	O4—C24—H24A	109.5
C4—C3—C2	120.1 (2)	O4—C24—H24B	109.5
C4—C3—H3	120.0	H24A—C24—H24B	109.5
C2—C3—H3	120.0	O4—C24—H24C	109.5
C5—C4—C3	120.44 (19)	H24A—C24—H24C	109.5
C5—C4—H4	119.8	H24B—C24—H24C	109.5
C3—C4—H4	119.8	O5—C25—H25A	109.5
C4—C5—C6	119.65 (19)	O5—C25—H25B	109.5
C4—C5—H5	120.2	H25A—C25—H25B	109.5
C6—C5—H5	120.2	O5—C25—H25C	109.5
C5—C6—C1	120.21 (19)	H25A—C25—H25C	109.5
C5—C6—H6	119.9	H25B—C25—H25C	109.5

supplementary materials

C1—C6—H6	119.9	O6—C26—H26A	109.5
N1—C7—N3	115.67 (17)	O6—C26—H26B	109.5
N1—C7—S1	125.33 (15)	H26A—C26—H26B	109.5
N3—C7—S1	118.99 (14)	O6—C26—H26C	109.5
O1—C8—N3	121.57 (18)	H26A—C26—H26C	109.5
O1—C8—C9	121.22 (18)	H26B—C26—H26C	109.5
N3—C8—C9	117.16 (17)	O7—C27—H27A	109.5
C10—C9—C14	120.91 (19)	O7—C27—H27B	109.5
C10—C9—C8	115.91 (17)	H27A—C27—H27B	109.5
C14—C9—C8	123.07 (18)	O7—C27—H27C	109.5
C11—C10—C9	119.59 (18)	H27A—C27—H27C	109.5
C11—C10—H10	120.2	H27B—C27—H27C	109.5
C9—C10—H10	120.2	O8—C28—H28A	109.5
O3—C11—C10	124.62 (18)	O8—C28—H28B	109.5
O3—C11—C12	114.95 (18)	H28A—C28—H28B	109.5
C10—C11—C12	120.41 (19)	O8—C28—H28C	109.5
O4—C12—C11	120.25 (19)	H28A—C28—H28C	109.5
O4—C12—C13	120.29 (18)	H28B—C28—H28C	109.5
C11—C12—C13	119.24 (18)	O9—C29—C30	112.1 (2)
O5—C13—C14	124.23 (19)	O9—C29—H29A	109.2
O5—C13—C12	115.03 (18)	C30—C29—H29A	109.2
C14—C13—C12	120.71 (18)	O9—C29—H29B	109.2
C13—C14—C9	119.08 (19)	C30—C29—H29B	109.2
C13—C14—H14	120.5	H29A—C29—H29B	107.9
C9—C14—H14	120.5	C29—C30—H30A	109.5
N2—C15—N4	114.99 (17)	C29—C30—H30B	109.5
N2—C15—S2	128.09 (15)	H30A—C30—H30B	109.5
N4—C15—S2	116.90 (14)	C29—C30—H30C	109.5
O2—C16—N4	121.94 (18)	H30A—C30—H30C	109.5
O2—C16—C17	122.15 (18)	H30B—C30—H30C	109.5
N4—C16—C17	115.90 (17)		
C7—N1—C1—C6	112.1 (2)	C11—C12—C13—O5	-179.33 (17)
C7—N1—C1—C2	-71.9 (2)	O4—C12—C13—C14	-171.96 (17)
C6—C1—C2—C3	-2.4 (3)	C11—C12—C13—C14	2.6 (3)
N1—C1—C2—C3	-178.42 (17)	O5—C13—C14—C9	-178.68 (18)
C6—C1—C2—N2	178.93 (17)	C12—C13—C14—C9	-0.8 (3)
N1—C1—C2—N2	2.9 (3)	C10—C9—C14—C13	-1.2 (3)
C15—N2—C2—C3	46.2 (3)	C8—C9—C14—C13	174.92 (18)
C15—N2—C2—C1	-135.2 (2)	C2—N2—C15—N4	176.38 (18)
C1—C2—C3—C4	3.5 (3)	C2—N2—C15—S2	-2.4 (3)
N2—C2—C3—C4	-177.94 (18)	C16—N4—C15—N2	-11.5 (3)
C2—C3—C4—C5	-1.6 (3)	C16—N4—C15—S2	167.45 (17)
C3—C4—C5—C6	-1.3 (3)	C15—N4—C16—O2	4.6 (3)
C4—C5—C6—C1	2.3 (3)	C15—N4—C16—C17	-174.03 (19)
C2—C1—C6—C5	-0.5 (3)	O2—C16—C17—C18	-28.7 (3)
N1—C1—C6—C5	175.64 (18)	N4—C16—C17—C18	149.99 (18)
C1—N1—C7—N3	-178.21 (16)	O2—C16—C17—C22	149.5 (2)
C1—N1—C7—S1	1.0 (3)	N4—C16—C17—C22	-31.8 (3)
C8—N3—C7—N1	11.8 (3)	C22—C17—C18—C19	-0.3 (3)

C8—N3—C7—S1	-167.47 (15)	C16—C17—C18—C19	177.93 (18)
C7—N3—C8—O1	0.1 (3)	C26—O6—C19—C18	-5.4 (3)
C7—N3—C8—C9	177.21 (17)	C26—O6—C19—C20	174.43 (17)
O1—C8—C9—C10	15.1 (3)	C17—C18—C19—O6	176.90 (17)
N3—C8—C9—C10	-162.03 (17)	C17—C18—C19—C20	-2.9 (3)
O1—C8—C9—C14	-161.21 (19)	C27—O7—C20—C19	65.1 (3)
N3—C8—C9—C14	21.6 (3)	C27—O7—C20—C21	-118.3 (2)
C14—C9—C10—C11	1.4 (3)	O6—C19—C20—O7	0.9 (3)
C8—C9—C10—C11	-174.97 (17)	C18—C19—C20—O7	-179.27 (17)
C23—O3—C11—C10	-2.1 (3)	O6—C19—C20—C21	-175.70 (17)
C23—O3—C11—C12	176.46 (17)	C18—C19—C20—C21	4.1 (3)
C9—C10—C11—O3	178.88 (18)	C28—O8—C21—C22	4.0 (3)
C9—C10—C11—C12	0.4 (3)	C28—O8—C21—C20	-177.79 (17)
C24—O4—C12—C11	93.5 (2)	O7—C20—C21—O8	2.8 (3)
C24—O4—C12—C13	-92.0 (2)	C19—C20—C21—O8	179.51 (17)
O3—C11—C12—O4	-6.4 (3)	O7—C20—C21—C22	-178.87 (17)
C10—C11—C12—O4	172.17 (17)	C19—C20—C21—C22	-2.2 (3)
O3—C11—C12—C13	178.98 (17)	O8—C21—C22—C17	177.19 (17)
C10—C11—C12—C13	-2.4 (3)	C20—C21—C22—C17	-1.0 (3)
C25—O5—C13—C14	-5.1 (3)	C18—C17—C22—C21	2.2 (3)
C25—O5—C13—C12	176.94 (18)	C16—C17—C22—C21	-175.89 (18)
O4—C12—C13—O5	6.1 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N4—H4A \cdots O1 ⁱ	0.90 (2)	2.51 (2)	3.403 (2)	173 (2)
N2—H2 \cdots S1	0.88 (2)	2.69 (2)	3.3527 (19)	133.0 (18)
N2—H2 \cdots O2	0.88 (2)	1.97 (2)	2.677 (2)	136 (2)
N1—H1 \cdots O1	0.89 (2)	1.90 (2)	2.621 (2)	137 (2)
N3—H3A \cdots O9	0.84 (2)	2.23 (2)	2.949 (2)	145 (2)
O9—H9 \cdots O2	0.80 (3)	2.13 (3)	2.905 (2)	163 (3)

Symmetry codes: (i) $-x+2, -y+1, -z+1$.

Fig. 1




Fig. 2

