

trans-Tetraiodidobis(tri-*p*-tolylphosphine oxide- κO)tin(IV)

Marina Hitosugi-Levesque and Joseph M. Tanski*

Department of Chemistry, Vassar College, Poughkeepsie, NY 12604, USA
Correspondence e-mail: jtanski@vassar.edu


Received 15 July 2008; accepted 28 July 2008

Key indicators: single-crystal X-ray study; $T = 125$ K; mean $\sigma(C-C) = 0.003$ Å;
 R factor = 0.019; wR factor = 0.049; data-to-parameter ratio = 22.7.

The centrosymmetric title compound, $[SnI_4(C_{21}H_{21}OP)_2]$, is a monomeric complex that displays a nearly octahedral coordination of tin(IV), with an Sn–O bond distance of 2.159 (2) Å and an average Sn–I bond distance of 2.79 (3) Å.

Related literature

For examples of structurally characterized tin(IV) halide complexes of phosphine oxide ligands, see: Tursina *et al.* (1985); Tursina, Aslanov *et al.* (1986); Tursina, Yatsenko *et al.* (1986); Tudela *et al.* (1993); Genge *et al.* (1999); Szymanska-Buzar *et al.* (2001); Davis, Clarke *et al.* (2006); Davis, Levason *et al.* (2006); Caldwell & Tanski (2008). For related literature, see: Levason *et al.* (2003); Woollins (2003).



Experimental

Crystal data

$[SnI_4(C_{21}H_{21}OP)_2]$
 $M_r = 1266.99$
Orthorhombic, $Pbca$

$a = 18.6013$ (11) Å
 $b = 11.9677$ (7) Å
 $c = 19.3335$ (11) Å

$V = 4303.9$ (4) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 3.57$ mm⁻¹
 $T = 125$ (2) K
 $0.20 \times 0.20 \times 0.05$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.535$, $T_{\max} = 0.842$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.019$
 $wR(F^2) = 0.048$
 $S = 1.07$
5331 reflections

235 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.56$ e Å⁻³
 $\Delta\rho_{\min} = -0.49$ e Å⁻³

Table 1
Selected geometric parameters (Å, °).

Sn–O1	2.1590 (15)	Sn–I2	2.8158 (2)
Sn–I1	2.76735 (17)	O1–P1	1.5207 (16)
O1–Sn–O1 ⁱ	180	I1–Sn–I1 ⁱ	180
O1–Sn–I1	89.84 (4)	O1–Sn–I2 ⁱ	93.59 (4)
O1 ⁱ –Sn–I1	90.16 (4)	O1 ⁱ –Sn–I2 ⁱ	86.41 (4)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by Vassar College. X-ray facilities were provided by the US National Science Foundation (grant No. 0521237 to JMT).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2089).

References

- Bruker (2007). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Caldwell, A. M. & Tanski, J. M. (2008). *Acta Cryst. E64*, m894.
- Davis, M. F., Clarke, M., Levason, W., Reid, G. & Webster, M. (2006). *Eur. J. Inorg. Chem.*, pp. 2773–2782.
- Davis, M. F., Levason, W., Reid, G. & Webster, M. (2006). *Polyhedron*, **25**, 930–936.
- Genge, A. R. J., Levason, W. & Reid, G. (1999). *Inorg. Chim. Acta*, **288**, 142–149.
- Levason, W., Patel, R. & Reid, G. (2003). *J. Organomet. Chem.*, **688**, 280–282.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Szymanska-Buzar, T., Glowik, T. & Czelusniak, I. (2001). *Main Group Met. Chem.*, **24**, 821–822.
- Tudela, D., Tornero, J. D., Monge, A. & Sánchez-Herencia, A. J. (1993). *Inorg. Chem.*, **32**, 392, 3928–3930.
- Tursina, A. I., Aslanov, L. A., Chernyshev, V. V., Medvedev, S. V. & Yatsenko, A. V. (1986). *Koord. Khim.*, **12**, 420–424.
- Tursina, A. I., Aslanov, L. A., Medvedev, S. V. & Yatsenko, A. V. (1985). *Koord. Khim.*, **11**, 417–424.
- Tursina, A. I., Yatsenko, A. V., Medvedev, S. V., Chernyshev, V. V. & Aslanov, L. A. (1986). *Zh. Strukt. Khim.*, **27**, 157–159.
- Woollins, J. D. (2003). *Inorganic Experiments*, 2nd ed., edited by J. D. Woollins, pp. 36–37. Weinheim: Wiley.

supplementary materials

Acta Cryst. (2008). E64, m1107 [doi:10.1107/S1600536808023945]

***trans*-Tetraiodidobis(tri-*p*-tolylphosphine oxide- κO)tin(IV)**

M. Hitosugi-Levesque and J. M. Tanski

Comment

Tin(IV) iodide may be readily prepared by oxidation of tin metal with iodine (Woollins, 2003). A relatively weak Lewis acid, SnI_4 nevertheless forms complexes with phosphines and phosphine oxides (Genge *et al.* 1999; Davis, Clarke *et al.* 2006; Caldwell & Tanski, 2008). The phosphine oxide complexes are chiefly obtained by air oxidation of the phosphine ligands in the presence of the tin(IV) halide (Levason *et al.* 2003). The crystal and molecular structures of the bis(triphenyl phosphine oxide) adducts of SnX_4 , where $X = \text{F}, \text{Cl}, \text{Br}$ and I , have all been previously reported. In the case of the fluoride, *trans*- $[\text{Ph}_3\text{PO}]_2\text{SnF}_4$, the phosphine oxides are mutually *trans* (Davis, Clarke *et al.* 2006). The chloride, *cis*- $[\text{Ph}_3\text{PO}]_2\text{SnCl}_4$, exhibits *cis* phosphine oxides ligands (Tursina *et al.* 1985; Szymanska-Buzar *et al.* 2001), although it has been reported that both the *cis* and *trans* isomers are observed in solution by ^{31}P NMR (Davis, Levason *et al.* 2006<). The structures of both *cis*- $[\text{Ph}_3\text{PO}]_2\text{SnBr}_4$ and *trans*- $[\text{Ph}_3\text{PO}]_2\text{SnBr}_4$ are known for the bromide (Tudela *et al.* 1993; Tursina, Yatsenko *et al.* 1986). In the structure of the iodide, the triphenyl phosphine oxide ligands of *cis*- $[\text{Ph}_3\text{PO}]_2\text{SnI}_4$ are found to be *cis* (Tursina, Aslanov *et al.* 1986). As reported here, tri(*p*-tolyl) phosphine oxide results in an iodide complex wherein the phosphine oxide ligands are found to be *trans*.

Reaction of SnI_4 with tri(*p*-tolyl) phosphine, (*p*-CH₃C₆H₄)₃P, in CHCl₃ in the presence of air afforded the title complex [(*p*-CH₃C₆H₄)₃PO]₂SnI₄, (I).

Complex (I) exhibits a nearly octahedral coordination at tin, which resides on a crystallographic inversion center. The phosphine oxide ligands are mutually *trans*, with an Sn—O distance of 2.159 (2) Å, and Sn—I distances of 2.7674 (2) and 2.8158 (2) Å. Relevant bond lengths and angles can be found in Table 1. Despite the *p*-CH₃ substituent and *trans* orientation of the phosphine oxide ligands in (I), the Sn—O and Sn—I distances in (I) are very similar to those found in *cis*- $[\text{Ph}_3\text{PO}]_2\text{SnI}_4$, wherein the Sn—O distances are 2.15 (2) and 2.11 (2) Å, and the Sn—I distances range from 2.781 (2) to 2.816 (2) Å (Tursina, Aslanov *et al.* 1986).

Experimental

Complex (I) was prepared by treating a chloroform (*ca* 10 ml) solution of SnI_4 (626 mg, 1.00 mmol) with an excess of (*p*-CH₃C₆H₄)₃P (647 mg, 2.13 mmol) in the presence of air. Suitable crystals for single-crystal X-ray analysis separated as orange plates within 2 weeks at room temperature.

Refinement

H atoms on carbon atoms were included in calculated positions using a riding model at C—H distances 0.95 and 0.98 Å and U_{iso}(H) = 1.2 and 1.5U_{eq}(C) of the aryl and methyl C-atoms, respectively.

supplementary materials

Figures




Fig. 1. A view of complex (I) with displacement ellipsoids shown at the 50% probability level. H atoms have been omitted for clarity. Symmetry code: i = -x, -y+1, -z+1

trans-Tetraiodidobis(tri-*p*-tolylphosphine oxide- κ O)tin(IV)

Crystal data

[SnI ₄ (C ₂₁ H ₂₁ OP) ₂]	$F_{000} = 2408$
$M_r = 1266.99$	$D_x = 1.955 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
Hall symbol: -P2ac2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 18.6013 (11) \text{ \AA}$	Cell parameters from 9901 reflections
$b = 11.9677 (7) \text{ \AA}$	$\theta = 2.8\text{--}28.3^\circ$
$c = 19.3335 (11) \text{ \AA}$	$\mu = 3.57 \text{ mm}^{-1}$
$V = 4303.9 (4) \text{ \AA}^3$	$T = 125 (2) \text{ K}$
$Z = 4$	Plate, orange
	$0.20 \times 0.20 \times 0.05 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	5331 independent reflections
Radiation source: fine-focus sealed tube	4841 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.032$
$T = 125(2) \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
ϕ and ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -24 \rightarrow 24$
$T_{\text{min}} = 0.535$, $T_{\text{max}} = 0.842$	$k = -15 \rightarrow 15$
53839 measured reflections	$l = -25 \rightarrow 25$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.019$	H-atom parameters constrained

$wR(F^2) = 0.048$	$w = 1/[\sigma^2(F_o^2) + (0.0232P)^2 + 4.4313P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} = 0.001$
5331 reflections	$\Delta\rho_{\max} = 0.56 \text{ e } \text{\AA}^{-3}$
235 parameters	$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. A suitable crystal was mounted in a nylon loop with Paratone-*N* cryoprotectant oil and data was collected on a Bruker *APEX2* CCD platform diffractometer. The structure was solved using direct methods and standard difference map techniques, and was refined by full-matrix least-squares procedures on F^2 with *SHELXTL* Version 6.14 (Sheldrick, 2008). All non-hydrogen atoms were refined anisotropically. Refinement of F^2 against ALL reflections. Refinement of F^2 against ALL reflections. The weighted *R*-factor wR and goodness of fit *S* are based on F^2 , conventional *R*-factors *R* are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on F , and *R*-factors based on ALL data will be even larger. EXTI refined to zero and was removed from the refinement.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn	0.0000	0.5000	0.5000	0.01453 (5)
I1	0.071249 (8)	0.378968 (13)	0.399124 (8)	0.02266 (4)
I2	0.042286 (8)	0.699279 (12)	0.434117 (8)	0.02186 (4)
O1	-0.09443 (8)	0.50578 (13)	0.43516 (8)	0.0188 (3)
P1	-0.15379 (3)	0.44482 (5)	0.39642 (3)	0.01603 (11)
C1	-0.17934 (12)	0.52832 (19)	0.32327 (11)	0.0194 (4)
C2	-0.23667 (13)	0.4945 (2)	0.28165 (13)	0.0265 (5)
H2A	-0.2605	0.4259	0.2910	0.032*
C3	-0.25902 (13)	0.5601 (2)	0.22692 (13)	0.0266 (5)
H3A	-0.2976	0.5355	0.1985	0.032*
C4	-0.22576 (12)	0.6619 (2)	0.21282 (12)	0.0221 (5)
C5	-0.25292 (14)	0.7344 (2)	0.15496 (13)	0.0291 (5)
H5A	-0.2205	0.7982	0.1488	0.044*
H5B	-0.3012	0.7615	0.1663	0.044*
H5C	-0.2547	0.6908	0.1121	0.044*
C6	-0.16762 (13)	0.6945 (2)	0.25375 (13)	0.0242 (5)
H6A	-0.1433	0.7624	0.2438	0.029*
C7	-0.14473 (13)	0.6286 (2)	0.30905 (12)	0.0222 (5)
H7A	-0.1055	0.6522	0.3369	0.027*
C8	-0.23153 (12)	0.43203 (19)	0.45048 (11)	0.0181 (4)

supplementary materials

C9	-0.24193 (13)	0.5130 (2)	0.50135 (12)	0.0224 (5)
H9A	-0.2060	0.5679	0.5092	0.027*
C10	-0.30407 (13)	0.5139 (2)	0.54033 (13)	0.0245 (5)
H10A	-0.3103	0.5696	0.5748	0.029*
C11	-0.35776 (13)	0.4346 (2)	0.53003 (13)	0.0242 (5)
C12	-0.42613 (15)	0.4397 (2)	0.57128 (16)	0.0351 (6)
H12A	-0.4445	0.5165	0.5713	0.053*
H12B	-0.4166	0.4161	0.6189	0.053*
H12C	-0.4620	0.3899	0.5505	0.053*
C13	-0.34695 (13)	0.3523 (2)	0.47927 (13)	0.0238 (5)
H13A	-0.3827	0.2970	0.4719	0.029*
C14	-0.28479 (12)	0.3509 (2)	0.43988 (12)	0.0209 (4)
H14A	-0.2782	0.2948	0.4056	0.025*
C15	-0.12942 (12)	0.31033 (19)	0.36393 (11)	0.0181 (4)
C16	-0.13571 (12)	0.21535 (19)	0.40494 (12)	0.0204 (4)
H16A	-0.1573	0.2205	0.4494	0.024*
C17	-0.11054 (14)	0.1132 (2)	0.38123 (13)	0.0246 (5)
H17A	-0.1159	0.0486	0.4094	0.029*
C18	-0.07758 (14)	0.1039 (2)	0.31675 (13)	0.0275 (5)
C19	-0.05059 (18)	-0.0075 (3)	0.29219 (16)	0.0406 (7)
H19A	-0.0021	0.0011	0.2731	0.061*
H19B	-0.0828	-0.0365	0.2563	0.061*
H19C	-0.0492	-0.0599	0.3311	0.061*
C20	-0.07179 (14)	0.1986 (2)	0.27588 (13)	0.0293 (5)
H20A	-0.0499	0.1932	0.2316	0.035*
C21	-0.09738 (13)	0.3010 (2)	0.29838 (12)	0.0249 (5)
H21A	-0.0933	0.3649	0.2695	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn	0.01273 (9)	0.01808 (10)	0.01278 (9)	-0.00027 (7)	0.00002 (7)	0.00045 (7)
I1	0.02116 (8)	0.02856 (9)	0.01826 (7)	0.00429 (6)	0.00170 (5)	-0.00372 (6)
I2	0.02329 (8)	0.02232 (8)	0.01999 (8)	-0.00454 (6)	0.00017 (6)	0.00311 (6)
O1	0.0151 (7)	0.0217 (8)	0.0195 (8)	-0.0017 (6)	-0.0029 (6)	0.0009 (6)
P1	0.0144 (2)	0.0187 (3)	0.0150 (3)	-0.0020 (2)	-0.0015 (2)	0.0013 (2)
C1	0.0182 (10)	0.0223 (11)	0.0177 (10)	-0.0002 (9)	-0.0020 (8)	0.0027 (8)
C2	0.0247 (12)	0.0271 (12)	0.0276 (12)	-0.0088 (10)	-0.0078 (10)	0.0076 (10)
C3	0.0203 (11)	0.0366 (14)	0.0229 (12)	-0.0047 (10)	-0.0068 (9)	0.0041 (10)
C4	0.0193 (10)	0.0288 (12)	0.0182 (11)	0.0042 (9)	0.0010 (9)	0.0037 (9)
C5	0.0272 (12)	0.0343 (14)	0.0259 (12)	0.0038 (11)	-0.0023 (10)	0.0100 (11)
C6	0.0237 (11)	0.0236 (11)	0.0253 (12)	-0.0030 (9)	0.0001 (9)	0.0056 (9)
C7	0.0202 (11)	0.0246 (12)	0.0216 (11)	-0.0041 (9)	-0.0039 (9)	0.0009 (9)
C8	0.0158 (10)	0.0217 (11)	0.0169 (10)	-0.0011 (8)	-0.0010 (8)	0.0022 (8)
C9	0.0197 (11)	0.0244 (11)	0.0231 (11)	-0.0022 (9)	-0.0029 (9)	-0.0030 (9)
C10	0.0249 (11)	0.0270 (12)	0.0215 (11)	0.0029 (10)	0.0007 (9)	-0.0019 (9)
C11	0.0209 (11)	0.0264 (12)	0.0254 (12)	0.0024 (9)	0.0041 (9)	0.0062 (10)
C12	0.0284 (13)	0.0333 (14)	0.0436 (16)	0.0016 (11)	0.0146 (12)	-0.0003 (12)

C13	0.0200 (11)	0.0226 (11)	0.0289 (12)	-0.0052 (9)	-0.0005 (9)	0.0041 (10)
C14	0.0197 (11)	0.0209 (11)	0.0220 (11)	-0.0013 (9)	-0.0012 (9)	-0.0009 (9)
C15	0.0169 (10)	0.0214 (11)	0.0161 (10)	-0.0022 (8)	-0.0013 (8)	-0.0020 (8)
C16	0.0213 (11)	0.0228 (11)	0.0172 (10)	-0.0007 (9)	-0.0009 (9)	-0.0010 (8)
C17	0.0297 (12)	0.0204 (11)	0.0236 (12)	-0.0003 (9)	-0.0027 (10)	-0.0011 (9)
C18	0.0270 (12)	0.0315 (13)	0.0239 (12)	0.0038 (10)	-0.0072 (10)	-0.0113 (10)
C19	0.0497 (17)	0.0370 (16)	0.0350 (15)	0.0097 (13)	-0.0074 (13)	-0.0184 (12)
C20	0.0285 (13)	0.0412 (15)	0.0183 (11)	0.0026 (11)	0.0011 (10)	-0.0072 (10)
C21	0.0251 (12)	0.0301 (13)	0.0193 (11)	-0.0016 (10)	0.0018 (9)	-0.0005 (10)

Geometric parameters (\AA , $^\circ$)

Sn—O1	2.1590 (15)	C9—C10	1.380 (3)
Sn—O1 ⁱ	2.1591 (15)	C9—H9A	0.9500
Sn—I1	2.76735 (17)	C10—C11	1.392 (3)
Sn—I1 ⁱ	2.76737 (17)	C10—H10A	0.9500
Sn—I2 ⁱ	2.81580 (19)	C11—C13	1.404 (4)
Sn—I2	2.8158 (2)	C11—C12	1.502 (3)
O1—P1	1.5207 (16)	C12—H12A	0.9800
P1—C15	1.786 (2)	C12—H12B	0.9800
P1—C8	1.791 (2)	C12—H12C	0.9800
P1—C1	1.796 (2)	C13—C14	1.385 (3)
C1—C7	1.389 (3)	C13—H13A	0.9500
C1—C2	1.396 (3)	C14—H14A	0.9500
C2—C3	1.382 (3)	C15—C16	1.391 (3)
C2—H2A	0.9500	C15—C21	1.405 (3)
C3—C4	1.393 (4)	C16—C17	1.387 (3)
C3—H3A	0.9500	C16—H16A	0.9500
C4—C6	1.396 (3)	C17—C18	1.394 (4)
C4—C5	1.503 (3)	C17—H17A	0.9500
C5—H5A	0.9800	C18—C20	1.387 (4)
C5—H5B	0.9800	C18—C19	1.502 (4)
C5—H5C	0.9800	C19—H19A	0.9800
C6—C7	1.395 (3)	C19—H19B	0.9800
C6—H6A	0.9500	C19—H19C	0.9800
C7—H7A	0.9500	C20—C21	1.384 (4)
C8—C9	1.394 (3)	C20—H20A	0.9500
C8—C14	1.402 (3)	C21—H21A	0.9500
O1—Sn—O1 ⁱ	180.0	C9—C8—P1	117.67 (17)
O1—Sn—I1	89.84 (4)	C14—C8—P1	122.98 (17)
O1—Sn—I1	89.84 (4)	C10—C9—C8	120.5 (2)
O1 ⁱ —Sn—I1	90.16 (4)	C10—C9—H9A	119.8
O1—Sn—I1 ⁱ	90.16 (4)	C8—C9—H9A	119.8
O1 ⁱ —Sn—I1 ⁱ	89.84 (4)	C9—C10—C11	121.2 (2)
I1—Sn—I1 ⁱ	180.0	C9—C10—H10A	119.4
O1—Sn—I2 ⁱ	93.59 (4)	C11—C10—H10A	119.4
O1 ⁱ —Sn—I2 ⁱ	86.41 (4)	C10—C11—C13	118.4 (2)

supplementary materials

I1—Sn—I2 ⁱ	90.529 (6)	C10—C11—C12	120.2 (2)
I1 ⁱ —Sn—I2 ⁱ	89.471 (6)	C13—C11—C12	121.4 (2)
O1—Sn—I2	86.41 (4)	C11—C12—H12A	109.5
O1 ⁱ —Sn—I2	93.59 (4)	C11—C12—H12B	109.5
I1—Sn—I2	89.469 (6)	H12A—C12—H12B	109.5
I1 ⁱ —Sn—I2	90.531 (6)	C11—C12—H12C	109.5
I2 ⁱ —Sn—I2	180.0	H12A—C12—H12C	109.5
P1—O1—Sn	149.47 (10)	H12B—C12—H12C	109.5
O1—P1—C15	114.91 (10)	C14—C13—C11	120.8 (2)
O1—P1—C8	109.87 (10)	C14—C13—H13A	119.6
C15—P1—C8	109.46 (11)	C11—C13—H13A	119.6
O1—P1—C1	108.25 (10)	C13—C14—C8	120.1 (2)
C15—P1—C1	106.96 (11)	C13—C14—H14A	120.0
C8—P1—C1	107.07 (10)	C8—C14—H14A	120.0
C7—C1—C2	119.4 (2)	C16—C15—C21	119.0 (2)
C7—C1—P1	120.92 (17)	C16—C15—P1	120.96 (17)
C2—C1—P1	119.66 (17)	C21—C15—P1	119.77 (18)
C3—C2—C1	120.4 (2)	C17—C16—C15	120.2 (2)
C3—C2—H2A	119.8	C17—C16—H16A	119.9
C1—C2—H2A	119.8	C15—C16—H16A	119.9
C2—C3—C4	120.9 (2)	C16—C17—C18	121.0 (2)
C2—C3—H3A	119.6	C16—C17—H17A	119.5
C4—C3—H3A	119.6	C18—C17—H17A	119.5
C3—C4—C6	118.5 (2)	C20—C18—C17	118.6 (2)
C3—C4—C5	120.1 (2)	C20—C18—C19	121.3 (2)
C6—C4—C5	121.4 (2)	C17—C18—C19	120.1 (3)
C4—C5—H5A	109.5	C18—C19—H19A	109.5
C4—C5—H5B	109.5	C18—C19—H19B	109.5
H5A—C5—H5B	109.5	H19A—C19—H19B	109.5
C4—C5—H5C	109.5	C18—C19—H19C	109.5
H5A—C5—H5C	109.5	H19A—C19—H19C	109.5
H5B—C5—H5C	109.5	H19B—C19—H19C	109.5
C7—C6—C4	120.9 (2)	C21—C20—C18	121.2 (2)
C7—C6—H6A	119.6	C21—C20—H20A	119.4
C4—C6—H6A	119.6	C18—C20—H20A	119.4
C1—C7—C6	119.9 (2)	C20—C21—C15	120.0 (2)
C1—C7—H7A	120.1	C20—C21—H21A	120.0
C6—C7—H7A	120.0	C15—C21—H21A	120.0
C9—C8—C14	119.1 (2)		
I1—Sn—O1—P1	60.78 (19)	C1—P1—C8—C14	84.5 (2)
I1 ⁱ —Sn—O1—P1	-119.22 (19)	C14—C8—C9—C10	-0.6 (3)
I2 ⁱ —Sn—O1—P1	-29.74 (19)	P1—C8—C9—C10	173.90 (18)
I2—Sn—O1—P1	150.26 (19)	C8—C9—C10—C11	0.0 (4)
Sn—O1—P1—C15	-31.2 (2)	C9—C10—C11—C13	0.7 (4)
Sn—O1—P1—C8	92.7 (2)	C9—C10—C11—C12	-177.7 (2)
Sn—O1—P1—C1	-150.70 (18)	C10—C11—C13—C14	-0.7 (4)
O1—P1—C1—C7	1.8 (2)	C12—C11—C13—C14	177.7 (2)

C15—P1—C1—C7	-122.5 (2)	C11—C13—C14—C8	0.1 (4)
C8—P1—C1—C7	120.2 (2)	C9—C8—C14—C13	0.5 (3)
O1—P1—C1—C2	-175.57 (19)	P1—C8—C14—C13	-173.65 (18)
C15—P1—C1—C2	60.1 (2)	O1—P1—C15—C16	86.7 (2)
C8—P1—C1—C2	-57.2 (2)	C8—P1—C15—C16	-37.4 (2)
C7—C1—C2—C3	-0.2 (4)	C1—P1—C15—C16	-153.10 (19)
P1—C1—C2—C3	177.2 (2)	O1—P1—C15—C21	-87.5 (2)
C1—C2—C3—C4	-1.1 (4)	C8—P1—C15—C21	148.36 (18)
C2—C3—C4—C6	2.2 (4)	C1—P1—C15—C21	32.7 (2)
C2—C3—C4—C5	-177.3 (2)	C21—C15—C16—C17	0.1 (3)
C3—C4—C6—C7	-2.2 (4)	P1—C15—C16—C17	-174.16 (18)
C5—C4—C6—C7	177.4 (2)	C15—C16—C17—C18	1.1 (4)
C2—C1—C7—C6	0.3 (4)	C16—C17—C18—C20	-1.4 (4)
P1—C1—C7—C6	-177.14 (19)	C16—C17—C18—C19	179.9 (2)
C4—C6—C7—C1	0.9 (4)	C17—C18—C20—C21	0.6 (4)
O1—P1—C8—C9	27.6 (2)	C19—C18—C20—C21	179.3 (3)
C15—P1—C8—C9	154.69 (18)	C18—C20—C21—C15	0.5 (4)
C1—P1—C8—C9	-89.72 (19)	C16—C15—C21—C20	-0.9 (3)
O1—P1—C8—C14	-158.13 (18)	P1—C15—C21—C20	173.47 (19)
C15—P1—C8—C14	-31.1 (2)		

Symmetry codes: (i) $-x, -y+1, -z+1$.

supplementary materials

Fig. 1

