

Bis[3-allyl-1-(4-cyanobenzyl)-2-methylbenzimidazolium] di- μ -bromido-bis-[bromidocuprate(I)]

Guang-Hai Xu and Wei Wang*

Ordered Matter Science Research Center, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: seuwangwei@gmail.com


Received 13 August 2008; accepted 20 August 2008

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.052; wR factor = 0.134; data-to-parameter ratio = 20.0.

The asymmetric unit of the title compound, $(\text{C}_{19}\text{H}_{18}\text{N}_3)_2[\text{Cu}_2\text{Br}_4]$, contains one cation and one half-anion; there is a centre of symmetry mid-way between the two Cu atoms. In the cation, the nearly planar benzimidazole ring system is oriented at dihedral angles of 75.31 (3) and 21.39 (3)° with respect to the cyanobenzyl and allyl groups, respectively. The dihedral angle between cyanobenzyl and allyl groups is 87.94 (3)°. In the crystal structure, intermolecular C–H···Br hydrogen bonds link the molecules. There is a C–H···π contact between the cyanobenzyl ring and the anion; π–π contacts also exist between the benzimidazole ring systems as well as between the anion and the cyanobenzyl ring [centroid–centroid distances = 4.024 (1) and 4.617 (1) Å, respectively].

Related literature

For related literature, see: Aaker *et al.* (2005).



Experimental

Crystal data

$(\text{C}_{19}\text{H}_{18}\text{N}_3)_2[\text{Cu}_2\text{Br}_4]$	$\gamma = 81.41 (2)^\circ$
$M_r = 1023.44$	$V = 998.93 (18)$ Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.6407 (5)$ Å	Mo $K\alpha$ radiation
$b = 10.1029 (11)$ Å	$\mu = 5.10$ mm ⁻¹
$c = 11.0389 (6)$ Å	$T = 294 (2)$ K
$\alpha = 87.27 (2)^\circ$	$0.20 \times 0.20 \times 0.20$ mm
$\beta = 69.990 (17)^\circ$	

Data collection

Rigaku, SCXmini diffractometer	10227 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku/MSC, 2005)	4517 independent reflections
	2765 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.052$
	$T_{\min} = 0.361$, $T_{\max} = 0.375$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	226 parameters
$wR(F^2) = 0.134$	H-atom parameters constrained
$S = 0.97$	$\Delta\rho_{\max} = 1.06$ e Å ⁻³
4517 reflections	$\Delta\rho_{\min} = -0.76$ e Å ⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
C8–H8A···Br1 ⁱ	0.97	2.90	3.863 (5)	175
C10–H10C···Br1	0.96	2.93	3.788 (5)	150
C17–H17A···Br1 ⁱⁱ	0.93	2.92	3.707 (5)	143
C4–H4a···Cg1 ⁱⁱⁱ	0.93	3.36	3.746 (2)	108 (1)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x, -y + 1, -z + 1$; (iii) $x, y, z - 1$. Cg1 is the centroid of the anion.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku/MSC, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful to the Starter Fund of Southeast University for financial support to purchase the CCD X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2513).

References

- Aaker, C. B., Desper, J. & Urbinam, J. F. (2005). *J. Cryst. Growth Des.* **5**, 1283–1293.
Rigaku/MSC. (2005). *CrystalClear* and *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1198 [doi:10.1107/S1600536808026937]

Bis[3-allyl-1-(4-cyanobenzyl)-2-methylbenzimidazolium] di- μ -bromido-bis[bromidocuprate(I)]

G.-H. Xu and W. Wang

Comment

The asymmetric unit of the title compound contains one cation and one-half anion (Fig. 1). In the cation, rings A (N2/N3/C9/C14/C19), B (C14—C19) and C (C2—C7) are, of course, planar and the dihedral angles between them are A/B = 1.47 (3) $^{\circ}$, A/C = 75.18 (4) $^{\circ}$ and B/C = 75.46 (4) $^{\circ}$. So, the benzimidazole ring system is nearly planar, and it is oriented with respect to the cyanobenzyl and allyl groups at dihedral angles of 75.31 (3) $^{\circ}$ and 21.39 (3) $^{\circ}$, respectively. The dihedral angle between cyanobenzyl and allyl groups is 87.94 (3) $^{\circ}$.

In the crystal structure, intermolecular C—H \cdots Br hydrogen bonds link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. There also exists a C—H \cdots π contact (Table 1) between the cyanobenzyl ring and the anion. The π — π contacts between the rings A and B as well as the anion and the ring C, $Cg2\cdots Cg4^i$ and $Cg3\cdots Cg1^{ii}$ [symmetry codes: (i) $-x, 1 - y, 1 - z$; (ii) $x, y, z - 1$, where $Cg1, Cg2, Cg3$ and $Cg4$ are centroids of the anion, the rings A, C and B, respectively] further stabilize the structure, with centroid-centroid distances of 4.024 (1) and 4.617 (1) Å, respectively.

Experimental

The synthesis of (1) (Scheme 1) was reported, previously (Aaker *et al.*, 2005). The mixture of (1) (2.47 g, 10 mmol), allyl bromide (1.21 g, 10 mmol) and Na_2CO_3 (0.53 g, 5 mmol) was stirred in THF (50 ml) under reflux for 24 h to obtain (2) (Scheme 1) with high yield (90%). The colorless crystals suitable for X-ray analysis were obtained from the mixture of (2) (73.6 mg, 0.2 mmol), CuBr (28.8 mg, 0.2 mmol) and methanol (2 ml) sealed in a glass tube maintaining at 353 K for 5 d.

Refinement

H atoms were positioned geometrically, with C—H = 0.93, 0.97 and 0.96 Å for aromatic and methylene (for allyl C13 atom), methylene and methyl H, respectively, and constrained to ride on their parent atoms with $U_{iso}(H) = xU_{eq}(C)$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level [symmetry code: (A) $-x, 1 - y, 2 - z$].

supplementary materials




Fig. 2. A partial packing diagram of the title compound, viewed along the a axis. Hydrogen bonds are shown as dashed lines.




Fig. 3. The formation of the title compound.

Bis[3-allyl-1-(4-cyanobenzyl)-2-methylbenzimidazolium] di- μ -bromido-bis[bromidocuprate(I)]

Crystal data

(C ₁₉ H ₁₈ N ₃) ₂ [Cu ₂ Br ₄]	$Z = 1$
$M_r = 1023.44$	$F_{000} = 504$
Triclinic, $P\bar{1}$	$D_x = 1.701 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 9.6407 (5) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 10.1029 (11) \text{ \AA}$	Cell parameters from 2257 reflections
$c = 11.0389 (6) \text{ \AA}$	$\theta = 2.8\text{--}27.5^\circ$
$\alpha = 87.27 (2)^\circ$	$\mu = 5.10 \text{ mm}^{-1}$
$\beta = 69.990 (17)^\circ$	$T = 294 (2) \text{ K}$
$\gamma = 81.41 (2)^\circ$	Prism, colorless
$V = 998.93 (18) \text{ \AA}^3$	$0.20 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Rigaku, SCXmini diffractometer	4517 independent reflections
Radiation source: fine-focus sealed tube	2765 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.052$
Detector resolution: 13.6612 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ$
$T = 294(2) \text{ K}$	$\theta_{\text{min}} = 2.8^\circ$
ω scans	$h = -12 \rightarrow 12$
Absorption correction: multi-scan (CrystalClear; Rigaku/MSC, 2005)	$k = -13 \rightarrow 13$
$T_{\text{min}} = 0.361, T_{\text{max}} = 0.375$	$l = -14 \rightarrow 14$
10227 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.052$	H-atom parameters constrained
$wR(F^2) = 0.134$	$w = 1/[\sigma^2(F_o^2) + (0.0636P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.97$	$(\Delta/\sigma)_{\max} < 0.001$
4517 reflections	$\Delta\rho_{\max} = 1.07 \text{ e \AA}^{-3}$
226 parameters	$\Delta\rho_{\min} = -0.76 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.37052 (6)	0.42643 (5)	0.73582 (5)	0.05337 (18)
Br2	-0.06239 (6)	0.34357 (6)	0.94360 (6)	0.0684 (2)
Cu1	0.14770 (8)	0.47171 (8)	0.90578 (7)	0.0703 (3)
N1	0.5950 (7)	0.0085 (5)	-0.2925 (5)	0.0873 (18)
N2	0.2273 (4)	0.3449 (4)	0.3653 (4)	0.0413 (9)
N3	0.1992 (4)	0.2221 (4)	0.5394 (3)	0.0406 (9)
C1	0.5454 (7)	0.0689 (6)	-0.1995 (6)	0.0586 (14)
C2	0.4844 (6)	0.1517 (5)	-0.0842 (5)	0.0486 (12)
C3	0.3336 (6)	0.1632 (5)	-0.0149 (5)	0.0554 (14)
H3A	0.2729	0.1148	-0.0396	0.067*
C4	0.2729 (6)	0.2476 (5)	0.0920 (5)	0.0511 (13)
H4A	0.1714	0.2562	0.1386	0.061*
C5	0.3641 (5)	0.3188 (5)	0.1291 (4)	0.0428 (11)
C6	0.5157 (5)	0.3059 (5)	0.0593 (5)	0.0522 (13)
H6A	0.5771	0.3529	0.0847	0.063*
C7	0.5758 (6)	0.2230 (6)	-0.0484 (5)	0.0551 (13)
H7A	0.6770	0.2155	-0.0960	0.066*
C8	0.2982 (6)	0.4137 (5)	0.2437 (4)	0.0468 (12)
H8A	0.3767	0.4580	0.2529	0.056*
H8B	0.2246	0.4819	0.2274	0.056*
C9	0.2985 (5)	0.2755 (5)	0.4384 (4)	0.0411 (11)
C10	0.4630 (5)	0.2599 (6)	0.4119 (5)	0.0538 (13)

supplementary materials

H10A	0.5095	0.3072	0.3345	0.081*
H10B	0.5034	0.1667	0.4014	0.081*
H10C	0.4818	0.2959	0.4829	0.081*
C11	0.2307 (6)	0.1405 (5)	0.6424 (5)	0.0511 (12)
H11A	0.1753	0.1848	0.7243	0.061*
H11B	0.3360	0.1347	0.6301	0.061*
C12	0.1921 (7)	0.0028 (6)	0.6483 (6)	0.0634 (15)
H12A	0.2200	-0.0443	0.5713	0.076*
C13	0.1227 (9)	-0.0555 (7)	0.7528 (8)	0.105 (3)
H13C	0.0932	-0.0112	0.8314	0.126*
H13A	0.1019	-0.1421	0.7498	0.126*
C14	0.0569 (5)	0.2589 (5)	0.5305 (4)	0.0397 (10)
C15	-0.0827 (5)	0.2331 (5)	0.6120 (5)	0.0496 (12)
H15A	-0.0947	0.1818	0.6861	0.060*
C16	-0.2025 (6)	0.2887 (6)	0.5754 (5)	0.0563 (14)
H16A	-0.2980	0.2738	0.6261	0.068*
C17	-0.1844 (6)	0.3659 (5)	0.4657 (5)	0.0539 (13)
H17A	-0.2686	0.4011	0.4456	0.065*
C18	-0.0473 (5)	0.3928 (5)	0.3850 (5)	0.0497 (12)
H18A	-0.0365	0.4449	0.3114	0.060*
C19	0.0746 (5)	0.3368 (5)	0.4208 (4)	0.0404 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0453 (3)	0.0604 (3)	0.0527 (3)	-0.0150 (2)	-0.0116 (2)	0.0021 (3)
Br2	0.0466 (3)	0.0807 (4)	0.0750 (4)	-0.0173 (3)	-0.0105 (3)	-0.0237 (3)
Cu1	0.0549 (5)	0.0929 (6)	0.0560 (4)	-0.0042 (4)	-0.0119 (3)	-0.0038 (4)
N1	0.114 (5)	0.076 (4)	0.059 (3)	-0.020 (3)	-0.007 (3)	-0.020 (3)
N2	0.037 (2)	0.048 (2)	0.037 (2)	-0.0122 (17)	-0.0077 (16)	-0.0018 (18)
N3	0.040 (2)	0.047 (2)	0.035 (2)	-0.0093 (17)	-0.0115 (17)	-0.0010 (18)
C1	0.071 (4)	0.054 (3)	0.053 (3)	-0.016 (3)	-0.021 (3)	0.001 (3)
C2	0.061 (3)	0.045 (3)	0.039 (3)	-0.008 (2)	-0.015 (2)	-0.002 (2)
C3	0.068 (4)	0.058 (3)	0.052 (3)	-0.025 (3)	-0.029 (3)	0.002 (3)
C4	0.044 (3)	0.062 (3)	0.046 (3)	-0.015 (2)	-0.011 (2)	-0.004 (3)
C5	0.049 (3)	0.045 (3)	0.036 (2)	-0.016 (2)	-0.013 (2)	0.004 (2)
C6	0.039 (3)	0.070 (4)	0.049 (3)	-0.020 (2)	-0.011 (2)	-0.009 (3)
C7	0.046 (3)	0.071 (4)	0.046 (3)	-0.014 (3)	-0.010 (2)	-0.008 (3)
C8	0.051 (3)	0.049 (3)	0.038 (3)	-0.014 (2)	-0.010 (2)	-0.001 (2)
C9	0.042 (3)	0.044 (3)	0.039 (3)	-0.007 (2)	-0.014 (2)	-0.005 (2)
C10	0.037 (3)	0.072 (4)	0.053 (3)	-0.014 (2)	-0.013 (2)	-0.001 (3)
C11	0.050 (3)	0.054 (3)	0.055 (3)	-0.013 (2)	-0.023 (2)	0.009 (3)
C12	0.075 (4)	0.057 (3)	0.060 (4)	-0.007 (3)	-0.027 (3)	0.003 (3)
C13	0.132 (7)	0.075 (5)	0.116 (7)	-0.046 (5)	-0.041 (6)	0.022 (5)
C14	0.035 (2)	0.044 (3)	0.039 (3)	-0.005 (2)	-0.0107 (19)	-0.007 (2)
C15	0.039 (3)	0.056 (3)	0.046 (3)	-0.005 (2)	-0.005 (2)	-0.002 (2)
C16	0.033 (3)	0.068 (4)	0.060 (3)	-0.009 (2)	-0.002 (2)	-0.010 (3)
C17	0.043 (3)	0.059 (3)	0.062 (3)	-0.003 (2)	-0.022 (3)	-0.004 (3)

C18	0.044 (3)	0.057 (3)	0.051 (3)	-0.006 (2)	-0.019 (2)	-0.002 (3)
C19	0.038 (3)	0.040 (3)	0.042 (3)	-0.007 (2)	-0.011 (2)	-0.006 (2)

Geometric parameters (\AA , $^\circ$)

Br1—Cu1	2.3181 (10)	C8—H8A	0.9700
Br2—Cu1 ⁱ	2.4134 (11)	C8—H8B	0.9700
Br2—Cu1	2.4710 (10)	C9—C10	1.496 (6)
Cu1—Br2 ⁱ	2.4134 (11)	C10—H10A	0.9600
Cu1—Cu1 ⁱ	2.8874 (15)	C10—H10B	0.9600
N1—C1	1.136 (7)	C10—H10C	0.9600
N2—C8	1.476 (6)	C11—C12	1.486 (7)
N2—C9	1.340 (6)	C11—H11A	0.9700
N2—C19	1.401 (5)	C11—H11B	0.9700
N3—C9	1.347 (6)	C12—C13	1.285 (8)
N3—C11	1.462 (6)	C12—H12A	0.9300
N3—C14	1.402 (6)	C13—H13C	0.9300
C2—C1	1.452 (7)	C13—H13A	0.9300
C2—C3	1.381 (7)	C14—C15	1.394 (6)
C2—C7	1.381 (7)	C15—H15A	0.9300
C3—C4	1.392 (7)	C16—C15	1.384 (7)
C3—H3A	0.9300	C16—C17	1.382 (7)
C4—H4A	0.9300	C16—H16A	0.9300
C5—C4	1.385 (6)	C17—H17A	0.9300
C5—C6	1.389 (6)	C18—C17	1.376 (7)
C5—C8	1.519 (6)	C18—H18A	0.9300
C6—H6A	0.9300	C19—C14	1.387 (6)
C7—C6	1.389 (7)	C19—C18	1.398 (7)
C7—H7A	0.9300		
Cu1 ⁱ —Br2—Cu1	72.46 (3)	N2—C9—C10	125.1 (4)
Br1—Cu1—Br2 ⁱ	129.22 (4)	N3—C9—C10	125.3 (4)
Br1—Cu1—Br2	122.83 (4)	C9—C10—H10A	109.5
Br2 ⁱ —Cu1—Br2	107.54 (3)	C9—C10—H10B	109.5
Br1—Cu1—Cu1 ⁱ	172.96 (5)	H10A—C10—H10B	109.5
Br2 ⁱ —Cu1—Cu1 ⁱ	54.69 (3)	C9—C10—H10C	109.5
Br2—Cu1—Cu1 ⁱ	52.85 (3)	H10A—C10—H10C	109.5
C9—N2—C19	108.6 (4)	H10B—C10—H10C	109.5
C9—N2—C8	125.8 (4)	N3—C11—C12	113.8 (4)
C19—N2—C8	125.5 (4)	N3—C11—H11A	108.8
C9—N3—C14	108.3 (4)	C12—C11—H11A	108.8
C9—N3—C11	127.0 (4)	N3—C11—H11B	108.8
C14—N3—C11	124.8 (4)	C12—C11—H11B	108.8
N1—C1—C2	177.3 (6)	H11A—C11—H11B	107.7
C7—C2—C3	120.7 (4)	C13—C12—C11	124.3 (6)
C7—C2—C1	119.9 (5)	C13—C12—H12A	117.8
C3—C2—C1	119.4 (5)	C11—C12—H12A	117.8
C2—C3—C4	119.8 (5)	C12—C13—H13C	120.0

supplementary materials

C2—C3—H3A	120.1	C12—C13—H13A	120.0
C4—C3—H3A	120.1	H13C—C13—H13A	120.0
C5—C4—C3	119.9 (5)	C19—C14—C15	122.2 (4)
C5—C4—H4A	120.1	C19—C14—N3	106.8 (4)
C3—C4—H4A	120.1	C15—C14—N3	130.9 (5)
C4—C5—C6	119.9 (4)	C16—C15—C14	115.7 (5)
C4—C5—C8	120.3 (4)	C16—C15—H15A	122.2
C6—C5—C8	119.8 (4)	C14—C15—H15A	122.2
C7—C6—C5	120.1 (4)	C17—C16—C15	121.9 (5)
C7—C6—H6A	119.9	C17—C16—H16A	119.0
C5—C6—H6A	119.9	C15—C16—H16A	119.0
C2—C7—C6	119.6 (5)	C18—C17—C16	122.9 (5)
C2—C7—H7A	120.2	C18—C17—H17A	118.5
C6—C7—H7A	120.2	C16—C17—H17A	118.5
N2—C8—C5	112.7 (4)	C17—C18—C19	115.6 (5)
N2—C8—H8A	109.0	C17—C18—H18A	122.2
C5—C8—H8A	109.0	C19—C18—H18A	122.2
N2—C8—H8B	109.0	C14—C19—C18	121.6 (4)
C5—C8—H8B	109.0	C14—C19—N2	106.7 (4)
H8A—C8—H8B	107.8	C18—C19—N2	131.7 (4)
N2—C9—N3	109.6 (4)		
Cu1 ⁱ —Br2—Cu1—Br1	-173.22 (6)	C7—C2—C3—C4	0.1 (8)
Cu1 ⁱ —Br2—Cu1—Br2 ⁱ	0.0	C1—C2—C7—C6	177.6 (5)
C9—N2—C8—C5	79.9 (6)	C3—C2—C7—C6	0.7 (8)
C19—N2—C8—C5	-97.3 (5)	C2—C3—C4—C5	-0.5 (8)
C8—N2—C9—N3	-177.6 (4)	C6—C5—C4—C3	0.1 (8)
C19—N2—C9—N3	0.0 (5)	C8—C5—C4—C3	178.5 (5)
C8—N2—C9—C10	2.5 (7)	C4—C5—C6—C7	0.7 (8)
C19—N2—C9—C10	-179.9 (4)	C8—C5—C6—C7	-177.7 (5)
C8—N2—C19—C14	177.4 (4)	C4—C5—C8—N2	61.6 (6)
C9—N2—C19—C14	-0.2 (5)	C6—C5—C8—N2	-119.9 (5)
C8—N2—C19—C18	-3.9 (8)	C2—C7—C6—C5	-1.1 (8)
C9—N2—C19—C18	178.5 (5)	N3—C11—C12—C13	-135.1 (7)
C11—N3—C9—N2	-179.4 (4)	N3—C14—C15—C16	178.1 (5)
C14—N3—C9—N2	0.2 (5)	C19—C14—C15—C16	0.8 (7)
C11—N3—C9—C10	0.5 (7)	C15—C16—C17—C18	0.2 (8)
C14—N3—C9—C10	-179.9 (5)	C17—C16—C15—C14	-0.6 (8)
C9—N3—C11—C12	-117.9 (5)	N2—C19—C14—N3	0.3 (5)
C14—N3—C11—C12	62.6 (6)	N2—C19—C14—C15	178.1 (4)
C9—N3—C14—C15	-177.9 (5)	C18—C19—C14—N3	-178.5 (4)
C11—N3—C14—C15	1.7 (8)	C18—C19—C14—C15	-0.7 (7)
C9—N3—C14—C19	-0.3 (5)	C19—C18—C17—C16	0.0 (8)
C11—N3—C14—C19	179.3 (4)	N2—C19—C18—C17	-178.3 (5)
C1—C2—C3—C4	-176.8 (5)	C14—C19—C18—C17	0.2 (7)

Symmetry codes: (i) $-x, -y+1, -z+2$.

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C8—H8A···Br1 ⁱⁱ	0.97	2.90	3.863 (5)	175
C10—H10C···Br1	0.96	2.93	3.788 (5)	150
C17—H17A···Br1 ⁱⁱⁱ	0.93	2.92	3.707 (5)	143
C4—H4a···Cg1 ^{iv}	0.93	3.36	3.746 (2)	107.5 (2)

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (iii) $-x, -y+1, -z+1$; (iv) $x, y, z-1$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

