

catena-Poly[[[(1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- $\kappa^2 O^3, O^4$)copper(II)]- μ -1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- $\kappa^3 N^{7'}:O^3, O^4$] tetrahydrate]

Shu-Ye Wang,^a Xue-Ming Song,^{a*} Li-Xiang Duan^a and Zhe An^b

^aThe First Affiliated Hospital, Harbin Medical University, Harbin, People's Republic of China, and ^bSchool of Chemistry and Life Science, Maoming University, Maoming 525000, People's Republic of China

Correspondence e-mail: song_xueming@sina.com


Received 16 August 2008; accepted 17 August 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.054; wR factor = 0.191; data-to-parameter ratio = 12.1.

In the title compound, $\{[Cu(C_{16}H_{17}FN_3O_3)_2] \cdot 4H_2O\}_n$, the Cu^{II} atom is bonded to two O, O' -bidentate 1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate (norf) monoanions and a symmetry-generated N -bonded norf anion, resulting in a distorted square-pyramidal coordination environment with the N atom occupying the apical site. The bridging norf anion results in one-dimensional chains propagating along $[010]$. A network of $O-H \cdots O$ and $N-H \cdots O$ hydrogen bonds helps to establish the crystal structure.

Related literature

For the iron, zinc and cobalt complexes of the norf anion, see: Chen *et al.* (2001); Qu *et al.* (2003); An *et al.* (2007). For background on the medicinal uses of Hnorf, see: Mizuki *et al.* (1996).



Experimental

Crystal data

$[Cu(C_{16}H_{17}FN_3O_3)_2] \cdot 4H_2O$

$M_r = 772.26$

Triclinic, $P\bar{1}$

$a = 10.023$ (2) Å

$b = 11.708$ (2) Å

$c = 16.219$ (3) Å

$\alpha = 97.22$ (3)°

$\beta = 107.05$ (3)°

$\gamma = 105.00$ (3)°

$V = 1715.2$ (6) Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 0.71$ mm⁻¹

$T = 296$ (2) K

$0.43 \times 0.28 \times 0.22$ mm

Data collection

Bruker APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2004)

$T_{\min} = 0.749$, $T_{\max} = 0.859$

13404 measured reflections

5960 independent reflections

4481 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.191$

$S = 1.00$

5960 reflections

492 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 1.90$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.72$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—O3	1.936 (3)	Cu1—O5	1.953 (3)
Cu1—O1	1.939 (3)	Cu1—N6 ⁱ	2.248 (3)
Cu1—O4	1.944 (3)		

Symmetry code: (i) $x, y - 1, z$.

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1 <i>W</i> —H1 <i>W</i> ···O4 <i>W</i>	0.83 (9)	2.08 (10)	2.735 (14)	136 (10)
O1 <i>W</i> —H2 <i>W</i> ···O3 <i>W</i> ⁱⁱ	0.83 (8)	1.78 (8)	2.580 (11)	163 (11)
O3 <i>W</i> —H5 <i>W</i> ···O1 <i>W</i> ⁱ	0.88 (6)	2.17 (11)	2.580 (11)	108 (9)
O3 <i>W</i> —H6 <i>W</i> ···O1	0.83 (8)	2.13 (8)	2.865 (8)	147 (9)
O4 <i>W</i> —H8 <i>W</i> ···O1 <i>W</i>	0.90 (9)	2.17 (11)	2.735 (14)	121 (10)
N6—H6 <i>A</i> ···O1 <i>W</i> ⁱⁱⁱ	1.05 (3)	2.12 (4)	3.105 (12)	155 (3)

Symmetry codes: (i) *x*, *y* − 1, *z*; (ii) *x*, *y* + 1, *z*; (iii) −*x* + 1, −*y* + 2, −*z*.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful for financial support from the Natural Science Foundation of Heilongjiang Province (D200672).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2782).

References

- An, Z., Xu, W. & Wang, R.-S. (2007). *Acta Cryst.* **E63**, m507–m508.
 Bruker (2004). *APEX2*, *SADABS* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chen, Z.-F., Xiong, R.-G., Zhang, J., Chen, X.-T., Xue, Z.-L. & You, X.-Z. (2001). *Inorg. Chem.* **40**, 4075–4077.
 Mizuki, Y., Fujiwara, I. & Yamaguchi, T. (1996). *J. Antimicrob. Chemother.* **37**, Suppl. A, 41–45.
 Qu, Z.-R., Zhao, H., Xing, L.-X., Wang, X.-S., Chen, Z.-F., Yu, Z., Xiong, R.-G. & You, X.-Z. (2003). *Eur. J. Inorg. Chem.* pp. 2920–2923.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m1201-m1202 [doi:10.1107/S1600536808026469]

***catena*-Poly[[[(1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- κ^2O^3,O^4)copper(II)]- μ -1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- $\kappa^3N^{7'}:O^3,O^4$] tetrahydrate]**

S.-Y. Wang, X.-M. Song, L.-X. Duan and Z. An

Comment

Norfloxacin (*H*-Norf, 1-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinoline carboxylic acid) is member of the class of quinolones that is used to treat infections (Mizuki *et al.* 1996). The zinc(II), iron(II) and cobalt(II) derivatives of norf have been reported (Chen. *et al.*, 2001; Qu *et al.* 2003; An *et al.*, 2007).

The title copper(II) derivative, (I), a one-dimensional coordination polymer in which the anion acts in a bridging mode, is reported here (Fig. 1).

The Cu(II) atom is coordinated (Table 1) by four oxygen atoms and one N atoms from three norfloxacin ligands (one monodentate-N and two O,*O*-bidentate) to form a one-dimensional coordination polymer (Fig. 2). A network of O—H \cdots O and N—H \cdots O hydrogen bonds (Table 2) helps to establish the packing.

Experimental

A mixture of Cu(CH₃COO)₂·H₂O (0.05 g, 0.25 mmol), Hnorf (0.16 g, 0.50 mmol), sodium hydroxide (0.02 g, 0.50 mmol) and water (12 ml) was stirred for 20 min in air. The mixture was then transferred to a 23 ml Teflon-lined hydrothermal bomb. The bomb was kept at 443 K for 96 h under autogenous pressure. After cooling, blue blocks of (I) were obtained from the reaction mixture.

Refinement

The N- and O-bonded H atoms were located in difference maps and their positions were freely refined with a fixed U_{iso} value. This has led to some very short H \cdots H intermolecular contacts and the positions of these H atoms should be regarded as less certain.

All the C-bonded H atoms were placed in calculated positions with C—H = 0.93 Å and refined as riding with $U_{iso}(H) = 1.2U_{eq}(\text{carrier})$.

Figures




Fig. 1. The asymmetric unit of (I), expanded to show the coordination of the Cu atom, drawn with 30% probability displacement ellipsoids for the non-hydrogen atoms (arbitrary spheres for the H atoms). Symmetry code: (i) $x, 1-y, z$.




Fig. 2. Part of the chain structure of (I).

catena-Poly[[[(1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- $\kappa^2\text{O}^3,\text{O}^4$)copper(II)]- μ -1-ethyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylato- $\kappa^3\text{N}^{7'}:\text{O}^3,\text{O}^4$] tetrahydrate]

Crystal data

$[\text{Cu}(\text{C}_{16}\text{H}_{17}\text{FN}_3\text{O}_3)_2] \cdot 4\text{H}_2\text{O}$

$M_r = 772.26$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.023$ (2) Å

$b = 11.708$ (2) Å

$c = 16.219$ (3) Å

$\alpha = 97.22$ (3)°

$\beta = 107.05$ (3)°

$\gamma = 105.00$ (3)°

$V = 1715.2$ (6) Å³

$Z = 2$

$F_{000} = 806$

$D_x = 1.495$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 5960 reflections

$\theta = 3.1\text{--}25.1^\circ$

$\mu = 0.71$ mm⁻¹

$T = 296$ (2) K

Block, blue

$0.43 \times 0.28 \times 0.22$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296$ (2) K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2004)

$T_{\min} = 0.749$, $T_{\max} = 0.859$

13404 measured reflections

5960 independent reflections

4481 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\text{max}} = 25.1^\circ$

$\theta_{\text{min}} = 3.1^\circ$

$h = -11 \rightarrow 11$

$k = -13 \rightarrow 13$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.191$

$S = 1.00$

5960 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: difmap and geom

H atoms treated by a mixture of
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.136P)^2 + 1.0468P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 1.90$ e Å⁻³

492 parameters

$$\Delta\rho_{\min} = -0.72 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.90729 (5)	0.04270 (4)	0.21869 (3)	0.0338 (2)
C1	0.6975 (4)	0.0019 (4)	0.3152 (3)	0.0439 (10)
C2	0.7856 (4)	-0.0629 (3)	0.3706 (3)	0.0373 (9)
C3	0.9119 (4)	-0.0878 (3)	0.3593 (3)	0.0342 (8)
C4	0.9788 (4)	-0.1560 (3)	0.4170 (3)	0.0362 (9)
C5	1.0984 (5)	-0.1904 (4)	0.4073 (3)	0.0436 (10)
H5	1.1368	-0.1669	0.3642	0.052*
C6	1.1580 (5)	-0.2569 (5)	0.4601 (3)	0.0524 (11)
C7	1.1120 (5)	-0.2918 (4)	0.5307 (3)	0.0442 (10)
C8	0.9952 (5)	-0.2557 (4)	0.5407 (3)	0.0408 (9)
H8	0.9617	-0.2748	0.5866	0.049*
C9	0.9262 (4)	-0.1917 (4)	0.4842 (3)	0.0377 (9)
C10	0.7405 (4)	-0.1000 (4)	0.4376 (3)	0.0424 (10)
H10	0.6590	-0.0819	0.4448	0.051*
C11	0.7415 (5)	-0.1957 (5)	0.5614 (3)	0.0532 (12)
H11A	0.6805	-0.1459	0.5693	0.064*
H11B	0.8203	-0.1801	0.6172	0.064*
C12	0.6511 (7)	-0.3266 (6)	0.5376 (5)	0.0801 (18)
H12A	0.5717	-0.3422	0.4830	0.120*
H12B	0.6122	-0.3452	0.5835	0.120*
H12C	0.7115	-0.3763	0.5309	0.120*
C13	1.3388 (5)	-0.3119 (5)	0.6269 (4)	0.0571 (12)
H13A	1.3626	-0.2425	0.6742	0.068*
H13B	1.3845	-0.2851	0.5847	0.068*
C14	1.3980 (6)	-0.4084 (5)	0.6642 (3)	0.0574 (12)
H14A	1.3819	-0.4739	0.6157	0.069*
H14B	1.5031	-0.3738	0.6944	0.069*
C15	1.1717 (6)	-0.5018 (5)	0.6819 (4)	0.0594 (13)
H15A	1.1267	-0.5291	0.7244	0.071*
H15B	1.1474	-0.5715	0.6348	0.071*

supplementary materials

C16	1.1055 (6)	-0.4098 (4)	0.6437 (3)	0.0505 (11)
H16A	1.0011	-0.4476	0.6124	0.061*
H16B	1.1180	-0.3442	0.6910	0.061*
C17	1.1846 (4)	0.1672 (4)	0.1965 (3)	0.0349 (9)
C18	1.1248 (4)	0.2682 (3)	0.1721 (2)	0.0320 (8)
C19	0.9742 (4)	0.2613 (3)	0.1532 (3)	0.0327 (8)
C20	0.9356 (4)	0.3645 (3)	0.1262 (3)	0.0319 (8)
C21	0.7879 (4)	0.3613 (3)	0.1007 (3)	0.0338 (9)
H21	0.7149	0.2912	0.0973	0.041*
C22	0.7527 (4)	0.4610 (3)	0.0811 (3)	0.0362 (9)
C23	0.8575 (4)	0.5732 (3)	0.0900 (3)	0.0333 (8)
C24	1.0026 (4)	0.5748 (3)	0.1113 (3)	0.0351 (9)
H24	1.0749	0.6456	0.1152	0.042*
C25	1.0414 (4)	0.4703 (3)	0.1269 (3)	0.0331 (8)
C26	1.2212 (4)	0.3712 (4)	0.1661 (3)	0.0383 (9)
H26	1.3180	0.3726	0.1767	0.046*
C27	1.3780 (7)	0.6550 (5)	0.2444 (5)	0.0809 (19)
H27A	1.3050	0.6809	0.2618	0.121*
H27B	1.4552	0.7247	0.2459	0.121*
H27C	1.4179	0.6088	0.2844	0.121*
C28	1.3088 (5)	0.5775 (4)	0.1518 (4)	0.0493 (11)
H28A	1.2717	0.6252	0.1112	0.059*
H28B	1.3826	0.5511	0.1344	0.059*
C29	0.7251 (5)	0.7081 (4)	0.1276 (3)	0.0423 (10)
H29A	0.7885	0.7424	0.1885	0.051*
H29B	0.6495	0.6362	0.1265	0.051*
C30	0.6551 (5)	0.7987 (4)	0.0915 (4)	0.0526 (12)
H30A	0.5873	0.7627	0.0317	0.063*
H30B	0.6001	0.8212	0.1275	0.063*
C31	0.8498 (5)	0.8720 (4)	0.0359 (3)	0.0469 (11)
H31A	0.9240	0.9430	0.0344	0.056*
H31B	0.7829	0.8360	-0.0240	0.056*
C32	0.9225 (5)	0.7818 (3)	0.0719 (3)	0.0387 (9)
H32A	0.9756	0.7589	0.0348	0.046*
H32B	0.9923	0.8188	0.1310	0.046*
F1	1.2644 (4)	-0.2977 (4)	0.4433 (2)	0.0794 (11)
F2	0.6077 (2)	0.4524 (2)	0.05141 (19)	0.0529 (7)
N1	0.8050 (4)	-0.1605 (3)	0.4935 (2)	0.0412 (8)
N2	1.1767 (4)	-0.3611 (4)	0.5827 (3)	0.0474 (9)
N3	1.3301 (5)	-0.4575 (4)	0.7252 (3)	0.0611 (11)
H3B	1.3751 (17)	-0.441 (4)	0.7842 (7)	0.073*
N4	1.1867 (3)	0.4699 (3)	0.1462 (2)	0.0366 (7)
N5	0.8124 (4)	0.6743 (3)	0.0745 (2)	0.0372 (8)
N6	0.7690 (4)	0.9081 (3)	0.0911 (2)	0.0392 (8)
H6A	0.723 (3)	0.9747 (16)	0.069 (3)	0.047*
O1	0.7412 (3)	0.0469 (3)	0.2558 (2)	0.0474 (7)
O2	0.5865 (4)	0.0120 (4)	0.3291 (3)	0.0752 (12)
O3	0.9694 (3)	-0.0539 (3)	0.3021 (2)	0.0424 (7)
O4	1.1027 (3)	0.0746 (2)	0.2105 (2)	0.0403 (7)

O5	0.8753 (3)	0.1742 (2)	0.1603 (2)	0.0419 (7)
O6	1.3139 (3)	0.1796 (3)	0.2032 (3)	0.0584 (9)
O1W	0.4376 (13)	0.9606 (8)	0.0102 (6)	0.182 (4)
O2W	0.2803 (6)	0.9249 (5)	0.2713 (5)	0.122 (2)
O3W	0.5350 (6)	0.1403 (7)	0.1444 (4)	0.109 (2)
O4W	0.2705 (8)	0.8303 (7)	0.0935 (7)	0.143 (3)
H1W	0.365 (9)	0.903 (8)	0.006 (7)	0.171*
H2W	0.473 (12)	1.008 (9)	0.059 (4)	0.171*
H3W	0.309 (13)	0.974 (7)	0.319 (4)	0.171*
H4W	0.240 (11)	0.855 (3)	0.274 (6)	0.171*
H5W	0.472 (9)	0.069 (4)	0.138 (8)	0.171*
H6W	0.615 (5)	0.143 (8)	0.180 (6)	0.171*
H7W	0.351 (6)	0.819 (10)	0.100 (9)	0.171*
H8W	0.265 (11)	0.888 (6)	0.062 (8)	0.171*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0306 (3)	0.0301 (3)	0.0431 (3)	0.0152 (2)	0.0094 (2)	0.0124 (2)
C1	0.030 (2)	0.045 (2)	0.061 (3)	0.0190 (18)	0.0139 (19)	0.016 (2)
C2	0.0310 (19)	0.0345 (19)	0.046 (2)	0.0103 (16)	0.0122 (17)	0.0070 (17)
C3	0.0313 (18)	0.0358 (19)	0.035 (2)	0.0104 (16)	0.0111 (16)	0.0074 (16)
C4	0.0335 (19)	0.0365 (19)	0.037 (2)	0.0114 (17)	0.0093 (16)	0.0087 (16)
C5	0.044 (2)	0.059 (3)	0.041 (2)	0.029 (2)	0.0185 (18)	0.022 (2)
C6	0.055 (3)	0.071 (3)	0.049 (3)	0.037 (2)	0.024 (2)	0.026 (2)
C7	0.045 (2)	0.054 (2)	0.037 (2)	0.021 (2)	0.0110 (18)	0.0162 (19)
C8	0.044 (2)	0.046 (2)	0.033 (2)	0.0144 (19)	0.0133 (17)	0.0109 (17)
C9	0.038 (2)	0.041 (2)	0.033 (2)	0.0136 (18)	0.0114 (16)	0.0066 (16)
C10	0.036 (2)	0.043 (2)	0.051 (3)	0.0161 (18)	0.0178 (19)	0.0069 (19)
C11	0.055 (3)	0.076 (3)	0.043 (3)	0.032 (3)	0.026 (2)	0.021 (2)
C12	0.066 (3)	0.095 (4)	0.082 (4)	0.009 (3)	0.036 (3)	0.034 (4)
C13	0.054 (3)	0.068 (3)	0.055 (3)	0.027 (2)	0.014 (2)	0.028 (2)
C14	0.061 (3)	0.074 (3)	0.044 (3)	0.038 (3)	0.012 (2)	0.020 (2)
C15	0.077 (3)	0.050 (3)	0.051 (3)	0.020 (3)	0.017 (2)	0.024 (2)
C16	0.061 (3)	0.054 (3)	0.043 (3)	0.021 (2)	0.020 (2)	0.020 (2)
C17	0.0322 (19)	0.040 (2)	0.038 (2)	0.0190 (17)	0.0122 (16)	0.0118 (16)
C18	0.0323 (18)	0.0334 (18)	0.032 (2)	0.0159 (16)	0.0085 (15)	0.0082 (15)
C19	0.0359 (19)	0.0289 (18)	0.035 (2)	0.0141 (16)	0.0102 (16)	0.0081 (15)
C20	0.0317 (18)	0.0259 (17)	0.035 (2)	0.0099 (15)	0.0067 (15)	0.0056 (15)
C21	0.0264 (17)	0.0283 (17)	0.041 (2)	0.0076 (15)	0.0035 (15)	0.0086 (15)
C22	0.0276 (18)	0.0297 (18)	0.047 (2)	0.0131 (16)	0.0036 (16)	0.0061 (16)
C23	0.0366 (19)	0.0266 (17)	0.034 (2)	0.0124 (16)	0.0062 (16)	0.0065 (15)
C24	0.036 (2)	0.0296 (18)	0.039 (2)	0.0080 (16)	0.0124 (16)	0.0115 (16)
C25	0.0346 (19)	0.0308 (18)	0.033 (2)	0.0119 (16)	0.0090 (15)	0.0082 (15)
C26	0.0303 (19)	0.044 (2)	0.045 (2)	0.0176 (17)	0.0122 (17)	0.0162 (18)
C27	0.062 (3)	0.056 (3)	0.092 (5)	-0.004 (3)	-0.002 (3)	0.017 (3)
C28	0.036 (2)	0.043 (2)	0.075 (3)	0.0081 (19)	0.027 (2)	0.024 (2)
C29	0.040 (2)	0.0281 (18)	0.063 (3)	0.0105 (17)	0.023 (2)	0.0081 (18)

supplementary materials

C30	0.040 (2)	0.029 (2)	0.078 (4)	0.0104 (19)	0.009 (2)	0.000 (2)
C31	0.069 (3)	0.0309 (19)	0.042 (2)	0.018 (2)	0.015 (2)	0.0146 (17)
C32	0.047 (2)	0.0285 (18)	0.047 (2)	0.0160 (17)	0.0207 (19)	0.0135 (17)
F1	0.090 (2)	0.135 (3)	0.077 (2)	0.090 (2)	0.0545 (19)	0.067 (2)
F2	0.0305 (12)	0.0325 (12)	0.0806 (19)	0.0100 (10)	-0.0026 (11)	0.0106 (11)
N1	0.0374 (17)	0.0491 (19)	0.040 (2)	0.0172 (16)	0.0145 (15)	0.0094 (16)
N2	0.051 (2)	0.056 (2)	0.042 (2)	0.0260 (18)	0.0144 (16)	0.0198 (17)
N3	0.079 (3)	0.065 (3)	0.042 (2)	0.034 (2)	0.009 (2)	0.022 (2)
N4	0.0315 (16)	0.0362 (17)	0.046 (2)	0.0113 (14)	0.0156 (14)	0.0152 (14)
N5	0.0391 (17)	0.0275 (15)	0.046 (2)	0.0146 (14)	0.0110 (15)	0.0127 (14)
N6	0.0406 (18)	0.0275 (15)	0.045 (2)	0.0171 (14)	0.0038 (15)	0.0051 (14)
O1	0.0407 (15)	0.0597 (18)	0.0563 (19)	0.0306 (14)	0.0186 (14)	0.0269 (15)
O2	0.051 (2)	0.101 (3)	0.114 (3)	0.048 (2)	0.048 (2)	0.062 (3)
O3	0.0378 (14)	0.0530 (17)	0.0492 (18)	0.0261 (13)	0.0171 (13)	0.0252 (14)
O4	0.0361 (14)	0.0392 (15)	0.0574 (19)	0.0239 (12)	0.0182 (13)	0.0205 (13)
O5	0.0308 (13)	0.0289 (13)	0.066 (2)	0.0114 (12)	0.0099 (13)	0.0200 (13)
O6	0.0397 (17)	0.059 (2)	0.094 (3)	0.0287 (15)	0.0296 (17)	0.0359 (18)
O1W	0.235 (12)	0.129 (7)	0.141 (7)	0.008 (6)	0.044 (8)	0.040 (5)
O2W	0.065 (3)	0.095 (3)	0.243 (8)	0.049 (3)	0.060 (4)	0.094 (4)
O3W	0.090 (3)	0.208 (6)	0.083 (3)	0.101 (4)	0.045 (3)	0.068 (4)
O4W	0.130 (5)	0.110 (5)	0.187 (8)	0.020 (4)	0.073 (6)	0.018 (5)

Geometric parameters (Å, °)

Cu1—O3	1.936 (3)	C18—C19	1.428 (5)
Cu1—O1	1.939 (3)	C19—O5	1.268 (5)
Cu1—O4	1.944 (3)	C19—C20	1.441 (5)
Cu1—O5	1.953 (3)	C20—C25	1.404 (5)
Cu1—N6 ⁱ	2.248 (3)	C20—C21	1.405 (5)
C1—O2	1.229 (6)	C21—C22	1.352 (5)
C1—O1	1.285 (6)	C21—H21	0.9300
C1—C2	1.489 (6)	C22—F2	1.362 (4)
C2—C10	1.371 (6)	C22—C23	1.414 (6)
C2—C3	1.426 (6)	C23—C24	1.388 (6)
C3—O3	1.281 (5)	C23—N5	1.398 (5)
C3—C4	1.435 (6)	C24—C25	1.406 (5)
C4—C5	1.403 (6)	C24—H24	0.9300
C4—C9	1.406 (6)	C25—N4	1.398 (5)
C5—C6	1.344 (6)	C26—N4	1.340 (5)
C5—H5	0.9300	C26—H26	0.9300
C6—F1	1.357 (6)	C27—C28	1.511 (8)
C6—C7	1.420 (7)	C27—H27A	0.9600
C7—C8	1.388 (6)	C27—H27B	0.9600
C7—N2	1.384 (6)	C27—H27C	0.9600
C8—C9	1.396 (6)	C28—N4	1.483 (5)
C8—H8	0.9300	C28—H28A	0.9700
C9—N1	1.398 (5)	C28—H28B	0.9700
C10—N1	1.341 (6)	C29—N5	1.481 (6)
C10—H10	0.9300	C29—C30	1.505 (6)

C11—N1	1.475 (6)	C29—H29A	0.9700
C11—C12	1.499 (8)	C29—H29B	0.9700
C11—H11A	0.9700	C30—N6	1.481 (5)
C11—H11B	0.9700	C30—H30A	0.9700
C12—H12A	0.9600	C30—H30B	0.9700
C12—H12B	0.9600	C31—N6	1.464 (6)
C12—H12C	0.9600	C31—C32	1.513 (6)
C13—N2	1.488 (6)	C31—H31A	0.9700
C13—C14	1.517 (7)	C31—H31B	0.9700
C13—H13A	0.9700	C32—N5	1.458 (5)
C13—H13B	0.9700	C32—H32A	0.9700
C14—N3	1.452 (7)	C32—H32B	0.9700
C14—H14A	0.9700	N3—H3B	0.900 (8)
C14—H14B	0.9700	N6—Cu1 ⁱⁱ	2.248 (3)
C15—N3	1.454 (7)	N6—H6A	1.05 (2)
C15—C16	1.511 (7)	O1W—O1W ⁱⁱⁱ	1.500 (18)
C15—H15A	0.9700	O1W—H1W	0.83 (9)
C15—H15B	0.9700	O1W—H2W	0.83 (8)
C16—N2	1.469 (6)	O2W—H3W	0.83 (7)
C16—H16A	0.9700	O2W—H4W	0.83 (4)
C16—H16B	0.9700	O3W—H5W	0.88 (3)
C17—O6	1.236 (5)	O3W—H6W	0.83 (8)
C17—O4	1.270 (5)	O4W—H7W	0.83 (8)
C17—C18	1.503 (5)	O4W—H8W	0.90 (9)
C18—C26	1.369 (6)		
O3—Cu1—O1	92.22 (13)	C21—C20—C19	119.3 (3)
O3—Cu1—O4	85.49 (12)	C22—C21—C20	119.4 (3)
O1—Cu1—O4	164.06 (13)	C22—C21—H21	120.3
O3—Cu1—O5	165.39 (13)	C20—C21—H21	120.3
O1—Cu1—O5	87.15 (13)	C21—C22—F2	117.8 (3)
O4—Cu1—O5	91.12 (12)	C21—C22—C23	123.8 (3)
O3—Cu1—N6 ⁱ	105.05 (13)	F2—C22—C23	118.4 (3)
O1—Cu1—N6 ⁱ	94.09 (14)	C24—C23—N5	123.3 (3)
O4—Cu1—N6 ⁱ	101.75 (14)	C24—C23—C22	116.5 (3)
O5—Cu1—N6 ⁱ	89.55 (13)	N5—C23—C22	120.1 (3)
O2—C1—O1	122.4 (4)	C23—C24—C25	120.6 (4)
O2—C1—C2	118.5 (4)	C23—C24—H24	119.7
O1—C1—C2	119.1 (4)	C25—C24—H24	119.7
C10—C2—C3	118.6 (4)	N4—C25—C20	118.1 (3)
C10—C2—C1	116.4 (4)	N4—C25—C24	121.2 (3)
C3—C2—C1	125.1 (4)	C20—C25—C24	120.7 (3)
O3—C3—C2	125.0 (4)	N4—C26—C18	125.0 (4)
O3—C3—C4	118.1 (4)	N4—C26—H26	117.5
C2—C3—C4	117.0 (4)	C18—C26—H26	117.5
C5—C4—C9	117.8 (4)	C28—C27—H27A	109.5
C5—C4—C3	120.8 (4)	C28—C27—H27B	109.5
C9—C4—C3	121.4 (4)	H27A—C27—H27B	109.5

supplementary materials

C6—C5—C4	120.4 (4)	C28—C27—H27C	109.5
C6—C5—H5	119.8	H27A—C27—H27C	109.5
C4—C5—H5	119.8	H27B—C27—H27C	109.5
C5—C6—F1	118.4 (4)	N4—C28—C27	111.2 (4)
C5—C6—C7	123.9 (4)	N4—C28—H28A	109.4
F1—C6—C7	117.6 (4)	C27—C28—H28A	109.4
C8—C7—N2	123.2 (4)	N4—C28—H28B	109.4
C8—C7—C6	115.3 (4)	C27—C28—H28B	109.4
N2—C7—C6	121.4 (4)	H28A—C28—H28B	108.0
C7—C8—C9	122.2 (4)	N5—C29—C30	110.5 (4)
C7—C8—H8	118.9	N5—C29—H29A	109.5
C9—C8—H8	118.9	C30—C29—H29A	109.6
C8—C9—N1	121.1 (4)	N5—C29—H29B	109.5
C8—C9—C4	120.3 (4)	C30—C29—H29B	109.5
N1—C9—C4	118.5 (4)	H29A—C29—H29B	108.1
N1—C10—C2	124.8 (4)	N6—C30—C29	110.2 (3)
N1—C10—H10	117.6	N6—C30—H30A	109.6
C2—C10—H10	117.6	C29—C30—H30A	109.6
N1—C11—C12	112.5 (4)	N6—C30—H30B	109.6
N1—C11—H11A	109.1	C29—C30—H30B	109.6
C12—C11—H11A	109.1	H30A—C30—H30B	108.1
N1—C11—H11B	109.1	N6—C31—C32	110.6 (4)
C12—C11—H11B	109.1	N6—C31—H31A	109.5
H11A—C11—H11B	107.8	C32—C31—H31A	109.5
C11—C12—H12A	109.5	N6—C31—H31B	109.5
C11—C12—H12B	109.5	C32—C31—H31B	109.5
H12A—C12—H12B	109.5	H31A—C31—H31B	108.1
C11—C12—H12C	109.5	N5—C32—C31	110.2 (3)
H12A—C12—H12C	109.5	N5—C32—H32A	109.6
H12B—C12—H12C	109.5	C31—C32—H32A	109.6
N2—C13—C14	110.1 (4)	N5—C32—H32B	109.6
N2—C13—H13A	109.6	C31—C32—H32B	109.6
C14—C13—H13A	109.6	H32A—C32—H32B	108.1
N2—C13—H13B	109.6	C10—N1—C9	119.6 (4)
C14—C13—H13B	109.6	C10—N1—C11	118.2 (4)
H13A—C13—H13B	108.1	C9—N1—C11	122.1 (4)
N3—C14—C13	113.2 (4)	C7—N2—C16	117.2 (4)
N3—C14—H14A	109.0	C7—N2—C13	116.2 (4)
C13—C14—H14A	109.0	C16—N2—C13	110.8 (4)
N3—C14—H14B	108.9	C14—N3—C15	109.1 (4)
C13—C14—H14B	108.9	C14—N3—H3B	124.2 (12)
H14A—C14—H14B	107.8	C15—N3—H3B	123.9 (12)
N3—C15—C16	114.5 (4)	C26—N4—C25	119.5 (3)
N3—C15—H15A	108.6	C26—N4—C28	117.4 (3)
C16—C15—H15A	108.6	C25—N4—C28	123.0 (3)
N3—C15—H15B	108.6	C23—N5—C32	117.2 (3)
C16—C15—H15B	108.6	C23—N5—C29	115.6 (3)
H15A—C15—H15B	107.6	C32—N5—C29	110.5 (3)
N2—C16—C15	110.3 (4)	C31—N6—C30	108.5 (3)

N2—C16—H16A	109.6	C31—N6—Cu1 ⁱⁱ	115.4 (3)
C15—C16—H16A	109.6	C30—N6—Cu1 ⁱⁱ	119.5 (3)
N2—C16—H16B	109.6	C31—N6—H6A	111.7 (13)
C15—C16—H16B	109.6	C30—N6—H6A	111.6 (13)
H16A—C16—H16B	108.1	Cu1 ⁱⁱ —N6—H6A	89 (2)
O6—C17—O4	122.8 (4)	C1—O1—Cu1	131.2 (3)
O6—C17—C18	117.7 (4)	C3—O3—Cu1	127.2 (3)
O4—C17—C18	119.6 (3)	C17—O4—Cu1	130.5 (2)
C26—C18—C19	118.9 (3)	C19—O5—Cu1	126.1 (2)
C26—C18—C17	117.4 (3)	O1W ⁱⁱⁱ —O1W—H1W	161 (7)
C19—C18—C17	123.7 (3)	O1W ⁱⁱⁱ —O1W—H2W	85 (7)
O5—C19—C18	125.4 (3)	H1W—O1W—H2W	113 (10)
O5—C19—C20	118.8 (3)	H3W—O2W—H4W	113 (8)
C18—C19—C20	115.8 (3)	H5W—O3W—H6W	108 (9)
C25—C20—C21	118.4 (3)	H7W—O4W—H8W	108 (11)
C25—C20—C19	122.2 (3)		

Symmetry codes: (i) $x, y-1, z$; (ii) $x, y+1, z$; (iii) $-x+1, -y+2, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1W \cdots O4W	0.83 (9)	2.08 (10)	2.735 (14)	136 (10)
O1W—H2W \cdots O3W ⁱⁱ	0.83 (8)	1.78 (8)	2.580 (11)	163 (11)
O3W—H5W \cdots O1W ⁱ	0.88 (6)	2.17 (11)	2.580 (11)	108 (9)
O3W—H6W \cdots O1	0.83 (8)	2.13 (8)	2.865 (8)	147 (9)
O4W—H8W \cdots O1W	0.90 (9)	2.17 (11)	2.735 (14)	121 (10)
N6—H6A \cdots O1W ⁱⁱⁱ	1.05 (3)	2.12 (4)	3.105 (12)	155 (3)

Symmetry codes: (ii) $x, y+1, z$; (i) $x, y-1, z$; (iii) $-x+1, -y+2, -z$.

Fig. 1




Fig. 2

