

## ***trans*-4-[(Phenylsulfonyloxy)methyl]cyclohexanecarboxylic acid**

**Yu-Feng Liang, Qing-Rong Qi and Hu Zheng\***

Department of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu 610041, People's Republic of China  
Correspondence e-mail: wcums416@yahoo.com.cn


Received 15 July 2008; accepted 31 July 2008

Key indicators: single-crystal X-ray study;  $T = 292$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005 \text{ \AA}$ ;  $R$  factor = 0.055;  $wR$  factor = 0.143; data-to-parameter ratio = 14.7.

The title compound,  $\text{C}_{14}\text{H}_{18}\text{O}_5\text{S}$ , is an important intermediate for the synthesis of poly(amidoamine) dendrimers. The cyclohexane ring adopts a chair conformation with its two substituents in equatorial positions. In the crystal structure, molecules form centrosymmetric dimers via  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For related literature, see: Ahmed *et al.* (2001); Grabchev *et al.* (2003); Wang *et al.* (2004).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_{18}\text{O}_5\text{S}$   
 $M_r = 298.34$   
Monoclinic,  $P2_1/c$   
 $a = 17.097 (5) \text{ \AA}$

$b = 5.960 (3) \text{ \AA}$   
 $c = 14.919 (4) \text{ \AA}$   
 $\beta = 107.09 (3)^\circ$   
 $V = 1453.2 (10) \text{ \AA}^3$

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.24 \text{ mm}^{-1}$

$T = 292 (2) \text{ K}$   
 $0.32 \times 0.32 \times 0.13 \text{ mm}$

#### Data collection

Enraf–Nonius CAD-4 diffractometer  
Absorption correction: none  
3702 measured reflections  
2691 independent reflections

1323 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.008$   
3 standard reflections every 250 reflections  
intensity decay: 0.9%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$   
 $wR(F^2) = 0.142$   
 $S = 1.00$   
2691 reflections

183 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4 $\cdots$ O5 <sup>i</sup>	0.82	1.86	2.677 (3)	174

Symmetry code: (i)  $-x + 1, -y, -z + 2$ .

Data collection: *DIFRAC* (Gabe & White, 1993); cell refinement: *DIFRAC*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2749).

### References

- Ahmed, S. M., Budd, P. M., McKeown, N. B., Evans, K. P., Beaumont, G. L., Donaldson, C. & Brennan, C. M. (2001). *Polymer*, **42**, 889–896.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Gabe, E. J. & White, P. S. (1993). *DIFRAC*. American Crystallographic Association Meeting, Pittsburgh, Abstract PA 104.
- Grabchev, I., Chovelon, J. M., Bojinov, V. & Ivanova, G. (2003). *Tetrahedron*, **59**, 9591–9598.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, B.-B., Zhang, X., Jia, X.-R., Luo, Y.-F., Sun, Z., Yang, L., Ji, Y. & Wei, Y. (2004). *Polymer*, **45**, 8395–8402.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o1703 [doi:10.1107/S1600536808024513]

### ***trans*-4-[(Phenylsulfonyloxy)methyl]cyclohexanecarboxylic acid**

**Y.-F. Liang, Q.-R. Qi and H. Zheng**

#### **Comment**

PAMAM (poly(amidoamine)) dendrimers have attracted much interest for their symmetry, high degree of branching and high density of terminal functional groups, which can participate in different reactions. The modification of periphery of PAMAM dendrimer which aimed to change the physical or chemical properties of PAMAM dendrimers, have been reported recently (Grabchev *et al.*, 2003; Ahmed *et al.*, 2001; Wang *et al.*, 2004). To improve the lipophilicity of PAMAM dendrimers and provide a new type of linker with special stereostructure, a series of cyclohexane derivatives were synthesized.

The crystal structure shows that molecules are mainly linked by O—H $\cdots$ O hydrogen bonds and the cyclohexane ring of the title compound exists in the chair conformation.

#### **Experimental**

*trans*-4-(methoxycarbonyl)cyclohexanemethanol (10 mmol), triethylamine (10 mmol) and a small amount of trimethylamine hydrochloride were suspended in dichloromethane (20 mL), benzenesulfonyl chloride (11 mmol) was dropped with vigorous stirring at room temperature, after 1 h the reaction was quenched by addition of water. The organic layer separated was evaporated to give an oil and the oil was hydrolyzed in methanol and aqueous NaOH (11 mmol) solution for 5 h at 323 K. Then the title compound was obtained by acidification with hydrochloride and recrystallized from acetone. Colorless crystals suitable for X-ray analysis were obtained by slow evaporation in cyclohexane and acetone at room temperature.

#### **Refinement**

H atoms were positioned geometrically (C—H = 0.93–0.98 Å, O—H = 0.82 Å) and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{O})$ .

#### **Figures**




Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level.

# supplementary materials

---




Fig. 2. A packing diagram of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

## *trans*-4-[(Phenylsulfonyloxy)methyl]cyclohexanecarboxylic acid

### Crystal data

C <sub>14</sub> H <sub>18</sub> O <sub>5</sub> S	$F_{000} = 632$
$M_r = 298.34$	$D_x = 1.364 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 17.097 (5) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 5.960 (3) \text{ \AA}$	Cell parameters from 24 reflections
$c = 14.919 (4) \text{ \AA}$	$\theta = 4.4\text{--}8.7^\circ$
$\beta = 107.09 (3)^\circ$	$\mu = 0.24 \text{ mm}^{-1}$
$V = 1453.2 (10) \text{ \AA}^3$	$T = 292 (2) \text{ K}$
$Z = 4$	Block, colourless
	$0.32 \times 0.32 \times 0.13 \text{ mm}$

### Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.008$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.5^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 1.3^\circ$
$T = 292(2) \text{ K}$	$h = -2\text{--}20$
$\omega/2\theta$ scans	$k = -7\text{--}0$
Absorption correction: none	$l = -18\text{--}17$
3702 measured reflections	3 standard reflections
2691 independent reflections	every 250 reflections
1323 reflections with $I > 2\sigma(I)$	intensity decay: 0.9%

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.054$	$w = 1/[\sigma^2(F_o^2) + (0.0729P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.143$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$

2691 reflections	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
183 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0076 (16)
Secondary atom site location: difference Fourier map	

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.15478 (7)	0.60525 (16)	0.42320 (6)	0.0631 (4)
O1	0.15724 (19)	0.8419 (4)	0.42806 (17)	0.0942 (11)
O2	0.10478 (16)	0.4829 (4)	0.46844 (17)	0.0782 (8)
O3	0.24614 (15)	0.5294 (4)	0.46482 (16)	0.0705 (8)
O4	0.4357 (2)	-0.1520 (5)	0.8999 (2)	0.1010 (11)
H4	0.4645	-0.1623	0.9545	0.151*
O5	0.46270 (19)	0.2062 (5)	0.92678 (18)	0.0973 (11)
C1	0.0928 (2)	0.3172 (6)	0.2785 (2)	0.0558 (10)
H1	0.0855	0.2167	0.3231	0.067*
C2	0.0687 (2)	0.2625 (7)	0.1846 (3)	0.0694 (12)
H2	0.0457	0.1227	0.1654	0.083*
C3	0.0785 (3)	0.4130 (9)	0.1199 (3)	0.0786 (14)
H3	0.0615	0.3752	0.0567	0.094*
C4	0.1127 (3)	0.6184 (9)	0.1467 (3)	0.0807 (13)
H4A	0.1184	0.7202	0.1017	0.097*
C5	0.1386 (2)	0.6739 (6)	0.2393 (2)	0.0636 (11)
H5	0.1632	0.8122	0.2579	0.076*
C6	0.12814 (18)	0.5243 (5)	0.3051 (2)	0.0434 (9)
C7	0.2647 (2)	0.2958 (6)	0.4899 (2)	0.0634 (11)
H7A	0.2877	0.2257	0.4447	0.076*
H7B	0.2147	0.2168	0.4885	0.076*
C8	0.3248 (2)	0.2787 (6)	0.5868 (2)	0.0543 (10)
H8	0.3740	0.3632	0.5870	0.065*
C9	0.2910 (2)	0.3764 (6)	0.6619 (2)	0.0560 (10)
H9A	0.2390	0.3056	0.6579	0.067*
H9B	0.2812	0.5356	0.6504	0.067*
C10	0.3488 (2)	0.3433 (6)	0.7602 (2)	0.0595 (11)

## supplementary materials

---

H10A	0.3985	0.4287	0.7669	0.071*
H10B	0.3232	0.3991	0.8058	0.071*
C11	0.3702 (2)	0.0971 (6)	0.7794 (2)	0.0540 (10)
H11	0.3193	0.0165	0.7753	0.065*
C12	0.4053 (2)	-0.0003 (6)	0.7056 (2)	0.0685 (12)
H12A	0.4147	-0.1597	0.7169	0.082*
H12B	0.4577	0.0699	0.7109	0.082*
C13	0.3484 (2)	0.0350 (6)	0.6063 (2)	0.0688 (12)
H13A	0.3754	-0.0171	0.5614	0.083*
H13B	0.2993	-0.0540	0.5981	0.083*
C14	0.4275 (2)	0.0573 (7)	0.8757 (3)	0.0637 (11)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0904 (8)	0.0477 (6)	0.0373 (5)	0.0035 (6)	-0.0028 (5)	-0.0010 (5)
O1	0.152 (3)	0.0478 (16)	0.0584 (17)	0.0040 (17)	-0.0069 (17)	-0.0068 (14)
O2	0.103 (2)	0.0828 (19)	0.0552 (16)	0.0009 (16)	0.0327 (15)	0.0129 (15)
O3	0.0844 (18)	0.0586 (15)	0.0488 (15)	-0.0147 (13)	-0.0111 (13)	0.0161 (12)
O4	0.129 (3)	0.0696 (19)	0.0663 (19)	-0.0009 (17)	-0.0307 (17)	0.0240 (15)
O5	0.125 (3)	0.0689 (18)	0.0601 (18)	-0.0064 (17)	-0.0309 (17)	0.0143 (15)
C1	0.060 (2)	0.053 (2)	0.049 (2)	-0.0035 (18)	0.0074 (18)	0.0019 (18)
C2	0.062 (2)	0.065 (3)	0.065 (3)	0.005 (2)	-0.006 (2)	-0.022 (2)
C3	0.086 (3)	0.104 (4)	0.034 (2)	0.034 (3)	-0.001 (2)	-0.006 (3)
C4	0.105 (3)	0.091 (3)	0.046 (2)	0.020 (3)	0.023 (2)	0.023 (3)
C5	0.077 (3)	0.058 (2)	0.051 (2)	0.004 (2)	0.012 (2)	0.014 (2)
C6	0.049 (2)	0.0432 (19)	0.0327 (18)	0.0033 (16)	0.0031 (15)	0.0025 (15)
C7	0.078 (3)	0.055 (2)	0.046 (2)	0.006 (2)	0.0014 (19)	0.0017 (19)
C8	0.057 (2)	0.053 (2)	0.044 (2)	0.0011 (17)	0.0010 (18)	0.0034 (18)
C9	0.060 (2)	0.054 (2)	0.046 (2)	0.0115 (18)	0.0023 (17)	0.0113 (19)
C10	0.068 (2)	0.064 (2)	0.0378 (19)	0.010 (2)	0.0006 (17)	0.0036 (18)
C11	0.051 (2)	0.057 (2)	0.045 (2)	0.0009 (18)	0.0007 (17)	0.0120 (19)
C12	0.072 (2)	0.063 (2)	0.058 (2)	0.017 (2)	0.000 (2)	0.009 (2)
C13	0.078 (3)	0.067 (3)	0.048 (2)	0.020 (2)	-0.003 (2)	-0.0003 (18)
C14	0.060 (2)	0.066 (3)	0.056 (2)	0.004 (2)	0.0020 (19)	0.020 (2)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

S1—O1	1.412 (3)	C7—H7A	0.9700
S1—O2	1.434 (3)	C7—H7B	0.9700
S1—O3	1.568 (3)	C8—C13	1.513 (5)
S1—C6	1.754 (3)	C8—C9	1.521 (5)
O3—C7	1.453 (4)	C8—H8	0.9800
O4—C14	1.295 (4)	C9—C10	1.521 (4)
O4—H4	0.8200	C9—H9A	0.9700
O5—C14	1.209 (4)	C9—H9B	0.9700
C1—C2	1.378 (5)	C10—C11	1.519 (4)
C1—C6	1.380 (4)	C10—H10A	0.9700
C1—H1	0.9300	C10—H10B	0.9700

C2—C3	1.364 (6)	C11—C14	1.501 (4)
C2—H2	0.9300	C11—C12	1.516 (5)
C3—C4	1.365 (6)	C11—H11	0.9800
C3—H3	0.9300	C12—C13	1.529 (4)
C4—C5	1.362 (5)	C12—H12A	0.9700
C4—H4A	0.9300	C12—H12B	0.9700
C5—C6	1.376 (4)	C13—H13A	0.9700
C5—H5	0.9300	C13—H13B	0.9700
C7—C8	1.510 (4)		
O1—S1—O2	119.8 (2)	C13—C8—H8	108.2
O1—S1—O3	104.85 (16)	C9—C8—H8	108.2
O2—S1—O3	109.22 (15)	C8—C9—C10	112.4 (3)
O1—S1—C6	108.74 (16)	C8—C9—H9A	109.1
O2—S1—C6	108.57 (16)	C10—C9—H9A	109.1
O3—S1—C6	104.66 (16)	C8—C9—H9B	109.1
C7—O3—S1	119.6 (2)	C10—C9—H9B	109.1
C14—O4—H4	109.5	H9A—C9—H9B	107.9
C2—C1—C6	118.6 (4)	C11—C10—C9	111.1 (3)
C2—C1—H1	120.7	C11—C10—H10A	109.4
C6—C1—H1	120.7	C9—C10—H10A	109.4
C3—C2—C1	120.1 (4)	C11—C10—H10B	109.4
C3—C2—H2	119.9	C9—C10—H10B	109.4
C1—C2—H2	119.9	H10A—C10—H10B	108.0
C2—C3—C4	120.9 (4)	C14—C11—C12	110.4 (3)
C2—C3—H3	119.5	C14—C11—C10	112.7 (3)
C4—C3—H3	119.5	C12—C11—C10	110.9 (3)
C5—C4—C3	119.9 (4)	C14—C11—H11	107.5
C5—C4—H4A	120.1	C12—C11—H11	107.5
C3—C4—H4A	120.1	C10—C11—H11	107.5
C4—C5—C6	119.6 (4)	C11—C12—C13	112.1 (3)
C4—C5—H5	120.2	C11—C12—H12A	109.2
C6—C5—H5	120.2	C13—C12—H12A	109.2
C5—C6—C1	120.8 (3)	C11—C12—H12B	109.2
C5—C6—S1	119.3 (3)	C13—C12—H12B	109.2
C1—C6—S1	119.8 (3)	H12A—C12—H12B	107.9
O3—C7—C8	110.3 (3)	C8—C13—C12	112.1 (3)
O3—C7—H7A	109.6	C8—C13—H13A	109.2
C8—C7—H7A	109.6	C12—C13—H13A	109.2
O3—C7—H7B	109.6	C8—C13—H13B	109.2
C8—C7—H7B	109.6	C12—C13—H13B	109.2
H7A—C7—H7B	108.1	H13A—C13—H13B	107.9
C7—C8—C13	108.5 (3)	O5—C14—O4	122.6 (3)
C7—C8—C9	112.4 (3)	O5—C14—C11	123.5 (3)
C13—C8—C9	111.3 (3)	O4—C14—C11	113.9 (3)
C7—C8—H8	108.2		
O1—S1—O3—C7	166.4 (3)	S1—O3—C7—C8	-132.0 (3)
O2—S1—O3—C7	36.9 (3)	O3—C7—C8—C13	-174.7 (3)
C6—S1—O3—C7	-79.2 (3)	O3—C7—C8—C9	61.8 (4)

## supplementary materials

---


C6—C1—C2—C3	−1.2 (5)	C7—C8—C9—C10	175.6 (3)
C1—C2—C3—C4	0.6 (6)	C13—C8—C9—C10	53.7 (4)
C2—C3—C4—C5	0.8 (6)	C8—C9—C10—C11	−55.3 (4)
C3—C4—C5—C6	−1.6 (6)	C9—C10—C11—C14	179.8 (3)
C4—C5—C6—C1	1.0 (5)	C9—C10—C11—C12	55.4 (4)
C4—C5—C6—S1	−175.4 (3)	C14—C11—C12—C13	179.5 (3)
C2—C1—C6—C5	0.4 (5)	C10—C11—C12—C13	−54.7 (4)
C2—C1—C6—S1	176.7 (3)	C7—C8—C13—C12	−176.5 (3)
O1—S1—C6—C5	20.1 (3)	C9—C8—C13—C12	−52.4 (4)
O2—S1—C6—C5	151.9 (3)	C11—C12—C13—C8	53.6 (5)
O3—S1—C6—C5	−91.5 (3)	C12—C11—C14—O5	113.9 (5)
O1—S1—C6—C1	−156.3 (3)	C10—C11—C14—O5	−10.8 (6)
O2—S1—C6—C1	−24.4 (3)	C12—C11—C14—O4	−66.6 (5)
O3—S1—C6—C1	92.1 (3)	C10—C11—C14—O4	168.7 (4)

### Hydrogen-bond geometry ( $\text{\AA}$ , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O4—H4 <sup>ii</sup> —O5 <sup>i</sup>	0.82	1.86	2.677 (3)	174

Symmetry codes: (i)  $-x+1, -y, -z+2$ .

Fig. 1



## supplementary materials

---

Fig. 2

