

**S-Benzylisothiouronium nitrate****P. Hemalatha<sup>a</sup> and V. Veeravazhuthi<sup>b\*</sup>**

<sup>a</sup>Department of Physics, PSG College of Technology, Coimbatore 641 004, TamilNadu, India, and <sup>b</sup>Department of Physics, PSG College of Arts and Science, Coimbatore 641 014, TamilNadu, India  
Correspondence e-mail: vv\_vazhuthi@rediffmail.com


Received 25 July 2008; accepted 18 August 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004 \text{ \AA}$ ;  $R$  factor = 0.074;  $wR$  factor = 0.229; data-to-parameter ratio = 18.3.

In the crystal structure of the title compound,  $\text{C}_8\text{H}_{11}\text{N}_2\text{S}^+\cdot\text{NO}_3^-$ , cations and anions are linked by intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, forming one-dimensional chains along [110].

**Related literature**

For related literature, see: Barker & Powell (1998); Boyd (1989); Hemalatha *et al.* (2006); Zaccaro *et al.* (1999); Zyss *et al.* (1984).

**Experimental***Crystal data*

$\text{C}_8\text{H}_{11}\text{N}_2\text{S}^+\cdot\text{NO}_3^-$   
 $M_r = 229.26$   
Monoclinic,  $P2_1/c$

$a = 5.8569 (4) \text{ \AA}$   
 $b = 7.5931 (5) \text{ \AA}$   
 $c = 23.9488 (16) \text{ \AA}$

$\beta = 93.304 (1)^\circ$   
 $V = 1063.28 (12) \text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.30 \text{ mm}^{-1}$   
 $T = 293 (2) \text{ K}$   
 $0.25 \times 0.21 \times 0.20 \text{ mm}$

*Data collection*

Bruker SMART APEX CCD area-detector diffractometer  
Absorption correction: none  
11620 measured reflections

2492 independent reflections  
2282 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.074$   
 $wR(F^2) = 0.229$   
 $S = 1.00$   
2492 reflections

136 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 1.08 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.79 \text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A $\cdots$ O2	0.86	1.98	2.803 (4)	160
N1—H1B $\cdots$ O3 <sup>i</sup>	0.86	2.23	3.009 (4)	151
N2—H2A $\cdots$ O1	0.86	2.21	3.040 (5)	164
N2—H2A $\cdots$ O2	0.86	2.56	3.240 (4)	136
N2—H2B $\cdots$ O1 <sup>ii</sup>	0.86	2.12	2.913 (4)	152

Symmetry codes: (i)  $-x + 2, -y, -z$ ; (ii)  $-x + 1, -y + 1, -z$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* and *PARST* (Nardelli, 1995).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2669).

**References**

- Barker, J. & Powell, H. R. (1998). *Acta Cryst. C* **54**, 2019–2021.  
Boyd, G. T. (1989). *J. Opt. Soc. Am. B* **6**, 685–688.  
Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Hemalatha, P., Veeravazhuthi, V., Mallika, J., Narayananadass, S. K. & Mangalaraj, D. (2006). *Cry. Res. Tec.* **41**, 775–779.  
Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.  
Zaccaro, J., Lorutet, F. & Ibanez, A. (1999). *J. Mater. Chem.* **9**, 1091–1094.  
Zyss, J., Nicoud, J. F. & Koquillay, M. (1984). *J. Chem. Phys.* **81**, 4160–4162.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o1805 [doi:10.1107/S1600536808026512]

### S-Benzylisothiouronium nitrate

P. Hemalatha and V. Veeravazhuthi

#### Comment

Organic molecular materials have many potential applications in integrated optics, and one of the most attractive applications is diode laser frequency doublers (Boyd, 1989). In the last two decades, extensive research has shown that organic crystals can exhibit nonlinear optical [NLO] efficiencies higher than those of inorganic materials (Zyss *et al.*, 1984 & Zaccaro *et al.*, 1999). Organic nonlinear optical materials are often formed by weak Vander Waals and hydrogen bonds and hence posses high degree of delocalization. Organic materials are molecular materials that offer unique opportunities for fundamental research as well as for technological applications. The title compound (I) is potentially in the above category of materials, therefore we have undertaken its crystal structure determination.

The title molecule is shown in Fig. 1. The C—N, S—C bond lengths and C—S—C and N—C—N bond angles are comparable with the similar structure reported earlier (Barker & Powell, 1998). The bond angles for O1—N3—O3 is 128.7 (4); O1—N3—O2 is 116.7 (4); O3—N3—O2 is 114.6 (3), indicating slight deviations in the bond angle from the expected 120° in terms of the  $sp^2$  hybridization. In the title crystal structure,  $C_8H_{11}N_2S$ ,  $NO_3^-$ , cations and anions are linked by intermolecular N—H···O hydrogen bonds to form one-dimensional chains along [110] (Fig. 2).

#### Experimental

*S*-benzylisothiouronium chloride (SBTC) was synthesized as reported earlier (Hemalatha *et al.*, 2006). The solutions of SBTC (5 g m) and potassium nitrate (5 g m) were prepared in water separately. These solutions were mixed together, and then stirred for 1 hr at room temperature. The precipitate was filtered off and washed with triple distilled water and the product was recrystallized from 0.2 M nitric acid. Single crystals were grown by slow evaporation of a solution of the title compound in water.

#### Refinement

All H-atoms were refined using a riding-model with  $d(C—H) = 0.93 \text{ \AA}$ ,  $U_{iso} = 1.2U_{eq}(C)$  for aromatic,  $0.97 \text{ \AA}$ ,  $U_{iso} = 1.2U_{eq}(C)$  for  $CH_2$  and  $0.86\text{\AA}$  for N—H with  $U_{iso}(H) = 1.2U_{eq}(N)$ .

#### Figures




Fig. 1. The molecular structure of title compound, showing 30% probability displacement ellipsoids.

# supplementary materials

---




Fig. 2. Part of the crystal structure of (I) showing hydrogen bonds as dashed lines.

## S-Benzylisothiouronium nitrate

### Crystal data

C <sub>8</sub> H <sub>11</sub> N <sub>2</sub> S <sup>+</sup> ·NO <sub>3</sub> <sup>-</sup>	F <sub>000</sub> = 480
M <sub>r</sub> = 229.26	D <sub>x</sub> = 1.432 Mg m <sup>-3</sup>
Monoclinic, P2 <sub>1</sub> /c	Mo K $\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda$ = 0.71073 Å
a = 5.8569 (4) Å	Cell parameters from 1296 reflections
b = 7.5931 (5) Å	$\theta$ = 1.7–28.0°
c = 23.9488 (16) Å	$\mu$ = 0.30 mm <sup>-1</sup>
$\beta$ = 93.304 (1)°	T = 293 (2) K
V = 1063.28 (12) Å <sup>3</sup>	Needle, colorless
Z = 4	0.25 × 0.21 × 0.20 mm

### Data collection

Bruker SMART APEXCCD area-detector diffractometer	2282 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}}$ = 0.021
Monochromator: graphite	$\theta_{\text{max}}$ = 28.0°
T = 293(2) K	$\theta_{\text{min}}$ = 1.7°
$\omega$ scans	$h = -7 \rightarrow 7$
Absorption correction: none	$k = -10 \rightarrow 9$
11620 measured reflections	$l = -31 \rightarrow 30$
2492 independent reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)]$ = 0.074	H-atom parameters constrained
$wR(F^2)$ = 0.229	$w = 1/[\sigma^2(F_o^2) + (0.1605P)^2 + 0.7215P]$ , where $P = (F_o^2 + 2F_c^2)/3$
$S$ = 1.00	$(\Delta/\sigma)_{\text{max}} < 0.001$
2492 reflections	$\Delta\rho_{\text{max}} = 1.08 \text{ e \AA}^{-3}$

136 parameters  $\Delta\rho_{\min} = -0.79 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct Extinction correction: none  
 methods

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4484 (5)	-0.0528 (3)	0.25594 (13)	0.0508 (6)
H1	0.5878	0.0053	0.2547	0.061*
C2	0.3462 (6)	-0.0682 (4)	0.30628 (13)	0.0593 (7)
H2	0.4173	-0.0216	0.3387	0.071*
C3	0.1385 (6)	-0.1527 (4)	0.30829 (13)	0.0604 (7)
H3	0.0683	-0.1617	0.3420	0.072*
C4	0.0359 (5)	-0.2230 (4)	0.26081 (14)	0.0582 (7)
H4	-0.1030	-0.2815	0.2626	0.070*
C5	0.1357 (4)	-0.2085 (3)	0.21013 (12)	0.0498 (6)
H5	0.0632	-0.2555	0.1779	0.060*
C6	0.3459 (4)	-0.1230 (3)	0.20739 (11)	0.0429 (5)
C7	0.4602 (5)	-0.1054 (4)	0.15294 (12)	0.0552 (7)
H7A	0.4431	-0.2135	0.1316	0.066*
H7B	0.6222	-0.0821	0.1600	0.066*
C8	0.5290 (4)	0.1362 (3)	0.06841 (10)	0.0436 (5)
N1	0.7023 (4)	0.0351 (3)	0.05812 (10)	0.0574 (6)
H1A	0.8017	0.0699	0.0355	0.069*
H1B	0.7168	-0.0662	0.0740	0.069*
N2	0.5046 (4)	0.2908 (3)	0.04458 (10)	0.0563 (6)
H2A	0.6031	0.3268	0.0219	0.068*
H2B	0.3900	0.3563	0.0516	0.068*
N3	0.9654 (5)	0.3266 (4)	-0.04824 (12)	0.0660 (7)
O1	0.7787 (6)	0.3965 (5)	-0.05407 (17)	0.1137 (13)
O2	1.0010 (6)	0.2296 (5)	-0.00529 (14)	0.0993 (10)
O3	1.1270 (5)	0.3341 (4)	-0.07925 (11)	0.0832 (8)
S1	0.32436 (11)	0.07656 (10)	0.11423 (3)	0.0538 (3)

### Atomic displacement parameters ( $\text{\AA}^2$ )

$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
----------	----------	----------	----------	----------	----------

## supplementary materials

---

C1	0.0463 (13)	0.0418 (12)	0.0640 (15)	-0.0041 (10)	0.0002 (11)	0.0033 (11)
C2	0.0728 (19)	0.0505 (15)	0.0540 (15)	-0.0015 (13)	-0.0025 (13)	-0.0016 (11)
C3	0.0714 (19)	0.0498 (15)	0.0619 (16)	0.0045 (13)	0.0204 (14)	0.0105 (12)
C4	0.0469 (14)	0.0474 (14)	0.0814 (19)	-0.0040 (10)	0.0136 (13)	0.0093 (13)
C5	0.0441 (12)	0.0435 (12)	0.0613 (14)	-0.0011 (10)	-0.0015 (11)	-0.0005 (11)
C6	0.0398 (11)	0.0346 (10)	0.0547 (13)	0.0050 (8)	0.0067 (9)	0.0049 (9)
C7	0.0588 (16)	0.0471 (13)	0.0612 (15)	0.0176 (11)	0.0168 (12)	0.0101 (11)
C8	0.0415 (12)	0.0452 (12)	0.0444 (11)	0.0058 (9)	0.0048 (9)	-0.0018 (9)
N1	0.0560 (14)	0.0561 (13)	0.0625 (13)	0.0195 (11)	0.0231 (11)	0.0085 (11)
N2	0.0581 (14)	0.0493 (12)	0.0632 (13)	0.0138 (10)	0.0174 (11)	0.0117 (10)
N3	0.0584 (14)	0.0686 (16)	0.0700 (16)	0.0209 (12)	-0.0037 (12)	-0.0270 (13)
O1	0.089 (2)	0.104 (2)	0.146 (3)	0.0491 (18)	-0.008 (2)	-0.031 (2)
O2	0.0851 (19)	0.126 (3)	0.0893 (18)	-0.0035 (19)	0.0280 (15)	0.0245 (19)
O3	0.0965 (19)	0.0875 (18)	0.0691 (14)	0.0096 (15)	0.0337 (13)	0.0199 (13)
S1	0.0424 (4)	0.0579 (5)	0.0624 (5)	0.0142 (2)	0.0143 (3)	0.0142 (3)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—C2	1.381 (4)	C7—H7A	0.9700
C1—C6	1.384 (4)	C7—H7B	0.9700
C1—H1	0.9300	C8—N1	1.307 (3)
C2—C3	1.379 (5)	C8—N2	1.309 (3)
C2—H2	0.9300	C8—S1	1.731 (3)
C3—C4	1.364 (5)	N1—H1A	0.8600
C3—H3	0.9300	N1—H1B	0.8600
C4—C5	1.382 (4)	N2—H2A	0.8600
C4—H4	0.9300	N2—H2B	0.8600
C5—C6	1.396 (3)	N3—O1	1.217 (4)
C5—H5	0.9300	N3—O3	1.237 (4)
C6—C7	1.506 (4)	N3—O2	1.273 (4)
C7—S1	1.821 (3)		
C2—C1—C6	120.8 (3)	C6—C7—H7A	110.2
C2—C1—H1	119.6	S1—C7—H7A	110.2
C6—C1—H1	119.6	C6—C7—H7B	110.2
C1—C2—C3	119.8 (3)	S1—C7—H7B	110.2
C1—C2—H2	120.1	H7A—C7—H7B	108.5
C3—C2—H2	120.1	N1—C8—N2	120.7 (2)
C4—C3—C2	120.0 (3)	N1—C8—S1	122.6 (2)
C4—C3—H3	120.0	N2—C8—S1	116.66 (19)
C2—C3—H3	120.0	C8—N1—H1A	120.0
C3—C4—C5	120.8 (3)	C8—N1—H1B	120.0
C3—C4—H4	119.6	H1A—N1—H1B	120.0
C5—C4—H4	119.6	C8—N2—H2A	120.0
C4—C5—C6	119.8 (3)	C8—N2—H2B	120.0
C4—C5—H5	120.1	H2A—N2—H2B	120.0
C6—C5—H5	120.1	O1—N3—O3	128.7 (4)
C1—C6—C5	118.7 (2)	O1—N3—O2	116.7 (4)
C1—C6—C7	120.0 (2)	O3—N3—O2	114.6 (3)
C5—C6—C7	121.3 (3)	C8—S1—C7	102.89 (12)

C6—C7—S1	107.78 (17)		
C6—C1—C2—C3	−0.7 (4)	C4—C5—C6—C7	179.7 (2)
C1—C2—C3—C4	0.9 (5)	C1—C6—C7—S1	−99.7 (3)
C2—C3—C4—C5	−1.0 (5)	C5—C6—C7—S1	79.9 (3)
C3—C4—C5—C6	0.9 (4)	N1—C8—S1—C7	15.6 (3)
C2—C1—C6—C5	0.6 (4)	N2—C8—S1—C7	−163.5 (2)
C2—C1—C6—C7	−179.8 (2)	C6—C7—S1—C8	158.3 (2)
C4—C5—C6—C1	−0.7 (4)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O2	0.86	1.98	2.803 (4)	160
N1—H1B···O3 <sup>i</sup>	0.86	2.23	3.009 (4)	151
N2—H2A···O1	0.86	2.21	3.040 (5)	164
N2—H2A···O2	0.86	2.56	3.240 (4)	136
N2—H2B···O1 <sup>ii</sup>	0.86	2.12	2.913 (4)	152

Symmetry codes: (i)  $-x+2, -y, -z$ ; (ii)  $-x+1, -y+1, -z$ .

## supplementary materials

---

Fig. 1




Fig. 2

