

(E,E)-1,5-Di-2-thienylpenta-1,4-dien-3-one

S. Murugavel,^a G. Ganesh,^b A. Subbiah Pandi,^{c*}
 Ramalingam Murugan^d and S. Sriman Narayanan^d

^aDepartment of Physics, Thanthai Periyar Government Institute of Technology, Vellore 632 002, India, ^bDepartment of Physics, SMK Fomra Institute of Technology, Thaipur, Chennai 603 103, India, ^cDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India, and ^dDepartment of Analytical Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India

Correspondence e-mail: a_spandian@yahoo.com


Received 30 July 2008; accepted 18 August 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.058; wR factor = 0.211; data-to-parameter ratio = 16.4.

In the title compound, $\text{C}_{13}\text{H}_{10}\text{OS}_2$, the dihedral angle between the thiophene rings is $14.3(1)^\circ$. The molecular structure is stabilized by $\text{C}-\text{H} \cdots \pi$ interactions between a thiophene H atom and an adjacent thiophene ring, and by intermolecular $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds.

Related literature

For the bioactivity of chalcones, see: Go *et al.* (2005). For uses in organic solid-state photochemistry, see: Gould *et al.* (1995); For a related structure, see: Arshad *et al.* (2008). For the nonlinear optical properties of bis-chalcones, see: Uchida *et al.* (1998).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{OS}_2$
 $M_r = 246.35$
 Orthorhombic, $Pbca$
 $a = 12.1141(4)$ Å

$b = 7.4449(3)$ Å
 $c = 27.246(1)$ Å
 $V = 2457.27(16)$ Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.41$ mm⁻¹

$T = 293(2)$ K
 $0.26 \times 0.15 \times 0.15$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.984$, $T_{\max} = 0.987$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.210$
 $S = 1.01$
 2373 reflections

145 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.36$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
C1—H1 ⁱ ···O1 ⁱ	0.93	2.47	3.374 (5)	165
C13—H13 ^j ···O1 ⁱⁱ	0.93	2.33	3.255 (4)	171
C11—H11 ^k ···Cg ⁱⁱⁱ	0.93	3.12	3.936 (5)	148

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (ii) $x - \frac{1}{2}, y, -z + \frac{1}{2}$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, z$. Cg is the centroid of the C10/C11/C12/C13/S1 thiophene ring.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia (1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2003).

SM and ASP thank Dr S. Pandi, Head of the Department of Physics, Presidency College (Autonomous), Chennai, for providing the necessary facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2066).

References

- Arshad, M. N., Tahir, M. N., Asghar, M. N., Khan, I. U. & Ashfaq, M. (2008). *Acta Cryst. E64*, o1413.
- Bruker (2004). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Go, M. L., Wu, X. & Liu, X. L. (2005). *Curr. Med. Chem. 12*, 483–499.
- Gould, B. S., Panneerelvam, K., Zacharias, D. E. & Desiraju, G. R. (1995). *J. Chem. Soc. Perkin Trans. 2*, pp. 325–330.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.
- Uchida, T., Kozawa, K., Sakai, T., Aoki, M., Yoguchi, H., Abduryim, A. & Watanabe, Y. (1998). *Mol. Cryst. Liq. Cryst. 315*, 135–140.

supplementary materials

Acta Cryst. (2008). E64, o1807 [doi:10.1107/S1600536808026603]

(E,E)-1,5-Di-2-thienylpenta-1,4-dien-3-one

S. Murugavel, G. Ganesh, A. SubbiahPandi, R. Murugan and S. SrimanNarayanan

Comment

Chalcones with the general formula Ar—CH=CH—CO—Ar are an important class of compounds, with the common structural entity being the central —CH=CH—C(=O)— group, in the H atoms can be substituted. The —C=C— double bond can be photoreactive and can produce various products through solid-state photo cycloaddition. Therefore, chalcones are widely used in organic solid-state photochemistry (Gould *et al.*, 1995). Reviews on the bioactivities of various chalcones have been reported (Go *et al.*, 2005). Bis-chalcones are also found to exhibit good NLO properties (Uchida *et al.*, 1998). In view of this biological importance, the crystal structure of the title compound (I), (1E, 4E)-1,5-Bis(2-thienyl)penta-1,4-dien-3-one (Fig. 1) has been determined and the results are presented here.

Compound (I) consists of two thiophene rings A and B. The central position of (I) shows double and single bonds orientating from O1 atom and five C atom behave like a backbone. The planarities of rings A and B are fairly good. The bond lengths in the (I) are normal and comparable with the corresponding values observed in the related structure (Arshad *et al.*, 2008). The dihedral angle between the two thiophene rings is 14.3 (1) $^{\circ}$. The crystal packing (Fig. 2) is stabilized by C—H \cdots π interactions between a thiophene H atom and a neighbouring thiophene ring, with a C11—H11 \cdots Cgⁱⁱⁱ separation of 2.34 Å (Fig. 2 and Table 1; Cg is the centroid of the C10/C11/C12/C13/S1 thiophene ring, symmetry code as in Fig. 2). The molecular packing is further stabilized by intermolecular C—H \cdots O hydrogen bonds (Fig. 2 and Table 1; symmetry code as in Fig. 2).

Experimental

A solution of sodium hydroxide (10 g, 0.25 mol) in water (50 ml) was added to a solution of acetone (5 ml) and 2-thiophene-carboxaldehyde (22.4 g, 0.2 mol) in methanol (50 ml) at 273 K. This mixture was stirred overnight and the product was filtered. Single crystals suitable for X-ray diffraction was obtained by slow evaporation of a solution of the title compound in ethyl acetate.

Refinement

All H atoms were fixed geometrically and allowed to ride on their parent atoms, with N—H=0.86 Å and C—H=0.93–0.98 Å with $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}$ (methyl H) and $1.2U_{\text{eq}}$ (for other H atoms).

Figures




Fig. 1. The molecular structure of title compound showing 30% probability displacement ellipsoids.

supplementary materials




Fig. 2. C—H··· π and C—H···O interactions (dotted lines) in the title compound. Cg denotes the ring centroid. [Symmetry code: (i) x-1/2, -y+1/2, -z+1; (ii)x-1/2, y, -z+1/2; (iii) x-1/2, y, -z+1/2; (iv) -x+3/2, y-1/2, z; (v)-x+1, y+1/2, -z+1/2.]

(E,E)-1,5-Di-2-thienylpenta-1,4-dien-3-one

Crystal data

C ₁₃ H ₁₀ OS ₂	$F_{000} = 1024$
$M_r = 246.35$	$D_x = 1.332 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
Hall symbol: -P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 12.1141 (4) \text{ \AA}$	Cell parameters from 10259 reflections
$b = 7.4449 (3) \text{ \AA}$	$\theta = 2.3\text{--}30.3^\circ$
$c = 27.246 (1) \text{ \AA}$	$\mu = 0.41 \text{ mm}^{-1}$
$V = 2457.27 (16) \text{ \AA}^3$	$T = 293 (2) \text{ K}$
$Z = 8$	Block, colourless
	$0.26 \times 0.15 \times 0.15 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	2373 independent reflections
Radiation source: fine-focus sealed tube	1719 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
Detector resolution: 10 pixels mm ⁻¹	$\theta_{\text{max}} = 26.0^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 1.5^\circ$
ω scans	$h = -11\text{--}14$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$k = -7\text{--}9$
$T_{\text{min}} = 0.984$, $T_{\text{max}} = 0.987$	$l = -32\text{--}31$
13976 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H-atom parameters constrained
$wR(F^2) = 0.211$	$w = 1/[\sigma^2(F_o^2) + (0.1295P)^2 + 1.4091P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
2373 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
145 parameters	$\Delta\rho_{\text{max}} = 0.49 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.52710 (7)	0.54328 (13)	0.23889 (4)	0.0733 (4)
S2	0.53529 (8)	0.23805 (14)	0.51370 (4)	0.0801 (4)
O1	0.82061 (18)	0.5548 (4)	0.38920 (9)	0.0734 (7)
C1	0.5312 (3)	0.2113 (6)	0.57495 (18)	0.0876 (13)
H1	0.4768	0.1455	0.5910	0.105*
C2	0.6135 (4)	0.2934 (6)	0.59817 (16)	0.0907 (12)
H2	0.6209	0.2904	0.6321	0.109*
C3	0.6903 (3)	0.3874 (5)	0.56716 (12)	0.0667 (9)
H3	0.7525	0.4507	0.5773	0.080*
C4	0.6519 (3)	0.3642 (4)	0.51665 (13)	0.0612 (8)
C5	0.7055 (3)	0.4350 (4)	0.47339 (13)	0.0610 (8)
H5	0.7755	0.4843	0.4774	0.073*
C6	0.6638 (3)	0.4359 (4)	0.42854 (13)	0.0612 (8)
H6	0.5938	0.3881	0.4236	0.073*
C7	0.7236 (2)	0.5093 (4)	0.38617 (12)	0.0573 (8)
C8	0.6611 (2)	0.5283 (4)	0.34019 (12)	0.0594 (8)
H8	0.5901	0.4810	0.3384	0.071*
C9	0.7027 (2)	0.6109 (4)	0.30106 (12)	0.0565 (7)
H9	0.7733	0.6583	0.3045	0.068*
C10	0.6513 (2)	0.6354 (4)	0.25413 (12)	0.0536 (7)
C11	0.6974 (2)	0.7299 (4)	0.21337 (11)	0.0541 (7)
H11	0.7653	0.7882	0.2132	0.065*
C12	0.6228 (3)	0.7207 (6)	0.17301 (14)	0.0794 (10)
H12	0.6364	0.7764	0.1431	0.095*
C13	0.5322 (3)	0.6260 (6)	0.18183 (15)	0.0774 (11)
H13	0.4771	0.6074	0.1586	0.093*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0542 (6)	0.0750 (6)	0.0909 (8)	-0.0090 (4)	-0.0089 (4)	-0.0068 (5)

supplementary materials

S2	0.0619 (6)	0.0821 (7)	0.0962 (9)	-0.0065 (4)	0.0047 (4)	0.0076 (5)
O1	0.0483 (13)	0.1008 (18)	0.0710 (16)	-0.0067 (11)	0.0009 (10)	-0.0029 (12)
C1	0.069 (2)	0.090 (3)	0.103 (3)	0.0109 (19)	0.018 (2)	0.033 (2)
C2	0.091 (3)	0.113 (3)	0.068 (2)	0.012 (3)	0.000 (2)	0.013 (2)
C3	0.071 (2)	0.073 (2)	0.056 (2)	-0.0057 (16)	-0.0001 (15)	0.0110 (15)
C4	0.0558 (17)	0.0542 (15)	0.074 (2)	0.0020 (13)	0.0024 (14)	0.0003 (14)
C5	0.0527 (16)	0.0593 (17)	0.071 (2)	-0.0027 (13)	-0.0005 (15)	-0.0028 (14)
C6	0.0510 (16)	0.0621 (17)	0.071 (2)	-0.0018 (13)	0.0012 (14)	-0.0036 (15)
C7	0.0497 (17)	0.0576 (16)	0.065 (2)	0.0021 (13)	0.0033 (13)	-0.0081 (14)
C8	0.0467 (15)	0.0646 (17)	0.067 (2)	-0.0029 (13)	0.0012 (14)	-0.0081 (15)
C9	0.0435 (14)	0.0592 (16)	0.067 (2)	-0.0007 (12)	-0.0031 (13)	-0.0097 (14)
C10	0.0457 (15)	0.0511 (14)	0.0641 (19)	0.0013 (12)	-0.0020 (13)	-0.0103 (13)
C11	0.0442 (14)	0.0624 (17)	0.0558 (18)	-0.0046 (12)	-0.0030 (12)	-0.0082 (13)
C12	0.081 (2)	0.096 (3)	0.062 (2)	0.013 (2)	0.0044 (18)	-0.0026 (18)
C13	0.063 (2)	0.091 (3)	0.078 (3)	0.0085 (18)	-0.0146 (17)	-0.022 (2)

Geometric parameters (Å, °)

S1—C13	1.673 (5)	C6—C7	1.468 (5)
S1—C10	1.704 (3)	C6—H6	0.9300
S2—C1	1.682 (5)	C7—C8	1.471 (5)
S2—C4	1.699 (3)	C8—C9	1.330 (5)
O1—C7	1.225 (4)	C8—H8	0.9300
C1—C2	1.330 (6)	C9—C10	1.434 (4)
C1—H1	0.9300	C9—H9	0.9300
C2—C3	1.438 (5)	C10—C11	1.429 (4)
C2—H2	0.9300	C11—C12	1.425 (5)
C3—C4	1.463 (5)	C11—H11	0.9300
C3—H3	0.9300	C12—C13	1.326 (5)
C4—C5	1.445 (5)	C12—H12	0.9300
C5—C6	1.322 (5)	C13—H13	0.9300
C5—H5	0.9300		
C13—S1—C10	92.62 (17)	O1—C7—C8	121.6 (3)
C1—S2—C4	92.5 (2)	C6—C7—C8	116.8 (3)
C2—C1—S2	113.3 (3)	C9—C8—C7	122.1 (3)
C2—C1—H1	123.3	C9—C8—H8	118.9
S2—C1—H1	123.3	C7—C8—H8	118.9
C1—C2—C3	115.4 (4)	C8—C9—C10	127.5 (3)
C1—C2—H2	122.3	C8—C9—H9	116.3
C3—C2—H2	122.3	C10—C9—H9	116.3
C2—C3—C4	106.8 (3)	C9—C10—C11	125.9 (3)
C2—C3—H3	126.6	C9—C10—S1	123.3 (2)
C4—C3—H3	126.6	C11—C10—S1	110.7 (2)
C5—C4—C3	125.6 (3)	C12—C11—C10	109.1 (3)
C5—C4—S2	122.5 (3)	C12—C11—H11	125.4
C3—C4—S2	112.0 (2)	C10—C11—H11	125.4
C6—C5—C4	125.7 (3)	C13—C12—C11	114.2 (4)
C6—C5—H5	117.1	C13—C12—H12	122.9
C4—C5—H5	117.1	C11—C12—H12	122.9

C5—C6—C7	122.7 (3)	C12—C13—S1	113.3 (3)
C5—C6—H6	118.7	C12—C13—H13	123.4
C7—C6—H6	118.7	S1—C13—H13	123.4
O1—C7—C6	121.5 (3)		
C4—S2—C1—C2	0.3 (4)	O1—C7—C8—C9	-6.5 (5)
S2—C1—C2—C3	-0.5 (5)	C6—C7—C8—C9	172.5 (3)
C1—C2—C3—C4	0.5 (5)	C7—C8—C9—C10	178.8 (3)
C2—C3—C4—C5	-179.2 (3)	C8—C9—C10—C11	177.9 (3)
C2—C3—C4—S2	-0.3 (4)	C8—C9—C10—S1	-5.4 (4)
C1—S2—C4—C5	179.0 (3)	C13—S1—C10—C9	-177.9 (3)
C1—S2—C4—C3	0.1 (3)	C13—S1—C10—C11	-0.8 (2)
C3—C4—C5—C6	-170.2 (3)	C9—C10—C11—C12	178.5 (3)
S2—C4—C5—C6	11.1 (5)	S1—C10—C11—C12	1.5 (3)
C4—C5—C6—C7	-179.5 (3)	C10—C11—C12—C13	-1.6 (4)
C5—C6—C7—O1	8.6 (5)	C11—C12—C13—S1	1.0 (4)
C5—C6—C7—C8	-170.4 (3)	C10—S1—C13—C12	-0.1 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C1—H1···O1 ⁱ	0.93	2.47	3.374 (5)	165
C13—H13···O1 ⁱⁱ	0.93	2.33	3.255 (4)	171
C11—H11···Cg ⁱⁱⁱ	0.93	3.12	3.936 (5)	148

Symmetry codes: (i) $x-1/2, -y+1/2, -z+1$; (ii) $x-1/2, y, -z+1/2$; (iii) $-x+3/2, y+1/2, z$.

supplementary materials

Fig. 1




Fig. 2

