

Quinoxalin-2-yl *o*-tolyl ether

Nor Duha Hassan, Hairul Anuar Tajuddin, Zanariah Abdullah and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 12 August 2008; accepted 19 August 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.066; wR factor = 0.184; data-to-parameter ratio = 12.1.

The dihedral angle between the two aromatic ring systems in the title compound, $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}$, is $85.9(1)^\circ$; The angle at the O atom is widened to $118.2(2)^\circ$. The quinoxalinyloxy part of the molecule lies on a mirror plane and the tolyl group is disordered over two positions about the mirror plane.

Related literature

The title compound exhibits fluorescence; see: Abdullah (2005); Kawai *et al.* (2001); Mohd Salleh *et al.* (2007).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}$	$V = 587.1(2) \text{ \AA}^3$
$M_r = 236.27$	$Z = 2$
Monoclinic, $P2_1/m$	Mo $K\alpha$ radiation
$a = 7.874(2) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$b = 7.413(1) \text{ \AA}$	$T = 100(2) \text{ K}$
$c = 10.596(2) \text{ \AA}$	$0.20 \times 0.20 \times 0.08 \text{ mm}$
$\beta = 108.332(3)^\circ$	

Data collection

Bruker SMART APEX diffractometer	1443 independent reflections
Absorption correction: none	1032 reflections with $I > 2\sigma(I)$
3392 measured reflections	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$	119 parameters
$wR(F^2) = 0.183$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
1443 reflections	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the University of Malaya for supporting this study (grant No. F2358/2008 A).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2769).

References

- Abdullah, Z. (2005). *Int. J. Chem. Sci.* **3**, 9–15.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Kawai, M., Lee, M. J., Evans, K. O. & Norlund, T. (2001). *J. Fluoresc.* **11**, 23–32.
 Mohd Salleh, N., Ling, L. P., Abdullah, Z. M. A. A. & Aiyub, Z. (2007). *Malays. J. Anal. Sci.* **11**, 229–236.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2008). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2008). E64, o1821 [doi:10.1107/S1600536808026810]

Quinoxalin-2-yl *o*-tolyl ether

N. D. Hassan, H. A. Tajuddin, Z. Abdullah and S. W. Ng

Comment

(type here to add)

Experimental

o-Cresol (0.54 g, 5 mmol) was dissolved in a small volume of water containing potassium hydroxide (0.20 g, 5 mmol). The mixture was heated to remove the water to give a brown compound. The compound and 2-chloroquinoxaline (0.82, g, 5 mmol) were heated in THF (15 ml) for 8 h. The mixture was in 1 N sodium hydroxide; the aqueous solution was extracted with dichloromethane. The organic phase was dried over sodium sulfate. Evaporation of the solvent gave a yellow product, which was washed with chloroform to remove impurities. Crystals were obtained upon recrystallization from an ethyl acetate/hexane mixture.

Refinement

The tolyl group is disordered about a mirror plane; the phenylene ring was refined as a rigid hexagon. The seven carbon atoms were allowed to refine off the symmetry element; the atoms were all given 0.5 site occupancy.

H-atoms were placed in calculated positions (C—H 0.95–0.98 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ fixed at 1.2–1.5 $U(\text{C})$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Quinoxalin-2-yl *o*-tolyl ether

Crystal data

$\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}$

$M_r = 236.27$

Monoclinic, $P2_1/m$

Hall symbol: -P 2yb

$a = 7.874(2)$ Å

$F_{000} = 248$

$D_x = 1.337$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 756 reflections

$\theta = 2.7\text{--}24.6^\circ$

supplementary materials

$b = 7.413 (1) \text{ \AA}$
 $c = 10.596 (2) \text{ \AA}$
 $\beta = 108.332 (3)^\circ$
 $V = 587.1 (2) \text{ \AA}^3$
 $Z = 2$

$\mu = 0.09 \text{ mm}^{-1}$
 $T = 100 (2) \text{ K}$
Block, colorless
 $0.20 \times 0.20 \times 0.08 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 100(2) \text{ K}$
 ω scans
Absorption correction: None
3392 measured reflections
1443 independent reflections

1032 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$
 $\theta_{\text{max}} = 27.5^\circ$
 $\theta_{\text{min}} = 2.0^\circ$
 $h = -10 \rightarrow 8$
 $k = -9 \rightarrow 8$
 $l = -12 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.183$
 $S = 1.03$
1443 reflections
119 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0776P)^2 + 0.4106P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.1778 (2)	0.2500	0.64890 (18)	0.0405 (6)	
N1	0.4752 (3)	0.2500	0.7791 (2)	0.0552 (9)	
N2	0.3518 (3)	0.2500	0.9997 (2)	0.0479 (8)	
C1	0.2266 (3)	0.1943 (3)	0.5368 (2)	0.0256 (8)	0.50
C2	0.2256 (5)	0.0171 (3)	0.4936 (3)	0.047 (2)	0.50
H2	0.2002	-0.0787	0.5445	0.056*	0.50
C3	0.2619 (5)	-0.0199 (3)	0.3761 (3)	0.0473 (11)	0.50
H3	0.2612	-0.1410	0.3466	0.057*	0.50
C4	0.2991 (4)	0.1203 (4)	0.3017 (2)	0.0417 (12)	0.50
H4	0.3239	0.0950	0.2213	0.050*	0.50
C5	0.3000 (3)	0.2975 (3)	0.3449 (2)	0.0345 (15)	0.50
H5	0.3255	0.3933	0.2940	0.041*	0.50

C6	0.2638 (3)	0.3345 (3)	0.4624 (3)	0.0313 (8)	0.50
C7	0.2599 (11)	0.5228 (12)	0.5056 (7)	0.0496 (17)	0.50
H7A	0.3449	0.5946	0.4761	0.074*	0.50
H7B	0.2932	0.5272	0.6028	0.074*	0.50
H7C	0.1391	0.5719	0.4666	0.074*	0.50
C8	0.3083 (3)	0.2500	0.7686 (2)	0.0300 (6)	
C9	0.2449 (4)	0.2500	0.8784 (3)	0.0393 (8)	
H9	0.1194	0.2500	0.8633	0.047*	
C10	0.5314 (3)	0.2500	1.0167 (2)	0.0284 (6)	
C11	0.6553 (4)	0.2500	1.1448 (3)	0.0429 (8)	
H11	0.6135	0.2500	1.2197	0.051*	
C12	0.8343 (4)	0.2500	1.1638 (3)	0.0398 (8)	
H12	0.9169	0.2500	1.2513	0.048*	
C13	0.8952 (4)	0.2500	1.0547 (3)	0.0631 (12)	
H13	1.0201	0.2500	1.0674	0.076*	
C14	0.7769 (4)	0.2500	0.9289 (3)	0.0889 (19)	
H14	0.8209	0.2500	0.8551	0.107*	
C15	0.5922 (3)	0.2500	0.9066 (3)	0.0378 (7)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0199 (9)	0.0771 (16)	0.0236 (9)	0.000	0.0055 (7)	0.000
N1	0.0196 (11)	0.123 (3)	0.0224 (12)	0.000	0.0065 (9)	0.000
N2	0.0248 (12)	0.093 (2)	0.0270 (12)	0.000	0.0090 (9)	0.000
C1	0.0198 (14)	0.033 (2)	0.0229 (16)	0.0038 (12)	0.0051 (12)	0.0008 (12)
C2	0.057 (4)	0.056 (5)	0.031 (3)	0.004 (3)	0.018 (3)	0.002 (3)
C3	0.072 (3)	0.041 (3)	0.032 (2)	0.015 (2)	0.021 (2)	0.0047 (19)
C4	0.040 (2)	0.064 (4)	0.024 (2)	0.015 (2)	0.0142 (17)	0.002 (2)
C5	0.0231 (15)	0.054 (5)	0.0263 (17)	-0.0015 (15)	0.0075 (13)	0.0121 (16)
C6	0.0175 (15)	0.0301 (19)	0.039 (2)	-0.0008 (15)	-0.0015 (15)	0.0020 (17)
C7	0.055 (3)	0.037 (3)	0.054 (4)	-0.001 (3)	0.013 (3)	0.008 (3)
C8	0.0220 (12)	0.0435 (16)	0.0233 (12)	0.000	0.0053 (10)	0.000
C9	0.0188 (12)	0.069 (2)	0.0304 (14)	0.000	0.0084 (11)	0.000
C10	0.0230 (12)	0.0369 (15)	0.0259 (13)	0.000	0.0086 (10)	0.000
C11	0.0303 (14)	0.076 (2)	0.0227 (13)	0.000	0.0093 (11)	0.000
C12	0.0265 (14)	0.065 (2)	0.0243 (13)	0.000	0.0020 (11)	0.000
C13	0.0211 (14)	0.137 (4)	0.0291 (15)	0.000	0.0053 (12)	0.000
C14	0.0238 (15)	0.221 (6)	0.0247 (16)	0.000	0.0111 (13)	0.000
C15	0.0218 (13)	0.067 (2)	0.0236 (13)	0.000	0.0063 (10)	0.000

Geometric parameters (\AA , $^\circ$)

O1—C8	1.359 (3)	C6—C7	1.472 (9)
O1—C1	1.420 (3)	C7—H7A	0.9800
O1—C1 ⁱ	1.420 (3)	C7—H7B	0.9800
N1—C8	1.283 (3)	C7—H7C	0.9800
N1—C15	1.375 (3)	C8—C9	1.402 (4)

supplementary materials

N2—C9	1.296 (4)	C9—H9	0.9500
N2—C10	1.369 (3)	C10—C15	1.392 (4)
C1—C2	1.3900	C10—C11	1.400 (4)
C1—C6	1.3900	C11—C12	1.360 (4)
C2—C3	1.3900	C11—H11	0.9500
C2—H2	0.9500	C12—C13	1.384 (4)
C3—C4	1.3900	C12—H12	0.9500
C3—H3	0.9500	C13—C14	1.365 (4)
C4—C5	1.3900	C13—H13	0.9500
C4—H4	0.9500	C14—C15	1.398 (4)
C5—C6	1.3900	C14—H14	0.9500
C5—H5	0.9500		
C8—O1—C1	117.19 (19)	O1—C8—C9	114.4 (2)
C8—N1—C15	115.8 (2)	N2—C9—C8	122.2 (2)
C9—N2—C10	116.9 (2)	N2—C9—H9	118.9
C2—C1—C6	120.0	C8—C9—H9	118.9
C2—C1—O1	125.13 (17)	N2—C10—C15	120.2 (2)
C6—C1—O1	114.68 (17)	N2—C10—C11	120.2 (2)
C1—C2—C3	120.0	C15—C10—C11	119.6 (2)
C1—C2—H2	120.0	C12—C11—C10	121.2 (3)
C3—C2—H2	120.0	C12—C11—H11	119.4
C2—C3—C4	120.0	C10—C11—H11	119.4
C2—C3—H3	120.0	C11—C12—C13	119.4 (3)
C4—C3—H3	120.0	C11—C12—H12	120.3
C5—C4—C3	120.0	C13—C12—H12	120.3
C5—C4—H4	120.0	C14—C13—C12	120.4 (3)
C3—C4—H4	120.0	C14—C13—H13	119.8
C4—C5—C6	120.0	C12—C13—H13	119.8
C4—C5—H5	120.0	C13—C14—C15	121.3 (3)
C6—C5—H5	120.0	C13—C14—H14	119.4
C5—C6—C1	120.0	C15—C14—H14	119.4
C5—C6—C7	119.7 (3)	N1—C15—C10	121.5 (2)
C1—C6—C7	120.3 (3)	N1—C15—C14	120.4 (2)
N1—C8—O1	122.3 (2)	C10—C15—C14	118.1 (3)
N1—C8—C9	123.4 (2)		
C8—O1—C1—C2	87.53 (18)	C1 ⁱ —O1—C8—C9	160.91 (12)
C1 ⁱ —O1—C1—C2	-173.48 (14)	C10—N2—C9—C8	0.000 (2)
C8—O1—C1—C6	-97.56 (16)	N1—C8—C9—N2	0.000 (2)
C1 ⁱ —O1—C1—C6	1.43 (19)	O1—C8—C9—N2	180.000 (1)
C6—C1—C2—C3	0.0	C9—N2—C10—C15	0.000 (2)
O1—C1—C2—C3	174.7 (2)	C9—N2—C10—C11	180.000 (2)
C1—C2—C3—C4	0.0	N2—C10—C11—C12	180.000 (2)
C2—C3—C4—C5	0.0	C15—C10—C11—C12	0.000 (2)
C3—C4—C5—C6	0.0	C10—C11—C12—C13	0.000 (2)
C4—C5—C6—C1	0.0	C11—C12—C13—C14	0.000 (2)
C4—C5—C6—C7	-178.3 (4)	C12—C13—C14—C15	0.000 (2)
C2—C1—C6—C5	0.0	C8—N1—C15—C10	0.000 (1)
O1—C1—C6—C5	-175.2 (2)	C8—N1—C15—C14	180.000 (1)

supplementary materials

C2—C1—C6—C7	178.2 (4)	N2—C10—C15—N1	0.000 (2)
O1—C1—C6—C7	3.0 (4)	C11—C10—C15—N1	180.000 (1)
C15—N1—C8—O1	180.000 (1)	N2—C10—C15—C14	180.000 (2)
C15—N1—C8—C9	0.000 (1)	C11—C10—C15—C14	0.000 (2)
C1—O1—C8—N1	19.09 (12)	C13—C14—C15—N1	180.000 (2)
C1 ⁱ —O1—C8—N1	-19.09 (12)	C13—C14—C15—C10	0.000 (2)
C1—O1—C8—C9	-160.91 (12)		

Symmetry codes: (i) $x, -y+1/2, z$.

Fig. 1

