

Diaquabis(4-bromo-2-formylphenolato- κ^2O,O')cobalt(II)

Yu Xiao* and Min Zhang

The Guangxi Key Laboratory of Environmental Engineering, Protection and Assessment (Department of Resources and Environmental Engineering, Guilin University of Technology), Guilin 541004, People's Republic of China
Correspondence e-mail: xiaoyuzh@yahoo.com.cn


Received 4 August 2008; accepted 13 August 2008

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.041; wR factor = 0.095; data-to-parameter ratio = 14.7.

In the title complex, $[\text{Co}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{H}_2\text{O})_2]$, the Co^{II} ion, which lies on a crystallographic inversion center, is coordinated by four O atoms from two bidentate 4-bromo-2-formylphenolato ligands and two O atoms from two water ligands in a slightly distorted octahedral environment. In the crystal structure, one-dimensional chains are formed through intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, which are further linked into a two-dimensional network through $\text{Br}\cdots\text{Br}$ interactions [$\text{Br}\cdots\text{Br} = 3.772(4)\text{ \AA}$].

Related literature

For related literature, see: Cohen *et al.* (1964); Desiraju (1989); Mathews & Manohar (1991); Willey *et al.* (1994); Zaman *et al.* (2004); Zhang *et al.* (2007); Zordan *et al.* (2005); Chen *et al.* (2008).



Experimental

Crystal data

$[\text{Co}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{H}_2\text{O})_2]$	$V = 1585.3(4)\text{ \AA}^3$
$M_w = 494.99$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 29.527(5)\text{ \AA}$	$\mu = 6.15\text{ mm}^{-1}$
$b = 4.7406(8)\text{ \AA}$	$T = 293(2)\text{ K}$
$c = 11.6314(18)\text{ \AA}$	$0.21 \times 0.19 \times 0.19\text{ mm}$
$\beta = 103.162(3)^\circ$	

Data collection

Bruker SMART-CCD diffractometer	1553 independent reflections
Absorption correction: none	1290 reflections with $I > 2\sigma(I)$
3884 measured reflections	$R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	106 parameters
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.55\text{ e \AA}^{-3}$
1553 reflections	$\Delta\rho_{\text{min}} = -0.32\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Co1—O2	2.013 (2)	Co1—O3	2.149 (3)
Co1—O1	2.099 (2)		
O2 ⁱ —Co1—O2	180	O1—Co1—O3 ⁱ	86.83 (10)
O2—Co1—O1 ⁱ	87.86 (10)	O2—Co1—O3	89.80 (10)
O2—Co1—O1 ⁱ	92.14 (10)	O1—Co1—O3	93.17 (10)
O1—Co1—O1 ⁱ	180	O3 ⁱ —Co1—O3	180
O2—Co1—O3 ⁱ	90.20 (10)		

Symmetry code: (i) $-x, -y, -z + 1$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3 ⁱⁱ —O1 ⁱⁱ	0.85	2.12	2.842 (4)	142
O3—H3B ⁱⁱⁱ —O2 ⁱⁱⁱ	0.85	1.93	2.725 (4)	155

Symmetry codes: (ii) $-x, -y - 1, -z + 1$; (iii) $x, y - 1, z$.

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *ORTEPIII* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

We acknowledge financial support by the Guangxi Key Laboratory of Environmental Engineering, Protection and Assessment, Guangxi, People's Republic of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2677).

References

- Bruker (2004). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Chen, F.-Y., Zhang, S.-H. & Ge, C.-M. (2008). *Acta Cryst. E64*, m1068.
- Cohen, M. D., Schmidt, G. M. J. & Sonntag, F. I. (1964). *J. Chem. Soc.* pp. 2000–2013.
- Desiraju, G. R. (1989). *Crystal Engineering: The Design of Organic Solids*. Amsterdam: Elsevier.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Mathews, I. I. & Manohar, H. (1991). *Acta Cryst. C47*, 1621–1624.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Willey, G. R., Palin, J., Lakin, M. T. & Alcock, N. W. (1994). *Transition Met. Chem. 19*, 187–190.
- Zaman, B., Udashin, K. A. & Ripmeester, J. A. (2004). *Cryst. Growth Des. 4*, 585–589.
- Zhang, S.-H., Li, G.-Z., Feng, X.-Z. & Liu, Z. (2007). *Acta Cryst. E63*, m1319–m1320.
- Zordan, F., Brammer, L. & Sherwood, P. (2005). *J. Am. Chem. Soc. 127*, 5979–5989.

supplementary materials

Acta Cryst. (2008). E64, m1231 [doi:10.1107/S1600536808026068]

Diaquabis(4-bromo-2-formylphenolato- κ^2O,O')cobalt(II)

Y. Xiao and M. Zhang

Comment

Halogens have a ubiquitous presence in both inorganic and organic chemistry. Schiff bases of bromo substituents on aromatic groups have aroused increasing interest in recent years because these halogenated compounds are an attractive target for use in supramolecular chemistry and crystal engineering wherein the halogen atoms are directly involved in forming intermolecular interactions (Cohen *et al.*, 1964; Zordan *et al.*, 2005; Desiraju, *et al.* 1989; Zaman *et al.*, 2004; Zhang, *et al.*, 2007; Chen, *et al.*, 2008). The title compound, (I), contains the bromo ligand 5-bromo-2-hydroxy-benzaldehyde, with one Br atom accessible at the periphery of each ligand.

In the molecular structure of (I), the Co^{II} ion is coordinated by four O atoms from two bidentate 5-bromo-2-hydroxy-benzaldehyde ligands and two O atoms from two H₂O ligands forming a slightly distorted octahedral geometry (Fig. 1). In the crystal structure, 1-D chains are formed through O–H···O hydrogen bonds (O₃···O1ⁱ, 2.842 (4) Å; O₃···O2ⁱⁱ, 2.725 (4); symmetry codes: (i) -x, -y-1, -z+1; (ii) x, y-1, z). Each molecule of (I) forms eight hydrogen bonds, four of which are donor hydrogen bonds and four are acceptor hydrogen bonds. The 1-D chains are further linked into a 2-D network *via* Br₁···Br₁ interactions. The shortest Br₁···Br₁ distance is 3.772 Å, (Mathews & Manohar, 1991; Willey *et al.*, 1994) observed between Br₁ and Br₁ⁱⁱⁱ, Br₁ and Br₁^{iv} [symmetry codes: (iii) 1/2-x, -1/2+y, 1/2-z; (iv) 1/2-x, 1/2+y, 1/2-z].

Experimental

Distilled water (30 ml) containing 5-bromo-2-hydroxy-benzaldehyde (0.201 g, 1 mmol) was dropwise added to an aqueous solution containing amino-methanesulfonic acid (0.111 g, 1 mmol) and sodium hydroxide (0.040 g, 1 mmol) with stirred during 10 min. After stirring for 1 h, an aqueous solution of cobalt chloride (0.237 g, 1 mmol) was added to the resulting solution and stirred for 2 h and filtrate. the filtration was left to stand at room temperature. After 12 days, red crystals were produced from the filtrate (yield: 76.4 %, based on Co).

Refinement

H atoms were positioned geometrically and were treated as riding atoms, with C–H distances of 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$, and with O–H distance of 0.85 Å and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. A view of (I), showing 30% probability displacement ellipsoids [symmetry code: (A) -x, -y, -z+1]

supplementary materials




Fig. 2. 1-D chain of (I). Dashed lines indicate hydrogen bonds.



Fig. 3. 2-D structure of (I). Blue dashed lines indicate Br...Br interactions and yellow dashed lines show hydrogen bonds.

Diaquabis(4-bromo-2-formylphenolato- κ^2 O,O')cobalt(II)

Crystal data

[Co(C ₇ H ₄ BrO ₂) ₂ (H ₂ O) ₂]	$F_{000} = 964$
$M_r = 494.99$	$D_x = 2.074 \text{ Mg m}^{-3}$
Monoclinic, C2/c	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 29.527 (5) \text{ \AA}$	Cell parameters from 3884 reflections
$b = 4.7406 (8) \text{ \AA}$	$\theta = 2.8\text{--}26.0^\circ$
$c = 11.6314 (18) \text{ \AA}$	$\mu = 6.15 \text{ mm}^{-1}$
$\beta = 103.162 (3)^\circ$	$T = 293 (2) \text{ K}$
$V = 1585.3 (4) \text{ \AA}^3$	Prism, red
$Z = 4$	$0.21 \times 0.19 \times 0.19 \text{ mm}$

Data collection

Bruker SMART-CCD diffractometer	1290 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.033$
Monochromator: graphite	$\theta_{\text{max}} = 26.0^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 2.8^\circ$
φ and ω scans	$h = -27 \rightarrow 36$
Absorption correction: none	$k = -5 \rightarrow 5$
3884 measured reflections	$l = -13 \rightarrow 14$
1553 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.095$	$w = 1/[\sigma^2(F_o^2) + (0.0409P)^2 + 2.4257P]$
	where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.04$	$(\Delta/\sigma)_{\max} < 0.001$
1553 reflections	$\Delta\rho_{\max} = 0.55 \text{ e \AA}^{-3}$
106 parameters	$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.0000	0.0000	0.5000	0.0288 (2)
Br1	0.222905 (17)	0.41219 (14)	0.34123 (5)	0.0652 (2)
O1	0.03315 (9)	-0.2051 (5)	0.3818 (2)	0.0354 (6)
O2	0.05893 (9)	0.2221 (5)	0.5574 (2)	0.0333 (6)
O3	0.02353 (10)	-0.3049 (5)	0.6373 (2)	0.0377 (6)
H3B	0.0416	-0.4195	0.6135	0.057*
H3	0.0003	-0.3971	0.6490	0.057*
C1	0.09460 (13)	0.2474 (8)	0.5103 (3)	0.0309 (8)
C2	0.13013 (14)	0.4390 (9)	0.5592 (4)	0.0405 (10)
H2	0.1280	0.5390	0.6266	0.049*
C3	0.16768 (15)	0.4831 (10)	0.5110 (4)	0.0447 (11)
H3A	0.1908	0.6094	0.5463	0.054*
C4	0.17148 (14)	0.3393 (10)	0.4089 (4)	0.0421 (10)
C5	0.13865 (13)	0.1447 (9)	0.3593 (3)	0.0377 (9)
H5	0.1419	0.0447	0.2928	0.045*
C6	0.09989 (13)	0.0949 (8)	0.4087 (3)	0.0304 (8)
C7	0.06866 (15)	-0.1242 (8)	0.3544 (3)	0.0367 (9)
H7	0.0763	-0.2157	0.2907	0.044*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0330 (4)	0.0264 (4)	0.0284 (4)	-0.0043 (3)	0.0104 (3)	-0.0022 (3)
Br1	0.0404 (3)	0.1038 (5)	0.0569 (3)	-0.0135 (3)	0.0226 (2)	0.0066 (3)
O1	0.0411 (16)	0.0322 (14)	0.0362 (15)	-0.0047 (12)	0.0157 (12)	-0.0056 (11)
O2	0.0323 (15)	0.0353 (15)	0.0340 (14)	-0.0067 (12)	0.0114 (12)	-0.0080 (11)
O3	0.0474 (17)	0.0312 (14)	0.0364 (15)	-0.0001 (12)	0.0138 (13)	-0.0008 (11)

supplementary materials

C1	0.031 (2)	0.032 (2)	0.030 (2)	0.0007 (16)	0.0078 (16)	0.0034 (15)
C2	0.037 (2)	0.049 (3)	0.036 (2)	-0.0060 (19)	0.0098 (19)	-0.0067 (18)
C3	0.035 (2)	0.054 (3)	0.043 (3)	-0.012 (2)	0.006 (2)	0.001 (2)
C4	0.031 (2)	0.056 (3)	0.041 (2)	-0.003 (2)	0.0123 (18)	0.009 (2)
C5	0.037 (2)	0.048 (3)	0.031 (2)	0.0001 (19)	0.0122 (17)	0.0015 (18)
C6	0.034 (2)	0.0287 (19)	0.0287 (19)	0.0008 (16)	0.0062 (16)	0.0003 (15)
C7	0.045 (3)	0.038 (2)	0.032 (2)	0.0022 (19)	0.0177 (18)	-0.0033 (17)

Geometric parameters (\AA , $^\circ$)

Co1—O2 ⁱ	2.013 (2)	C1—C2	1.406 (6)
Co1—O2	2.013 (2)	C1—C6	1.424 (5)
Co1—O1	2.099 (2)	C2—C3	1.368 (6)
Co1—O1 ⁱ	2.099 (2)	C2—H2	0.9300
Co1—O3 ⁱ	2.149 (3)	C3—C4	1.395 (6)
Co1—O3	2.149 (3)	C3—H3A	0.9300
Br1—C4	1.894 (4)	C4—C5	1.367 (6)
O1—C7	1.225 (5)	C5—C6	1.412 (5)
O2—C1	1.299 (4)	C5—H5	0.9300
O3—H3B	0.8500	C6—C7	1.436 (6)
O3—H3	0.8500	C7—H7	0.9300
O2 ⁱ —Co1—O2	180	O2—C1—C6	123.8 (3)
O2 ⁱ —Co1—O1	92.14 (10)	C2—C1—C6	116.8 (3)
O2—Co1—O1	87.86 (10)	C3—C2—C1	122.1 (4)
O2 ⁱ —Co1—O1 ⁱ	87.86 (10)	C3—C2—H2	118.9
O2—Co1—O1 ⁱ	92.14 (10)	C1—C2—H2	118.9
O1—Co1—O1 ⁱ	180	C2—C3—C4	120.3 (4)
O2 ⁱ —Co1—O3 ⁱ	89.80 (10)	C2—C3—H3A	119.9
O2—Co1—O3 ⁱ	90.20 (10)	C4—C3—H3A	119.9
O1—Co1—O3 ⁱ	86.83 (10)	C5—C4—C3	120.1 (4)
O1 ⁱ —Co1—O3 ⁱ	93.17 (10)	C5—C4—Br1	120.4 (3)
O2 ⁱ —Co1—O3	90.20 (10)	C3—C4—Br1	119.5 (3)
O2—Co1—O3	89.80 (10)	C4—C5—C6	120.3 (4)
O1—Co1—O3	93.17 (10)	C4—C5—H5	119.9
O1 ⁱ —Co1—O3	86.83 (10)	C6—C5—H5	119.9
O3 ⁱ —Co1—O3	180	C5—C6—C1	120.3 (3)
C7—O1—Co1	125.4 (2)	C5—C6—C7	116.2 (3)
C1—O2—Co1	129.1 (2)	C1—C6—C7	123.5 (3)
Co1—O3—H3B	107.9	O1—C7—C6	127.9 (4)
Co1—O3—H3	109.2	O1—C7—H7	116.1
H3B—O3—H3	108.2	C6—C7—H7	116.1
O2—C1—C2	119.4 (3)		

Symmetry codes: (i) $-x, -y, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O3—H3···O1 ⁱⁱ	0.85	2.12	2.842 (4)	142
O3—H3B···O2 ⁱⁱⁱ	0.85	1.93	2.725 (4)	155

Symmetry codes: (ii) $-x, -y-1, -z+1$; (iii) $x, y-1, z$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

