

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***trans*-Dichloridobis(2,2-dimethylpropane-1,3-diamine- κ^2N,N')chromium(III) perchlorate**Jong-Ha Choi,^a Sang Hak Lee^b and Uk Lee^{c*}

^aDepartment of Chemistry, Andong National University, Andong 760-749, Republic of Korea, ^bDepartment of Chemistry, Kyungpook National University, Daegu 702-701, Republic of Korea, and ^cDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong Nam-gu, Busan 608-737, Republic of Korea
Correspondence e-mail: uklee@pknu.ac.kr


Received 8 October 2008; accepted 11 October 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.051; wR factor = 0.146; data-to-parameter ratio = 19.8.

In the title salt, $[\text{CrCl}_2(\text{C}_5\text{H}_{14}\text{N}_2)_2]\text{ClO}_4$, the Cr atom is in a *trans*- CrCl_2N_4 octahedral environment comprising the four N atoms of two chelating 2,2-dimethylpropane-1,3-diamine ligands and two Cl atoms. The two six-membered CrC_3N_2 rings in the cation adopt *anti* chair–chair conformations with respect to each other. The perchlorate anion is disordered over two positions in respect of the Cl and an O atom in a 6:4 ratio. $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the cations and anions into a layer structure.

Related literature

For the synthesis, see: House (1986). For related structures, see: Choi *et al.* (2002, 2007). For the spectroscopic properties, see: Choi (2000); Poon & Pun (1980).



Experimental

Crystal data

$[\text{CrCl}_2(\text{C}_5\text{H}_{14}\text{N}_2)_2]\text{ClO}_4$
 $M_r = 426.71$

Monoclinic, $P2_1/c$
 $a = 6.6373$ (6) Å

$b = 20.767$ (2) Å
 $c = 13.878$ (2) Å
 $\beta = 100.249$ (9)°
 $V = 1882.4$ (4) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 1.05$ mm⁻¹
 $T = 298$ (2) K
 $0.32 \times 0.30 \times 0.25$ mm

Data collection

Stoe Stadi-4 diffractometer
Absorption correction: numerical
(*X-SHAPE*; Stoe & Cie, 1996)
 $T_{\text{min}} = 0.805$, $T_{\text{max}} = 0.942$
4305 measured reflections

4305 independent reflections
3453 reflections with $I > 2\sigma(I)$
3 standard reflections
frequency: 60 min
intensity decay: 2.7%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.146$
 $S = 1.11$
4305 reflections

217 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.54$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.81$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1BN}\cdots\text{O2}^i$	0.90	2.29	3.030 (5)	139
$\text{N2}-\text{H2AN}\cdots\text{O3}^{ii}$	0.90	2.23	3.099 (6)	162
$\text{N2}-\text{H2AN}\cdots\text{O4A}^{ii}$	0.90	2.42	3.183 (6)	143
$\text{N2}-\text{H2BN}\cdots\text{O4B}$	0.90	2.36	3.217 (9)	159
$\text{N3}-\text{H3AN}\cdots\text{O4A}^{ii}$	0.90	2.60	3.482 (7)	168
$\text{N4}-\text{H4BN}\cdots\text{O2}^i$	0.90	2.14	3.030 (5)	172
$\text{N4}-\text{H4AN}\cdots\text{O4A}^{iii}$	0.90	2.54	3.403 (8)	161

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x - 1, y, z$; (iii) $x - 1, -y + \frac{1}{2}, z + \frac{1}{2}$; (iv) $x + 1, y, z$.

Data collection: *STADI-4* (Stoe & Cie, 1996); cell refinement: *STADI-4*; data reduction: *X-RED* (Stoe & Cie, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

This work was supported by funds for 2008 of Andong National University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2499).

References

- Choi, J. H. (2000). *Chem. Phys.* **256**, 29–35.
Choi, J. H., Clegg, W., Nichol, G. S., Lee, S. H., Park, Y. C. & Habibi, M. H. (2007). *Spectrochim. Acta A*, **68**, 796–801.
Choi, J.-H., Suzuki, T. & Kaizaki, S. (2002). *Acta Cryst. C* **58**, m539–m541.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
House, D. A. (1986). *Inorg. Chem.* **25**, 1671–1674.
Poon, C. K. & Pun, K. C. (1980). *Inorg. Chem.* **19**, 568–569.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Stoe & Cie (1996). *STADI-4, X-RED and X-SHAPE*. Stoe & Cie GmbH, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2008). E64, m1429 [doi:10.1107/S1600536808032911]

***trans*-Dichloridobis(2,2-dimethylpropane-1,3-diamine- κ^2N,N')chromium(III) perchlorate**

J.-H. Choi, S. H. Lee and U. Lee

Comment

The $[\text{Cr}(\text{Me}_2\text{tn})_2\text{L}_2]^+$ (Me_2tn =2,2-dimethylpropane-1,3-diamine, L = monodentate) cation can exist as *trans* and *cis* geometric isomers. Especially, there are two possible conformations with respect to the six-membered rings in the *trans* isomer. The carbon atoms of the two chelate rings of the two conformers may be on the same side (*syn* conformer) or on opposite side (*anti* conformer) of the coordination plane (Choi *et al.*, 2002; Choi *et al.*, 2007). The *syn* or *anti* conformational stereochemistry of the six-membered chelate rings can not be readily determined by spectroscopic and physicochemical methods (Poon & Pun, 1980; Choi, 2000). In order to examine the influences of counter anions and packing forces of the crystal on the conformations, we have undertaken the X-ray structural analysis of *trans*- $[\text{Cr}(\text{Me}_2\text{tn})_2\text{Cl}_2]\text{ClO}_4$, (I).

The title complex has approximate C_i symmetry. The two chelate rings in the complex cation are only in anti chair-chair conformation with respect to each other (Fig.1). The Cr—N and Cr—Cl bond length are very similar to those of the *trans*- $[\text{Cr}(\text{Me}_2\text{tn})_2\text{Cl}_2]\text{Cl}$ (Choi *et al.*, 2007). However, the significant difference between these two crystal structures is the orientations with respect to the six-membered chelate rings of two Me_2tn ligands in the same *trans* geometry. The complex is stabilized by the formation of the extensive hydrogen bonds (Table 1).

Experimental

The complex *trans*- $[\text{Cr}(\text{Me}_2\text{tn})_2\text{Cl}_2]\text{ClO}_4$ was prepared according to the literature (House, 1986). The crystalline product deposited with ice-bath cooling was filtered off, and washed with cold 2-propanol and then diethyl ether. Recrystallization of the crude precipitate from 0.5M HCl and 70% HClO_4 solution afforded dark green crystals suitable for X-ray analysis. Anal. Found: C, 28.02; H, 6.50; N, 13.08%. Calc. for $\text{C}_{10}\text{H}_{28}\text{Cl}_3\text{CrN}_4\text{O}_4$: C, 28.15; H, 6.61; N, 13.13%.

Refinement

H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.97 (methylene), 0.96 (methyl) Å, and N—H = 0.90 Å respectively, $\text{Uiso}(\text{H}) = 1.2\text{Ueq}(\text{C} \ \& \ \text{N})$. The Cl and one O atoms in the perchlorate anion are disordered over two positions with site-occupancy factors fixed at 0.60 (for atoms labelled A) and 0.40 (for atoms labelled B) in the final refinement.

Figures




Fig. 1. Perspective view (30% probability level) of title compound.

trans-Dichloridobis(2,2-dimethylpropane-1,3-diamine- κ^2N,N')chromium(III) perchlorate

Crystal data

$[\text{CrCl}_2(\text{C}_5\text{H}_{14}\text{N}_2)_2]\text{ClO}_4$

$M_r = 426.71$

Monoclinic, $P2_1/c$

Hall symbol: $-P_2ybc$

$a = 6.6373$ (6) Å

$b = 20.767$ (2) Å

$c = 13.878$ (2) Å

$\beta = 100.249$ (9)°

$V = 1882.4$ (4) Å³

$Z = 4$

$F_{000} = 892$

$D_x = 1.506$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71069$ Å

Cell parameters from 54 reflections

$\theta = 9.5\text{--}10.4^\circ$

$\mu = 1.05$ mm⁻¹

$T = 298$ (2) K

Block, green

$0.32 \times 0.30 \times 0.25$ mm

Data collection

Stoe Stadi-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298$ (2) K

$\omega/2\text{--}\theta$ scans

Absorption correction: Numerical
(X-SHAPE; Stoe & Cie, 1996)

$T_{\min} = 0.805$, $T_{\max} = 0.942$

4305 measured reflections

4305 independent reflections

3453 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.0000$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 1.8^\circ$

$h = -8 \rightarrow 8$

$k = 0 \rightarrow 26$

$l = 0 \rightarrow 18$

3 standard reflections

every 60 min

intensity decay: 2.7%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.146$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0733P)^2 + 1.4729P]$

$S = 1.11$

4305 reflections

217 parameters

Primary atom site location: structure-invariant direct methods

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.54 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.81 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cr	0.43254 (6)	0.23278 (2)	0.57694 (3)	0.03056 (15)	
Cl1	0.67222 (13)	0.25634 (5)	0.47898 (7)	0.0540 (2)	
Cl2	0.18895 (12)	0.21076 (4)	0.67312 (7)	0.0493 (2)	
Cl3A	0.9541 (3)	0.11988 (10)	0.31249 (14)	0.0471 (4)	0.60
Cl3B	0.8697 (4)	0.11095 (15)	0.3227 (2)	0.0434 (6)	0.40
O4B	0.6845 (14)	0.1083 (5)	0.3541 (7)	0.096 (3)	0.40
O4A	1.1593 (9)	0.1344 (3)	0.3048 (5)	0.0913 (18)	0.60
O1	0.8766 (6)	0.06609 (19)	0.2505 (3)	0.0957 (12)	
O2	0.8510 (7)	0.1748 (2)	0.2793 (3)	0.1124 (15)	
O3	0.9840 (9)	0.1052 (3)	0.4093 (3)	0.140 (2)	
N1	0.6658 (4)	0.20091 (13)	0.6894 (2)	0.0420 (6)	
H1AN	0.7834	0.2010	0.6656	0.050*	
H1BN	0.6795	0.2305	0.7375	0.050*	
N2	0.4181 (4)	0.13895 (13)	0.5210 (2)	0.0466 (7)	
H2AN	0.2998	0.1354	0.4782	0.056*	
H2BN	0.5200	0.1348	0.4865	0.056*	
N3	0.1983 (4)	0.26418 (12)	0.46576 (19)	0.0385 (6)	
H3AN	0.1880	0.2354	0.4167	0.046*	
H3BN	0.0806	0.2622	0.4893	0.046*	
N4	0.4506 (4)	0.32609 (13)	0.6350 (2)	0.0448 (6)	
H4AN	0.3526	0.3297	0.6719	0.054*	
H4BN	0.5717	0.3295	0.6758	0.054*	
C1	0.6476 (5)	0.13679 (16)	0.7353 (2)	0.0444 (7)	
H1A	0.5282	0.1372	0.7667	0.053*	
H1B	0.7668	0.1302	0.7860	0.053*	
C2	0.6295 (5)	0.07995 (15)	0.6641 (3)	0.0411 (7)	

supplementary materials

C3	0.6162 (7)	0.01867 (19)	0.7255 (3)	0.0644 (11)
H3A	0.6049	-0.0184	0.6836	0.077*
H3B	0.4982	0.0212	0.7565	0.077*
H3C	0.7374	0.0151	0.7746	0.077*
C4	0.8169 (6)	0.07509 (19)	0.6154 (3)	0.0569 (9)
H4A	0.8011	0.0393	0.5708	0.068*
H4B	0.9368	0.0688	0.6645	0.068*
H4C	0.8311	0.1141	0.5801	0.068*
C5	0.4308 (5)	0.08283 (16)	0.5890 (3)	0.0495 (8)
H5A	0.4181	0.0435	0.5507	0.059*
H5B	0.3161	0.0846	0.6235	0.059*
C6	0.2096 (5)	0.32895 (16)	0.4223 (2)	0.0441 (7)
H6A	0.0874	0.3356	0.3735	0.053*
H6B	0.3261	0.3300	0.3889	0.053*
C7	0.2293 (5)	0.38464 (15)	0.4953 (2)	0.0410 (7)
C8	0.2337 (8)	0.4469 (2)	0.4358 (4)	0.0715 (12)
H8A	0.2455	0.4833	0.4791	0.086*
H8B	0.1095	0.4503	0.3886	0.086*
H8C	0.3488	0.4459	0.4026	0.086*
C9	0.4311 (5)	0.38267 (15)	0.5681 (3)	0.0469 (8)
H9A	0.5435	0.3821	0.5320	0.056*
H9B	0.4433	0.4217	0.6071	0.056*
C10	0.0472 (6)	0.38715 (19)	0.5480 (3)	0.0571 (9)
H10A	0.0640	0.4224	0.5935	0.069*
H10B	0.0388	0.3475	0.5827	0.069*
H10C	-0.0762	0.3932	0.5010	0.069*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cr	0.0243 (2)	0.0293 (2)	0.0375 (3)	0.00052 (16)	0.00378 (17)	-0.00218 (18)
Cl1	0.0375 (4)	0.0574 (5)	0.0726 (6)	0.0075 (4)	0.0244 (4)	0.0073 (4)
Cl2	0.0377 (4)	0.0561 (5)	0.0578 (5)	0.0062 (3)	0.0185 (4)	0.0123 (4)
Cl3A	0.0561 (12)	0.0449 (9)	0.0375 (8)	0.0080 (9)	0.0011 (9)	-0.0001 (6)
Cl3B	0.0381 (13)	0.0507 (14)	0.0405 (12)	0.0016 (11)	0.0046 (11)	0.0027 (9)
O4B	0.077 (5)	0.122 (8)	0.096 (6)	-0.025 (5)	0.034 (5)	-0.052 (6)
O4A	0.065 (3)	0.100 (4)	0.099 (4)	-0.011 (3)	-0.013 (3)	0.000 (4)
O1	0.110 (3)	0.091 (3)	0.092 (2)	-0.029 (2)	0.034 (2)	-0.038 (2)
O2	0.126 (3)	0.077 (2)	0.122 (3)	0.025 (2)	-0.010 (3)	0.024 (2)
O3	0.167 (5)	0.178 (5)	0.062 (2)	0.004 (4)	-0.019 (3)	0.017 (3)
N1	0.0319 (12)	0.0373 (13)	0.0526 (16)	0.0014 (10)	-0.0036 (11)	-0.0012 (12)
N2	0.0489 (15)	0.0369 (14)	0.0502 (16)	0.0038 (12)	-0.0017 (12)	-0.0068 (12)
N3	0.0332 (12)	0.0391 (14)	0.0420 (14)	0.0029 (10)	0.0031 (10)	-0.0021 (11)
N4	0.0485 (15)	0.0358 (13)	0.0460 (15)	0.0014 (11)	-0.0029 (12)	-0.0048 (12)
C1	0.0439 (17)	0.0408 (17)	0.0467 (17)	0.0064 (13)	0.0035 (14)	0.0064 (14)
C2	0.0373 (15)	0.0323 (15)	0.0554 (19)	0.0035 (12)	0.0126 (13)	0.0059 (13)
C3	0.070 (3)	0.044 (2)	0.084 (3)	0.0058 (18)	0.024 (2)	0.0195 (19)
C4	0.0474 (19)	0.053 (2)	0.076 (3)	0.0068 (16)	0.0259 (18)	0.0017 (19)

C5	0.0434 (17)	0.0322 (16)	0.071 (2)	-0.0017 (13)	0.0040 (16)	-0.0052 (15)
C6	0.0467 (17)	0.0487 (18)	0.0371 (16)	0.0131 (14)	0.0080 (13)	0.0041 (14)
C7	0.0417 (16)	0.0352 (15)	0.0479 (17)	0.0068 (12)	0.0133 (13)	0.0069 (13)
C8	0.083 (3)	0.045 (2)	0.086 (3)	0.007 (2)	0.015 (2)	0.022 (2)
C9	0.0448 (18)	0.0321 (15)	0.064 (2)	-0.0009 (13)	0.0094 (16)	-0.0003 (14)
C10	0.049 (2)	0.052 (2)	0.077 (3)	0.0090 (16)	0.0259 (19)	-0.0037 (18)

Geometric parameters (Å, °)

Cr—N3	2.090 (3)	C1—H1A	0.9700
Cr—N2	2.093 (3)	C1—H1B	0.9700
Cr—N4	2.094 (3)	C2—C4	1.522 (4)
Cr—N1	2.100 (3)	C2—C5	1.530 (5)
Cr—Cl2	2.3179 (9)	C2—C3	1.543 (5)
Cr—Cl1	2.3212 (9)	C3—H3A	0.9600
Cl3A—Cl3B	0.629 (2)	C3—H3B	0.9600
Cl3A—O3	1.357 (4)	C3—H3C	0.9600
Cl3A—O2	1.368 (4)	C4—H4A	0.9600
Cl3A—O4A	1.418 (6)	C4—H4B	0.9600
Cl3A—O1	1.447 (4)	C4—H4C	0.9600
Cl3B—O3	1.308 (5)	C5—H5A	0.9700
Cl3B—O1	1.374 (5)	C5—H5B	0.9700
Cl3B—O4B	1.377 (9)	C6—C7	1.528 (5)
Cl3B—O2	1.453 (5)	C6—H6A	0.9700
N1—C1	1.490 (4)	C6—H6B	0.9700
N1—H1AN	0.9000	C7—C10	1.520 (5)
N1—H1BN	0.9000	C7—C9	1.528 (5)
N2—C5	1.492 (4)	C7—C8	1.537 (5)
N2—H2AN	0.9000	C8—H8A	0.9600
N2—H2BN	0.9000	C8—H8B	0.9600
N3—C6	1.481 (4)	C8—H8C	0.9600
N3—H3AN	0.9000	C9—H9A	0.9700
N3—H3BN	0.9000	C9—H9B	0.9700
N4—C9	1.488 (4)	C10—H10A	0.9600
N4—H4AN	0.9000	C10—H10B	0.9600
N4—H4BN	0.9000	C10—H10C	0.9600
C1—C2	1.530 (5)		
N3—Cr—N2	92.17 (10)	N1—C1—C2	114.6 (3)
N3—Cr—N4	88.82 (10)	N1—C1—H1A	108.6
N2—Cr—N4	179.01 (11)	C2—C1—H1A	108.6
N3—Cr—N1	179.46 (11)	N1—C1—H1B	108.6
N2—Cr—N1	87.78 (11)	C2—C1—H1B	108.6
N4—Cr—N1	91.24 (11)	H1A—C1—H1B	107.6
N3—Cr—Cl2	89.04 (8)	C4—C2—C5	111.9 (3)
N2—Cr—Cl2	92.25 (9)	C4—C2—C1	111.2 (3)
N4—Cr—Cl2	87.65 (9)	C5—C2—C1	111.7 (3)
N1—Cr—Cl2	90.42 (8)	C4—C2—C3	108.8 (3)
N3—Cr—Cl1	89.95 (8)	C5—C2—C3	106.4 (3)
N2—Cr—Cl1	88.28 (9)	C1—C2—C3	106.6 (3)

supplementary materials

N4—Cr—Cl1	91.84 (9)	C2—C3—H3A	109.5
N1—Cr—Cl1	90.59 (8)	C2—C3—H3B	109.5
Cl2—Cr—Cl1	178.87 (4)	H3A—C3—H3B	109.5
O3—Cl3A—O2	119.8 (4)	C2—C3—H3C	109.5
Cl3B—Cl3A—O4A	170.0 (5)	H3A—C3—H3C	109.5
O3—Cl3A—O4A	98.6 (4)	H3B—C3—H3C	109.5
O2—Cl3A—O4A	103.2 (4)	C2—C4—H4A	109.5
O3—Cl3A—O1	112.9 (3)	C2—C4—H4B	109.5
O2—Cl3A—O1	109.9 (3)	H4A—C4—H4B	109.5
O4A—Cl3A—O1	111.3 (3)	C2—C4—H4C	109.5
O3—Cl3A—O4B	70.5 (4)	H4A—C4—H4C	109.5
O2—Cl3A—O4B	77.1 (4)	H4B—C4—H4C	109.5
O4A—Cl3A—O4B	166.7 (4)	N2—C5—C2	114.0 (3)
O1—Cl3A—O4B	80.6 (3)	N2—C5—H5A	108.8
O3—Cl3B—O1	121.3 (4)	C2—C5—H5A	108.8
O3—Cl3B—O4B	96.3 (5)	N2—C5—H5B	108.8
O1—Cl3B—O4B	110.6 (4)	C2—C5—H5B	108.8
O3—Cl3B—O2	117.1 (4)	H5A—C5—H5B	107.7
O1—Cl3B—O2	109.2 (3)	N3—C6—C7	115.0 (3)
O4B—Cl3B—O2	98.8 (6)	N3—C6—H6A	108.5
O3—Cl3B—O4A	74.2 (3)	C7—C6—H6A	108.5
O1—Cl3B—O4A	85.3 (3)	N3—C6—H6B	108.5
O4B—Cl3B—O4A	164.1 (5)	C7—C6—H6B	108.5
O2—Cl3B—O4A	75.2 (3)	H6A—C6—H6B	107.5
C1—N1—Cr	119.62 (19)	C10—C7—C6	111.3 (3)
C1—N1—H1AN	107.4	C10—C7—C9	111.2 (3)
Cr—N1—H1AN	107.4	C6—C7—C9	112.3 (3)
C1—N1—H1BN	107.4	C10—C7—C8	108.8 (3)
Cr—N1—H1BN	107.4	C6—C7—C8	106.7 (3)
H1AN—N1—H1BN	106.9	C9—C7—C8	106.2 (3)
C5—N2—Cr	119.9 (2)	C7—C8—H8A	109.5
C5—N2—H2AN	107.3	C7—C8—H8B	109.5
Cr—N2—H2AN	107.3	H8A—C8—H8B	109.5
C5—N2—H2BN	107.3	C7—C8—H8C	109.5
Cr—N2—H2BN	107.3	H8A—C8—H8C	109.5
H2AN—N2—H2BN	106.9	H8B—C8—H8C	109.5
C6—N3—Cr	119.9 (2)	N4—C9—C7	113.7 (3)
C6—N3—H3AN	107.3	N4—C9—H9A	108.8
Cr—N3—H3AN	107.3	C7—C9—H9A	108.8
C6—N3—H3BN	107.3	N4—C9—H9B	108.8
Cr—N3—H3BN	107.3	C7—C9—H9B	108.8
H3AN—N3—H3BN	106.9	H9A—C9—H9B	107.7
C9—N4—Cr	119.9 (2)	C7—C10—H10A	109.5
C9—N4—H4AN	107.4	C7—C10—H10B	109.5
Cr—N4—H4AN	107.4	H10A—C10—H10B	109.5
C9—N4—H4BN	107.4	C7—C10—H10C	109.5
Cr—N4—H4BN	107.4	H10A—C10—H10C	109.5
H4AN—N4—H4BN	106.9	H10B—C10—H10C	109.5
O2—Cl3A—Cl3B—O3	-123.6 (4)	O1—Cl3A—O2—Cl3B	67.3 (4)

O4A—C13A—C13B—O3	20 (3)	O4B—C13A—O2—C13B	-7.5 (5)
O1—C13A—C13B—O3	123.2 (4)	O3—C13B—O2—C13A	67.5 (5)
O4B—C13A—C13B—O3	-80 (3)	O1—C13B—O2—C13A	-75.3 (4)
O3—C13A—C13B—O1	-123.2 (4)	O4B—C13B—O2—C13A	169.2 (6)
O2—C13A—C13B—O1	113.2 (3)	O4A—C13B—O2—C13A	4.3 (4)
O4A—C13A—C13B—O1	-103 (3)	O1—C13B—O3—C13A	76.7 (5)
O4B—C13A—C13B—O1	157 (3)	O4B—C13B—O3—C13A	-164.5 (7)
O3—C13A—C13B—O4B	80 (3)	O2—C13B—O3—C13A	-61.3 (5)
O2—C13A—C13B—O4B	-43 (3)	O4A—C13B—O3—C13A	2.4 (4)
O4A—C13A—C13B—O4B	100 (4)	O2—C13A—O3—C13B	72.8 (5)
O1—C13A—C13B—O4B	-157 (3)	O4A—C13A—O3—C13B	-176.6 (6)
O3—C13A—C13B—O2	123.6 (4)	O1—C13A—O3—C13B	-59.0 (5)
O4A—C13A—C13B—O2	143 (3)	O4B—C13A—O3—C13B	11.2 (5)
O1—C13A—C13B—O2	-113.2 (3)	N2—Cr—N1—C1	-41.0 (2)
O4B—C13A—C13B—O2	43 (3)	N4—Cr—N1—C1	138.9 (2)
O3—C13A—C13B—O4A	-20 (3)	Cl2—Cr—N1—C1	51.3 (2)
O2—C13A—C13B—O4A	-143 (3)	Cl1—Cr—N1—C1	-129.2 (2)
O1—C13A—C13B—O4A	103 (3)	N3—Cr—N2—C5	-137.9 (2)
O4B—C13A—C13B—O4A	-100 (4)	N1—Cr—N2—C5	41.6 (2)
O3—C13B—O4B—C13A	78 (3)	Cl2—Cr—N2—C5	-48.8 (2)
O1—C13B—O4B—C13A	-155 (3)	Cl1—Cr—N2—C5	132.2 (2)
O2—C13B—O4B—C13A	-41 (2)	N2—Cr—N3—C6	-141.5 (2)
O4A—C13B—O4B—C13A	25.7 (16)	N4—Cr—N3—C6	38.6 (2)
O3—C13A—O4B—C13B	-96 (3)	Cl2—Cr—N3—C6	126.3 (2)
O2—C13A—O4B—C13B	135 (3)	Cl1—Cr—N3—C6	-53.2 (2)
O4A—C13A—O4B—C13B	-132 (3)	N3—Cr—N4—C9	-39.7 (2)
O1—C13A—O4B—C13B	22 (2)	N1—Cr—N4—C9	140.8 (2)
O3—C13A—O4A—C13B	19 (3)	Cl2—Cr—N4—C9	-128.8 (2)
O2—C13A—O4A—C13B	142 (3)	Cl1—Cr—N4—C9	50.2 (2)
O1—C13A—O4A—C13B	-100 (3)	Cr—N1—C1—C2	59.1 (3)
O4B—C13A—O4A—C13B	53 (3)	N1—C1—C2—C4	60.6 (4)
O3—C13B—O4A—C13A	-160 (3)	N1—C1—C2—C5	-65.2 (4)
O1—C13B—O4A—C13A	76 (3)	N1—C1—C2—C3	179.0 (3)
O4B—C13B—O4A—C13A	-105 (4)	Cr—N2—C5—C2	-60.0 (4)
O2—C13B—O4A—C13A	-35 (3)	C4—C2—C5—N2	-60.1 (4)
O3—C13B—O1—C13A	-75.2 (5)	C1—C2—C5—N2	65.2 (4)
O4B—C13B—O1—C13A	173.4 (8)	C3—C2—C5—N2	-178.8 (3)
O2—C13B—O1—C13A	65.8 (4)	Cr—N3—C6—C7	-56.9 (3)
O4A—C13B—O1—C13A	-6.8 (4)	N3—C6—C7—C10	-60.2 (4)
O3—C13A—O1—C13B	59.8 (5)	N3—C6—C7—C9	65.2 (4)
O2—C13A—O1—C13B	-76.8 (5)	N3—C6—C7—C8	-178.8 (3)
O4A—C13A—O1—C13B	169.6 (6)	Cr—N4—C9—C7	58.7 (3)
O4B—C13A—O1—C13B	-4.3 (5)	C10—C7—C9—N4	59.8 (4)
O3—C13A—O2—C13B	-65.8 (5)	C6—C7—C9—N4	-65.7 (4)
O4A—C13A—O2—C13B	-173.9 (6)	C8—C7—C9—N4	178.0 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
---------------	-------	-------------	-------------	---------------

supplementary materials

N1—H1BN...O2 ⁱ	0.90	2.29	3.030 (5)	139
N2—H2AN...O3 ⁱⁱ	0.90	2.23	3.099 (6)	162
N2—H2AN...O4A ⁱⁱ	0.90	2.42	3.183 (6)	143
N2—H2BN...O4B	0.90	2.36	3.217 (9)	159
N3—H3AN...O4A ⁱⁱ	0.90	2.60	3.482 (7)	168
N4—H4BN...O2 ⁱ	0.90	2.14	3.030 (5)	172
N4—H4AN...O4A ⁱⁱⁱ	0.90	2.54	3.403 (8)	161
N1—H1AN...Cl2 ^{iv}	0.90	2.68	3.525 (3)	156
N3—H3BN...Cl1 ⁱⁱ	0.90	2.69	3.533 (3)	156

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $x-1, y, z$; (iii) $x-1, -y+1/2, z+1/2$; (iv) $x+1, y, z$.

Fig. 1

