

1,2-Diazinium hydrogen chloranilate

Kazuma Gotoh and Hiroyuki Ishida*

Department of Chemistry, Faculty of Science, Okayama University, Okayama 700-8530, Japan

Correspondence e-mail: ishidah@cc.okayama-u.ac.jp


Received 30 September 2008; accepted 5 October 2008

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.030; wR factor = 0.082; data-to-parameter ratio = 18.4.

In the crystal structure of the title compound, $\text{C}_4\text{H}_5\text{N}_2^+\cdot\text{C}_6\text{HCl}_2\text{O}_4^-$, there are three crystallographically independent 1,2-diazinium cations and hydrogen chloranilate anions. The anions are held together by pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds to form two types of dimers, one of which is centrosymmetric. The 1,2-diazinium cations are linked on both sides of each dimer via bifurcated $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds to give two kinds of 2–2 cation–anion associations. The 2–2 associations are linked by intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds, forming a molecular tape along the [230] direction. The tapes are further connected by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For general background, see: Gotoh *et al.* (2007). For related compounds, see: Gotoh *et al.* (2008); Ishida & Kashino (1999).



Experimental

Crystal data



$M_r = 289.07$

Monoclinic, $P2_1/c$

$a = 25.6849(10)\text{ \AA}$

$b = 7.0516(2)\text{ \AA}$

$c = 18.1388(6)\text{ \AA}$

$\beta = 97.5822(13)^\circ$

$V = 3256.56(18)\text{ \AA}^3$

$Z = 12$

Mo $K\alpha$ radiation

$\mu = 0.61\text{ mm}^{-1}$

$T = 173(1)\text{ K}$

$0.40 \times 0.22 \times 0.12\text{ mm}$

Data collection

Rigaku R-AXIS RAPIDII

diffractometer

Absorption correction: numerical
(*ABSCOR*; Higashi, 1999)

$T_{\min} = 0.855$, $T_{\max} = 0.930$

42942 measured reflections

9398 independent reflections

7981 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.082$

$S = 1.04$

9398 reflections

511 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.47\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O2	0.889 (17)	1.753 (17)	2.6138 (13)	162.2 (17)
N1—H1 \cdots O3	0.889 (17)	2.578 (18)	3.1583 (14)	123.7 (14)
N3—H3 \cdots O6	0.886 (18)	1.811 (17)	2.6621 (13)	160.4 (17)
N3—H3 \cdots O7	0.886 (18)	2.431 (18)	3.0066 (14)	123.0 (14)
O4—H4 \cdots O1	0.79 (2)	2.26 (2)	2.6619 (13)	113 (2)
O4—H4 \cdots O5	0.79 (2)	1.94 (2)	2.6723 (12)	155 (2)
N5—H5 \cdots O10	0.890 (18)	1.881 (18)	2.7450 (14)	163.2 (16)
N5—H5 \cdots O11	0.890 (18)	2.410 (17)	2.9659 (13)	120.7 (14)
O8—H8 \cdots O1	0.80 (2)	1.86 (2)	2.5930 (12)	152 (2)
O8—H8 \cdots O5	0.80 (2)	2.25 (2)	2.6582 (13)	112.7 (18)
O12—H12 \cdots O9	0.79 (2)	2.23 (2)	2.6595 (13)	115.1 (18)
O12—H12 \cdots O9 ⁱ	0.79 (2)	2.02 (2)	2.6801 (12)	142 (2)
C20—H20 \cdots O2 ⁱⁱ	0.95	2.57	3.1097 (15)	116
C21—H21 \cdots O3 ⁱⁱⁱ	0.95	2.36	3.2754 (15)	162
C22—H22 \cdots O3	0.95	2.60	3.1913 (15)	121
C25—H25 \cdots O10	0.95	2.47	3.4056 (15)	169
C26—H26 \cdots O7	0.95	2.44	3.0363 (15)	121
C26—H26 \cdots N6	0.95	2.51	3.2665 (17)	137
C27—H27 \cdots O7	0.95	2.37	3.2765 (15)	159
C28—H28 \cdots O6 ^{iv}	0.95	2.47	3.1091 (15)	124
C30—H30 \cdots O11	0.95	2.34	2.9551 (15)	122

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 2, -y + 3, -z + 1$; (iv) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *PROCESS-AUTO* (Rigaku/MSC, 2004); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure* and *PLATON* (Spek, 2003).

This work was supported by a Grant-in-Aid for Scientific Research (C) (No. 19550018) from the Japan Society for the Promotion of Science.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2158).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gotoh, K., Asaji, T. & Ishida, H. (2007). *Acta Cryst.* **C63**, o17–o20.
- Gotoh, K., Asaji, T. & Ishida, H. (2008). *Acta Cryst.* **C64**, o550–o553.
- Higashi, T. (1999). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Ishida, H. & Kashino, S. (1999). *Acta Cryst.* **C55**, 1149–1152.
- Rigaku/MSC (2004). *PROCESS-AUTO* and *CrystalStructure*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o2095 [doi:10.1107/S1600536808031978]

1,2-Diazinium hydrogen chloranilate

K. Gotoh and H. Ishida

Comment

The title compound, (I), was prepared in order to extend our study on $D-\text{H}\cdots A$ hydrogen bonding ($D = \text{N}, \text{O}, \text{or C}; A = \text{N}, \text{O} \text{ or Cl}$) in amine–chloranilic acid (systematic name: 2,5-dichloro-3,6-dihydroxy-1,4-benzoquinone) systems (Gotoh *et al.*, 2007). The crystal structure of 1,2-diazine (pyridazine)–chloranilic acid (2/1) was already reported (Ishida & Kashino, 1999; Gotoh *et al.*, 2008).

The asymmetric unit in (I) contains three 1,2-diazinium cations and three hydrogen chloranilate anions. Hydrogen chloranilate anions are held together by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds (Table 1) to form two crystallographically independent dimers. The 1,2-diazinium cations are linked on both sides of each dimer *via* bifurcated $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds to give centro- and acentrosymmetric 2:2 salts of 1,2-diazine and chloranilic acid (Fig. 1). The $\text{N}\cdots\text{O}$ distances [2.6138 (13)–3.1583 (14) Å] in the hydrogen bonds are longer than those [2.582 (3) Å at 299 K (Ishida & Kashino, 1999) and 2.5549 (12) Å at 110 K (Gotoh *et al.*, 2008)] in 1,2-diazine–chloranilic acid (2/1) and the H atoms are located at the N atom sites. The dihedral angles between C1–C6 and C7–C12 rings, C1–C6 and N1/N2/C19–C22 rings, C7–C12 and N3/N4/C23–C26 rings, and C13–C18 and N5/N6/C27–C30 rings are 3.40 (5), 11.83 (5), 4.88 (6) and 11.41 (5)°, respectively. The 2:2 units are linked by intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds, forming a molecular tape along the [230] direction. The tapes are further connected by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network (Table 1 and Fig. 2).

Experimental

Single crystals were obtained by slow evaporation from a methanol solution (30 ml) of chloranilic acid (400 mg) and 1,2-diazine (155 mg).

Refinement

C-bound H atoms were positioned geometrically ($\text{C}-\text{H} = 0.95$ Å) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms in the $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds were located in a difference Fourier map and refined isotropically (refined distances given in Table 1).

Figures




Fig. 1. The structure of the 2:2 units of (I). The dashed lines indicate $\text{O}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds (symmetry code as Table 1). Displacement ellipsoids of non-H atoms are drawn at the 50% probability level.



Fig. 2. A partial packing diagram of (I), viewed down the b axis. The dashed lines indicate $\text{O}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds.

supplementary materials

1,2-Diazinium hydrogen chloranilate

Crystal data

$C_4H_5N_2^+ \cdot C_6HCl_2O_4^-$	$F_{000} = 1752.00$
$M_r = 289.07$	$D_x = 1.769 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71075 \text{ \AA}$
$a = 25.6849 (10) \text{ \AA}$	Cell parameters from 36292 reflections
$b = 7.0516 (2) \text{ \AA}$	$\theta = 3.0\text{--}30.0^\circ$
$c = 18.1388 (6) \text{ \AA}$	$\mu = 0.61 \text{ mm}^{-1}$
$\beta = 97.5822 (13)^\circ$	$T = 173 (1) \text{ K}$
$V = 3256.56 (18) \text{ \AA}^3$	Prism, brown
$Z = 12$	$0.40 \times 0.22 \times 0.12 \text{ mm}$

Data collection

Rigaku R-AXIS RAPIDII diffractometer	7981 reflections with $I > 2\sigma(I)$
Detector resolution: 10.00 pixels mm^{-1}	$R_{\text{int}} = 0.025$
ω scans	$\theta_{\text{max}} = 30.0^\circ$
Absorption correction: numerical (ABSCOR; Higashi, 1999)	$h = -36 \rightarrow 36$
$T_{\text{min}} = 0.855$, $T_{\text{max}} = 0.930$	$k = -9 \rightarrow 8$
42942 measured reflections	$l = -24 \rightarrow 25$
9398 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0465P)^2 + 0.8542P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} = 0.002$
9398 reflections	$\Delta\rho_{\text{max}} = 0.47 \text{ e \AA}^{-3}$
511 parameters	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.788813 (11)	1.00777 (4)	0.695195 (15)	0.02287 (7)
Cl2	0.813873 (11)	1.33053 (4)	0.375637 (15)	0.01939 (6)
Cl3	0.546309 (11)	0.98926 (4)	0.291828 (15)	0.02164 (7)
Cl4	0.529070 (10)	0.68298 (4)	0.616552 (15)	0.01938 (6)
Cl5	0.134170 (11)	0.31600 (4)	0.406398 (15)	0.02014 (6)
Cl6	0.126530 (11)	0.00278 (4)	0.732528 (15)	0.02182 (7)
O1	0.70886 (3)	1.03911 (14)	0.56092 (5)	0.02672 (19)
O2	0.88525 (3)	1.17619 (12)	0.64597 (5)	0.02155 (17)
O3	0.89647 (3)	1.29114 (12)	0.50763 (5)	0.02036 (16)
O4	0.71996 (3)	1.15850 (13)	0.42465 (5)	0.02316 (18)
O5	0.62655 (3)	0.99280 (14)	0.42628 (5)	0.0258 (2)
O6	0.45198 (3)	0.80783 (12)	0.34672 (5)	0.02140 (17)
O7	0.44599 (3)	0.69004 (12)	0.48544 (5)	0.02138 (17)
O8	0.61970 (3)	0.86803 (14)	0.56295 (5)	0.02294 (18)
O9	0.04325 (3)	0.14786 (13)	0.47288 (5)	0.02485 (19)
O10	0.22343 (3)	0.29297 (12)	0.53149 (5)	0.02021 (17)
O11	0.21688 (3)	0.17839 (12)	0.67058 (5)	0.02290 (18)
O12	0.04124 (3)	0.00464 (13)	0.60811 (5)	0.02206 (18)
N1	0.97488 (4)	1.35579 (15)	0.65484 (6)	0.0219 (2)
N2	0.99547 (4)	1.31774 (15)	0.72462 (6)	0.0247 (2)
N3	0.36001 (4)	0.64298 (15)	0.35724 (6)	0.0212 (2)
N4	0.33565 (4)	0.66986 (16)	0.28834 (6)	0.0265 (2)
N5	0.31069 (4)	0.37842 (15)	0.63023 (6)	0.02006 (19)
N6	0.34283 (4)	0.45730 (16)	0.58711 (6)	0.0239 (2)
C1	0.75362 (4)	1.09400 (16)	0.55266 (6)	0.0170 (2)
C2	0.79706 (4)	1.09715 (16)	0.60871 (6)	0.01556 (19)
C3	0.84590 (4)	1.16825 (15)	0.59736 (6)	0.0152 (2)
C4	0.85282 (4)	1.24116 (15)	0.51915 (6)	0.01446 (19)
C5	0.80717 (4)	1.24251 (15)	0.46242 (6)	0.01550 (19)
C6	0.76101 (4)	1.16835 (15)	0.47678 (6)	0.0164 (2)
C7	0.58355 (4)	0.92184 (16)	0.43505 (6)	0.0165 (2)
C8	0.54005 (4)	0.90299 (16)	0.37928 (6)	0.0157 (2)
C9	0.49245 (4)	0.82418 (15)	0.39314 (6)	0.0155 (2)

supplementary materials

C10	0.48814 (4)	0.75267 (15)	0.47232 (6)	0.01508 (19)
C11	0.53375 (4)	0.76601 (15)	0.52856 (6)	0.01517 (19)
C12	0.57836 (4)	0.84830 (16)	0.51193 (6)	0.0162 (2)
C13	0.08582 (4)	0.15642 (15)	0.51475 (6)	0.0164 (2)
C14	0.13318 (4)	0.23100 (15)	0.49570 (6)	0.01558 (19)
C15	0.17974 (4)	0.23689 (15)	0.54582 (6)	0.01537 (19)
C16	0.17708 (4)	0.16726 (15)	0.62576 (6)	0.0154 (2)
C17	0.12835 (4)	0.08899 (16)	0.64439 (6)	0.0159 (2)
C18	0.08570 (4)	0.07922 (15)	0.59219 (6)	0.0156 (2)
C19	1.04186 (5)	1.39362 (18)	0.74659 (7)	0.0234 (2)
H19	1.0576	1.3710	0.7962	0.028*
C20	1.06920 (5)	1.50533 (17)	0.70098 (7)	0.0232 (2)
H20	1.1027	1.5564	0.7190	0.028*
C21	1.04646 (5)	1.53937 (18)	0.62952 (7)	0.0231 (2)
H21	1.0637	1.6133	0.5962	0.028*
C22	0.99713 (5)	1.46098 (17)	0.60787 (7)	0.0228 (2)
H22	0.9795	1.4834	0.5592	0.027*
C23	0.28768 (5)	0.59892 (19)	0.27484 (7)	0.0279 (3)
H23	0.2688	0.6165	0.2266	0.034*
C24	0.26300 (5)	0.49954 (18)	0.32732 (8)	0.0250 (3)
H24	0.2286	0.4497	0.3148	0.030*
C25	0.28964 (5)	0.47601 (18)	0.39706 (7)	0.0252 (2)
H25	0.2746	0.4101	0.4347	0.030*
C26	0.33997 (5)	0.55323 (18)	0.41044 (7)	0.0247 (2)
H26	0.3599	0.5409	0.4582	0.030*
C27	0.38814 (5)	0.51986 (19)	0.62180 (7)	0.0252 (3)
H27	0.4118	0.5787	0.5928	0.030*
C28	0.40324 (5)	0.50458 (18)	0.69840 (7)	0.0242 (2)
H28	0.4363	0.5508	0.7209	0.029*
C29	0.36891 (5)	0.42094 (18)	0.74013 (7)	0.0242 (2)
H29	0.3774	0.4059	0.7924	0.029*
C30	0.32121 (5)	0.35891 (18)	0.70316 (7)	0.0238 (2)
H30	0.2961	0.3023	0.7303	0.029*
H1	0.9443 (7)	1.298 (2)	0.6418 (10)	0.040 (5)*
H4	0.6953 (9)	1.112 (3)	0.4388 (13)	0.070 (7)*
H8	0.6436 (8)	0.921 (3)	0.5477 (12)	0.064 (7)*
H3	0.3919 (7)	0.693 (2)	0.3649 (10)	0.037 (5)*
H5	0.2801 (7)	0.340 (2)	0.6060 (10)	0.040 (5)*
H12	0.0205 (8)	-0.001 (3)	0.5722 (11)	0.052 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.02091 (13)	0.03338 (16)	0.01460 (12)	-0.00592 (10)	0.00337 (9)	0.00278 (11)
Cl2	0.02201 (13)	0.02140 (13)	0.01473 (12)	-0.00058 (9)	0.00235 (9)	0.00273 (10)
Cl3	0.02047 (13)	0.03130 (15)	0.01329 (12)	-0.00316 (10)	0.00276 (9)	0.00268 (10)
Cl4	0.02048 (13)	0.02307 (13)	0.01446 (12)	-0.00125 (9)	0.00187 (9)	0.00367 (10)
Cl5	0.02339 (13)	0.02148 (13)	0.01490 (12)	-0.00022 (10)	0.00008 (9)	0.00332 (10)

Cl6	0.01993 (13)	0.03180 (15)	0.01370 (12)	-0.00260 (10)	0.00209 (9)	0.00197 (10)
O1	0.0137 (4)	0.0443 (5)	0.0221 (4)	-0.0085 (3)	0.0020 (3)	0.0002 (4)
O2	0.0153 (4)	0.0311 (4)	0.0170 (4)	-0.0058 (3)	-0.0026 (3)	0.0029 (3)
O3	0.0155 (4)	0.0264 (4)	0.0195 (4)	-0.0057 (3)	0.0034 (3)	-0.0006 (3)
O4	0.0137 (4)	0.0360 (5)	0.0185 (4)	-0.0030 (3)	-0.0026 (3)	0.0037 (4)
O5	0.0148 (4)	0.0416 (5)	0.0206 (4)	-0.0087 (3)	0.0009 (3)	0.0036 (4)
O6	0.0156 (4)	0.0296 (4)	0.0177 (4)	-0.0056 (3)	-0.0028 (3)	0.0023 (3)
O7	0.0163 (4)	0.0282 (4)	0.0195 (4)	-0.0075 (3)	0.0020 (3)	0.0000 (3)
O8	0.0135 (4)	0.0372 (5)	0.0169 (4)	-0.0064 (3)	-0.0024 (3)	0.0047 (4)
O9	0.0161 (4)	0.0313 (5)	0.0248 (4)	-0.0035 (3)	-0.0062 (3)	0.0067 (4)
O10	0.0159 (4)	0.0265 (4)	0.0179 (4)	-0.0047 (3)	0.0014 (3)	0.0001 (3)
O11	0.0167 (4)	0.0305 (4)	0.0198 (4)	-0.0041 (3)	-0.0036 (3)	0.0035 (3)
O12	0.0138 (4)	0.0323 (5)	0.0196 (4)	-0.0045 (3)	0.0003 (3)	0.0014 (3)
N1	0.0144 (4)	0.0246 (5)	0.0262 (5)	-0.0023 (4)	0.0010 (4)	-0.0009 (4)
N2	0.0249 (5)	0.0260 (5)	0.0233 (5)	-0.0044 (4)	0.0041 (4)	0.0014 (4)
N3	0.0153 (4)	0.0247 (5)	0.0225 (5)	-0.0032 (4)	-0.0015 (4)	-0.0013 (4)
N4	0.0262 (5)	0.0306 (6)	0.0216 (5)	-0.0059 (4)	-0.0017 (4)	0.0021 (4)
N5	0.0151 (4)	0.0238 (5)	0.0207 (5)	-0.0023 (4)	0.0001 (3)	-0.0016 (4)
N6	0.0214 (5)	0.0310 (5)	0.0189 (5)	-0.0043 (4)	0.0017 (4)	-0.0003 (4)
C1	0.0141 (5)	0.0205 (5)	0.0164 (5)	-0.0015 (4)	0.0018 (4)	-0.0019 (4)
C2	0.0143 (5)	0.0204 (5)	0.0120 (4)	-0.0022 (4)	0.0019 (3)	0.0002 (4)
C3	0.0134 (4)	0.0169 (5)	0.0152 (5)	-0.0013 (4)	0.0012 (4)	-0.0013 (4)
C4	0.0145 (4)	0.0143 (5)	0.0144 (5)	-0.0014 (4)	0.0016 (4)	-0.0022 (4)
C5	0.0159 (5)	0.0170 (5)	0.0133 (4)	0.0005 (4)	0.0009 (4)	0.0004 (4)
C6	0.0142 (5)	0.0188 (5)	0.0155 (5)	0.0014 (4)	-0.0004 (4)	-0.0012 (4)
C7	0.0138 (5)	0.0206 (5)	0.0152 (5)	-0.0011 (4)	0.0014 (4)	0.0002 (4)
C8	0.0152 (5)	0.0195 (5)	0.0123 (4)	-0.0006 (4)	0.0012 (4)	0.0002 (4)
C9	0.0146 (5)	0.0170 (5)	0.0145 (5)	-0.0009 (4)	0.0008 (4)	0.0000 (4)
C10	0.0144 (4)	0.0154 (5)	0.0153 (5)	-0.0014 (4)	0.0013 (4)	-0.0013 (4)
C11	0.0157 (5)	0.0170 (5)	0.0127 (4)	-0.0005 (4)	0.0013 (4)	0.0014 (4)
C12	0.0134 (5)	0.0193 (5)	0.0153 (5)	0.0004 (4)	-0.0006 (4)	-0.0002 (4)
C13	0.0157 (5)	0.0155 (5)	0.0172 (5)	0.0015 (4)	-0.0007 (4)	0.0010 (4)
C14	0.0162 (5)	0.0167 (5)	0.0133 (4)	0.0001 (4)	-0.0001 (4)	0.0014 (4)
C15	0.0156 (5)	0.0143 (5)	0.0158 (5)	-0.0006 (4)	0.0005 (4)	-0.0017 (4)
C16	0.0142 (5)	0.0158 (5)	0.0156 (5)	0.0001 (4)	0.0001 (4)	-0.0010 (4)
C17	0.0151 (5)	0.0186 (5)	0.0140 (5)	0.0001 (4)	0.0016 (4)	-0.0004 (4)
C18	0.0130 (4)	0.0170 (5)	0.0169 (5)	0.0006 (4)	0.0022 (4)	-0.0014 (4)
C19	0.0229 (6)	0.0258 (6)	0.0204 (5)	-0.0011 (4)	-0.0012 (4)	-0.0004 (5)
C20	0.0164 (5)	0.0262 (6)	0.0265 (6)	-0.0046 (4)	0.0017 (4)	-0.0065 (5)
C21	0.0228 (6)	0.0233 (6)	0.0241 (6)	-0.0060 (4)	0.0065 (4)	-0.0004 (5)
C22	0.0205 (5)	0.0246 (6)	0.0224 (6)	0.0013 (4)	-0.0006 (4)	0.0009 (5)
C23	0.0265 (6)	0.0303 (6)	0.0241 (6)	-0.0045 (5)	-0.0075 (5)	0.0015 (5)
C24	0.0164 (5)	0.0265 (6)	0.0309 (6)	-0.0037 (4)	-0.0019 (4)	-0.0038 (5)
C25	0.0213 (6)	0.0293 (6)	0.0252 (6)	-0.0060 (5)	0.0034 (4)	0.0002 (5)
C26	0.0209 (6)	0.0312 (6)	0.0205 (5)	-0.0034 (5)	-0.0034 (4)	0.0011 (5)
C27	0.0204 (6)	0.0325 (6)	0.0231 (6)	-0.0072 (5)	0.0038 (4)	-0.0011 (5)
C28	0.0176 (5)	0.0300 (6)	0.0238 (6)	-0.0038 (4)	-0.0016 (4)	-0.0052 (5)
C29	0.0240 (6)	0.0303 (6)	0.0172 (5)	-0.0025 (5)	-0.0015 (4)	-0.0008 (5)
C30	0.0212 (5)	0.0284 (6)	0.0217 (6)	-0.0041 (4)	0.0030 (4)	0.0003 (5)

supplementary materials

Geometric parameters (\AA , $^\circ$)

C11—C2	1.7293 (11)	C3—C4	1.5412 (15)
C12—C5	1.7218 (11)	C4—C5	1.4545 (14)
C13—C8	1.7262 (11)	C5—C6	1.3522 (15)
C14—C11	1.7192 (11)	C7—C8	1.4110 (14)
C15—C14	1.7305 (11)	C7—C12	1.5102 (15)
C16—C17	1.7168 (11)	C8—C9	1.3958 (15)
O1—C1	1.2408 (13)	C9—C10	1.5398 (15)
O2—C3	1.2517 (13)	C10—C11	1.4515 (14)
O3—C4	1.2193 (13)	C11—C12	1.3533 (15)
O4—C6	1.3218 (13)	C13—C14	1.4096 (15)
O4—H4	0.78 (2)	C13—C18	1.5069 (15)
O5—C7	1.2417 (13)	C14—C15	1.4040 (14)
O6—C9	1.2536 (13)	C15—C16	1.5408 (15)
O7—C10	1.2213 (13)	C16—C17	1.4481 (15)
O8—C12	1.3207 (12)	C17—C18	1.3520 (14)
O8—H8	0.80 (2)	C19—C20	1.3969 (17)
O9—C13	1.2476 (13)	C19—H19	0.9500
O10—C15	1.2491 (13)	C20—C21	1.3708 (17)
O11—C16	1.2218 (13)	C20—H20	0.9500
O12—C18	1.3232 (13)	C21—C22	1.3906 (16)
O12—H12	0.79 (2)	C21—H21	0.9500
N1—C22	1.3158 (16)	C22—H22	0.9500
N1—N2	1.3331 (15)	C23—C24	1.3994 (19)
N1—H1	0.888 (18)	C23—H23	0.9500
N2—C19	1.3189 (15)	C24—C25	1.3666 (18)
N3—C26	1.3144 (16)	C24—H24	0.9500
N3—N4	1.3355 (14)	C25—C26	1.3942 (16)
N3—H3	0.886 (17)	C25—H25	0.9500
N4—C23	1.3223 (16)	C26—H26	0.9500
N5—C30	1.3221 (16)	C27—C28	1.3966 (18)
N5—N6	1.3314 (14)	C27—H27	0.9500
N5—H5	0.891 (18)	C28—C29	1.3690 (17)
N6—C27	1.3236 (15)	C28—H28	0.9500
C1—C2	1.4069 (14)	C29—C30	1.3875 (16)
C1—C6	1.5079 (15)	C29—H29	0.9500
C2—C3	1.3913 (14)	C30—H30	0.9500
C6—O4—H4	113.0 (17)	C14—C13—C18	118.12 (9)
C12—O8—H8	112.9 (16)	C15—C14—C13	122.76 (10)
C18—O12—H12	110.5 (15)	C15—C14—Cl5	118.65 (8)
C22—N1—N2	125.20 (10)	C13—C14—Cl5	118.58 (8)
C22—N1—H1	122.0 (12)	O10—C15—C14	125.90 (10)
N2—N1—H1	112.8 (12)	O10—C15—C16	116.70 (9)
C19—N2—N1	115.76 (10)	C14—C15—C16	117.40 (9)
C26—N3—N4	125.10 (10)	O11—C16—C17	122.83 (10)
C26—N3—H3	121.5 (11)	O11—C16—C15	117.95 (10)
N4—N3—H3	113.4 (11)	C17—C16—C15	119.21 (9)

C23—N4—N3	115.44 (11)	C18—C17—C16	120.10 (10)
C30—N5—N6	124.80 (10)	C18—C17—Cl6	120.99 (8)
C30—N5—H5	120.8 (12)	C16—C17—Cl6	118.88 (8)
N6—N5—H5	114.4 (12)	O12—C18—C17	120.69 (10)
C27—N6—N5	115.74 (10)	O12—C18—C13	117.06 (9)
O1—C1—C2	125.21 (10)	C17—C18—C13	122.23 (10)
O1—C1—C6	116.67 (10)	N2—C19—C20	123.68 (11)
C2—C1—C6	118.09 (9)	N2—C19—H19	118.2
C3—C2—C1	122.90 (10)	C20—C19—H19	118.2
C3—C2—Cl1	119.09 (8)	C21—C20—C19	118.28 (11)
C1—C2—Cl1	118.00 (8)	C21—C20—H20	120.9
O2—C3—C2	124.89 (10)	C19—C20—H20	120.9
O2—C3—C4	116.93 (9)	C20—C21—C22	117.14 (11)
C2—C3—C4	118.17 (9)	C20—C21—H21	121.4
O3—C4—C5	123.41 (10)	C22—C21—H21	121.4
O3—C4—C3	118.33 (9)	N1—C22—C21	119.91 (11)
C5—C4—C3	118.24 (9)	N1—C22—H22	120.0
C6—C5—C4	120.45 (10)	C21—C22—H22	120.0
C6—C5—Cl2	120.77 (8)	N4—C23—C24	123.93 (12)
C4—C5—Cl2	118.72 (8)	N4—C23—H23	118.0
O4—C6—C5	121.29 (10)	C24—C23—H23	118.0
O4—C6—C1	116.75 (10)	C25—C24—C23	118.30 (11)
C5—C6—C1	121.96 (10)	C25—C24—H24	120.8
O5—C7—C8	125.43 (10)	C23—C24—H24	120.8
O5—C7—C12	116.43 (9)	C24—C25—C26	116.95 (12)
C8—C7—C12	118.14 (9)	C24—C25—H25	121.5
C9—C8—C7	122.55 (10)	C26—C25—H25	121.5
C9—C8—Cl3	119.72 (8)	N3—C26—C25	120.28 (11)
C7—C8—Cl3	117.69 (8)	N3—C26—H26	119.9
O6—C9—C8	125.73 (10)	C25—C26—H26	119.9
O6—C9—C10	116.17 (9)	N6—C27—C28	123.96 (12)
C8—C9—C10	118.09 (9)	N6—C27—H27	118.0
O7—C10—C11	122.72 (10)	C28—C27—H27	118.0
O7—C10—C9	118.35 (9)	C29—C28—C27	117.95 (11)
C11—C10—C9	118.92 (9)	C29—C28—H28	121.0
C12—C11—C10	119.98 (10)	C27—C28—H28	121.0
C12—C11—Cl4	121.19 (8)	C28—C29—C30	117.43 (11)
C10—C11—Cl4	118.79 (8)	C28—C29—H29	121.3
O8—C12—C11	121.01 (10)	C30—C29—H29	121.3
O8—C12—C7	116.74 (9)	N5—C30—C29	120.10 (11)
C11—C12—C7	122.25 (9)	N5—C30—H30	119.9
O9—C13—C14	125.75 (10)	C29—C30—H30	119.9
O9—C13—C18	116.14 (10)		
C22—N1—N2—C19	-0.02 (18)	C10—C11—C12—C7	-2.38 (17)
C26—N3—N4—C23	0.15 (19)	Cl4—C11—C12—C7	179.80 (8)
C30—N5—N6—C27	-0.32 (18)	O5—C7—C12—O8	0.75 (15)
O1—C1—C2—C3	-176.99 (12)	C8—C7—C12—O8	-179.96 (10)
C6—C1—C2—C3	1.03 (16)	O5—C7—C12—C11	-179.30 (11)
O1—C1—C2—Cl1	3.15 (17)	C8—C7—C12—C11	-0.01 (16)

supplementary materials

C6—C1—C2—Cl1	-178.82 (8)	O9—C13—C14—C15	179.75 (11)
C1—C2—C3—O2	178.89 (11)	C18—C13—C14—C15	-0.34 (16)
Cl1—C2—C3—O2	-1.25 (16)	O9—C13—C14—Cl5	-1.12 (16)
C1—C2—C3—C4	-1.82 (16)	C18—C13—C14—Cl5	178.79 (8)
Cl1—C2—C3—C4	178.04 (8)	C13—C14—C15—O10	176.55 (11)
O2—C3—C4—O3	4.40 (15)	Cl5—C14—C15—O10	-2.58 (16)
C2—C3—C4—O3	-174.94 (10)	C13—C14—C15—C16	-3.15 (16)
O2—C3—C4—C5	-177.03 (10)	Cl5—C14—C15—C16	177.72 (7)
C2—C3—C4—C5	3.63 (14)	O10—C15—C16—O11	2.96 (15)
O3—C4—C5—C6	173.62 (11)	C14—C15—C16—O11	-177.31 (10)
C3—C4—C5—C6	-4.87 (15)	O10—C15—C16—C17	-175.99 (10)
O3—C4—C5—Cl2	-3.52 (15)	C14—C15—C16—C17	3.74 (15)
C3—C4—C5—Cl2	177.99 (8)	O11—C16—C17—C18	-179.53 (11)
C4—C5—C6—O4	-176.09 (10)	C15—C16—C17—C18	-0.63 (15)
Cl2—C5—C6—O4	1.00 (15)	O11—C16—C17—Cl6	-1.56 (15)
C4—C5—C6—C1	4.23 (16)	C15—C16—C17—Cl6	177.34 (8)
Cl2—C5—C6—C1	-178.68 (8)	C16—C17—C18—O12	178.68 (10)
O1—C1—C6—O4	-3.73 (15)	Cl6—C17—C18—O12	0.75 (15)
C2—C1—C6—O4	178.08 (10)	C16—C17—C18—C13	-3.07 (16)
O1—C1—C6—C5	175.96 (11)	Cl6—C17—C18—C13	179.00 (8)
C2—C1—C6—C5	-2.23 (16)	O9—C13—C18—O12	1.95 (15)
O5—C7—C8—C9	-179.02 (11)	C14—C13—C18—O12	-177.98 (10)
C12—C7—C8—C9	1.75 (16)	O9—C13—C18—C17	-176.36 (11)
O5—C7—C8—Cl3	-1.27 (16)	C14—C13—C18—C17	3.71 (16)
C12—C7—C8—Cl3	179.50 (8)	N1—N2—C19—C20	-0.85 (18)
C7—C8—C9—O6	178.05 (11)	N2—C19—C20—C21	0.50 (19)
Cl3—C8—C9—O6	0.34 (16)	C19—C20—C21—C22	0.68 (18)
C7—C8—C9—C10	-1.04 (16)	N2—N1—C22—C21	1.20 (19)
Cl3—C8—C9—C10	-178.74 (8)	C20—C21—C22—N1	-1.48 (18)
O6—C9—C10—O7	-1.84 (15)	N3—N4—C23—C24	-0.8 (2)
C8—C9—C10—O7	177.33 (10)	N4—C23—C24—C25	0.9 (2)
O6—C9—C10—C11	179.47 (10)	C23—C24—C25—C26	-0.24 (19)
C8—C9—C10—C11	-1.36 (15)	N4—N3—C26—C25	0.4 (2)
O7—C10—C11—C12	-175.60 (11)	C24—C25—C26—N3	-0.35 (19)
C9—C10—C11—C12	3.03 (15)	N5—N6—C27—C28	0.92 (19)
O7—C10—C11—Cl4	2.27 (15)	N6—C27—C28—C29	-0.4 (2)
C9—C10—C11—Cl4	-179.10 (8)	C27—C28—C29—C30	-0.63 (19)
C10—C11—C12—O8	177.57 (10)	N6—N5—C30—C29	-0.7 (2)
Cl4—C11—C12—O8	-0.24 (16)	C28—C29—C30—N5	1.20 (19)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1···O2	0.889 (17)	1.753 (17)	2.6138 (13)	162.2 (17)
N1—H1···O3	0.889 (17)	2.578 (18)	3.1583 (14)	123.7 (14)
N3—H3···O6	0.886 (18)	1.811 (17)	2.6621 (13)	160.4 (17)
N3—H3···O7	0.886 (18)	2.431 (18)	3.0066 (14)	123.0 (14)
O4—H4···O1	0.79 (2)	2.26 (2)	2.6619 (13)	113 (2)
O4—H4···O5	0.79 (2)	1.94 (2)	2.6723 (12)	155 (2)

supplementary materials

N5—H5···O10	0.890 (18)	1.881 (18)	2.7450 (14)	163.2 (16)
N5—H5···O11	0.890 (18)	2.410 (17)	2.9659 (13)	120.7 (14)
O8—H8···O1	0.80 (2)	1.86 (2)	2.5930 (12)	152 (2)
O8—H8···O5	0.80 (2)	2.25 (2)	2.6582 (13)	112.7 (18)
O12—H12···O9	0.79 (2)	2.23 (2)	2.6595 (13)	115.1 (18)
O12—H12···O9 ⁱ	0.79 (2)	2.02 (2)	2.6801 (12)	142 (2)
C20—H20···O2 ⁱⁱ	0.95	2.57	3.1097 (15)	116
C21—H21···O3 ⁱⁱⁱ	0.95	2.36	3.2754 (15)	162
C22—H22···O3	0.95	2.60	3.1913 (15)	121
C25—H25···O10	0.95	2.47	3.4056 (15)	169
C26—H26···O7	0.95	2.44	3.0363 (15)	121
C26—H26···N6	0.95	2.51	3.2665 (17)	137
C27—H27···O7	0.95	2.37	3.2765 (15)	159
C28—H28···O6 ^{iv}	0.95	2.47	3.1091 (15)	124
C30—H30···O11	0.95	2.34	2.9551 (15)	122

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+2, y+1/2, -z+3/2$; (iii) $-x+2, -y+3, -z+1$; (iv) $x, -y+3/2, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

