

Ethyl 1,3-dimethyl-1*H*-indole-2-carboxylate

A. Marx,^a G. Chakkaravarthi,^b G. Gobirajeshwaran,^c
A. K. Mohanakrishnan^c and V. Manivannan^{a*}

^aDepartment of Physics, Presidency College, Chennai 600 005, India, ^bDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, and ^cDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India
Correspondence e-mail: manivan_1999@yahoo.com


Received 9 October 2008; accepted 19 October 2008

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$; R factor = 0.056; wR factor = 0.184; data-to-parameter ratio = 24.5.

In the title compound, $\text{C}_{13}\text{H}_{15}\text{NO}_2$, the plane of the indole ring forms a dihedral angle of $5.26(6)^\circ$ with the ester group and the ethyl side-chain C atoms. The crystal packing is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For biological activities of indole derivatives, see: Okabe & Adachi (1998); Schollmeyer *et al.* (1995). For related structures, see: Chakkaravarthi *et al.* (2007, 2008).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{15}\text{NO}_2$	$c = 12.9449(5) \text{ \AA}$
$M_r = 217.26$	$\beta = 105.488(2)^\circ$
Monoclinic, $P2_1/c$	$V = 1153.71(9) \text{ \AA}^3$
$a = 7.5511(3) \text{ \AA}$	$Z = 4$
$b = 12.2476(6) \text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.08 \text{ mm}^{-1}$
 $T = 295(2) \text{ K}$

$0.25 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker Kappa APEXII
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.979$, $T_{\max} = 0.983$

15440 measured reflections
3620 independent reflections
2068 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.184$
 $S = 1.03$
3620 reflections

148 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.21 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C2—H2···O1 ⁱ	0.93	2.53	3.401 (2)	156
C12—H12A···Cg1 ⁱⁱ	0.97	2.76	3.646 (2)	152

Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 2, -y, -z + 1$. Cg1 is the centroid of the C1–C6 ring.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT; data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97.

The authors acknowledge the Sophisticated Analytical Instrument Facility, Indian Institute of Technology, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2100).

References

- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2007). *Acta Cryst. E63*, o3673.
Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008). *Acta Cryst. E64*, o542.
Okabe, N. & Adachi, Y. (1998). *Acta Cryst. C54*, 386–387.
Schollmeyer, D., Fischer, G. & Pindur, U. (1995). *Acta Cryst. C51*, 2572–2575.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o2200 [doi:10.1107/S1600536808034156]

Ethyl 1,3-dimethyl-1*H*-indole-2-carboxylate

A. Marx, G. Chakkavarthi, G. Gobirajeshwaran, A. K. Mohanakrishnan and V. Manivannan

Comment

The indole derivatives are found to possess antibacterial (Okabe and Adachi, 1998) and antitumour activities (Schollmeyer *et al.*, 1995). In continuation to our studies in indole derivatives, we determine the crystal structure of the title compound (I). The geometric parameters of the molecule of (I) (Fig. 1) agree well with the reported structures (Chakkavarthi *et al.*, 2007; Chakkavarthi *et al.*, 2008).

The five- (N1/C1/C6/C7/C8) and six- (C1—C6) membered rings in the indane group are almost planar, with a dihedral angle of 1.67 (6) $^{\circ}$ between these rings. The plane of indole ring forms a dihedral angle of 5.26 (6) $^{\circ}$ with the ester group. The molecular packing is stabilized by weak intramolecular C—H···O interaction and the crystal packing of (I) is stabilized by weak intermolecular C—H···O and C—H··· π interactions (Table 1) (Fig. 2).

Experimental

To a stirred suspension of NaH (0.6 mmol, hexane washed) in THF (2 ml), a solution of 2-carbethoxy-3-methyl indole (0.5 mmol) in THF (2 ml) was added and stirred for 30 minutes at room temperature. To the reaction mixture, a solution of Iodomethane (0.6 mmol) was added and stirring was continued for further 6 hr. After the indole was consumed (monitored by TLC), the reaction mixture was quenched with cold dil HCl (25 ml), extracted with ethyl acetate (2 x 10 ml) and dried (Na_2SO_4). Removal of solvent followed by crystallization (hexane) afforded as yellow crystal.

Refinement

H atoms were positioned geometrically and refined using riding model with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2\text{U}_{\text{eq}}(\text{C})$ for aromatic H atoms, C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2\text{U}_{\text{eq}}(\text{C})$ for CH₂ and C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5\text{U}_{\text{eq}}(\text{C})$ for methyl H atoms.

Figures




Fig. 1. The molecular structure of (I), with atom labeling scheme. Displacement ellipsoids are drawn at 50% probability level.

supplementary materials




Fig. 2. The crystal structure of (I), viewed down the a face. For the sake of clarity, H atoms not involved in interaction have been omitted.

Ethyl 1,3-dimethyl-1*H*-indole-2-carboxylate

Crystal data

C ₁₃ H ₁₅ NO ₂	$F_{000} = 464$
$M_r = 217.26$	$D_x = 1.251 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 7.5511 (3) \text{ \AA}$	Cell parameters from 4312 reflections
$b = 12.2476 (6) \text{ \AA}$	$\theta = 2.3\text{--}24.1^\circ$
$c = 12.9449 (5) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\beta = 105.488 (2)^\circ$	$T = 295 (2) \text{ K}$
$V = 1153.71 (9) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.25 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker Kappa APEXII diffractometer	3620 independent reflections
Radiation source: fine-focus sealed tube	2068 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 295(2) \text{ K}$	$\theta_{\text{max}} = 31.0^\circ$
ω and φ scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.979$, $T_{\text{max}} = 0.983$	$k = -17 \rightarrow 17$
15440 measured reflections	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H-atom parameters constrained
$wR(F^2) = 0.184$	$w = 1/[\sigma^2(F_o^2) + (0.0844P)^2 + 0.1542P]$
	where $P = (F_o^2 + 2F_c^2)/3$

S = 1.03	$(\Delta/\sigma)_{\max} < 0.001$
3620 reflections	$\Delta\rho_{\max} = 0.21 \text{ e \AA}^{-3}$
148 parameters	$\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7616 (2)	-0.19859 (13)	0.57067 (13)	0.0527 (4)
C2	0.7747 (2)	-0.30504 (16)	0.61130 (15)	0.0653 (5)
H2	0.8324	-0.3194	0.6829	0.078*
C3	0.7003 (3)	-0.38637 (16)	0.54252 (18)	0.0750 (5)
H3	0.7075	-0.4577	0.5679	0.090*
C4	0.6132 (3)	-0.36648 (16)	0.43493 (18)	0.0738 (5)
H4	0.5646	-0.4246	0.3901	0.089*
C5	0.5983 (2)	-0.26368 (16)	0.39465 (14)	0.0655 (4)
H5	0.5386	-0.2510	0.3230	0.079*
C6	0.67448 (19)	-0.17669 (13)	0.46276 (12)	0.0525 (4)
C7	0.68784 (19)	-0.06336 (13)	0.44615 (11)	0.0501 (4)
C8	0.78272 (19)	-0.01966 (13)	0.54312 (11)	0.0512 (4)
C9	0.9305 (3)	-0.09425 (17)	0.73226 (14)	0.0757 (5)
H9A	0.9384	-0.1651	0.7650	0.114*
H9B	0.8689	-0.0449	0.7687	0.114*
H9C	1.0519	-0.0675	0.7371	0.114*
C10	0.6076 (2)	-0.00822 (15)	0.34166 (13)	0.0670 (5)
H10A	0.5464	0.0575	0.3534	0.100*
H10B	0.5209	-0.0561	0.2955	0.100*
H10C	0.7038	0.0095	0.3088	0.100*
C11	0.8333 (2)	0.09395 (14)	0.57069 (13)	0.0559 (4)
C12	0.8335 (2)	0.27250 (13)	0.50367 (14)	0.0644 (4)
H12A	0.9652	0.2837	0.5286	0.077*
H12B	0.7771	0.3007	0.5573	0.077*
C13	0.7583 (3)	0.32875 (18)	0.40024 (17)	0.0805 (6)
H13A	0.8079	0.2962	0.3465	0.121*
H13B	0.7913	0.4046	0.4077	0.121*
H13C	0.6269	0.3219	0.3793	0.121*
N1	0.82742 (18)	-0.10221 (11)	0.61980 (10)	0.0568 (4)
O1	0.9033 (2)	0.12730 (12)	0.65918 (10)	0.0943 (5)
O2	0.79272 (16)	0.15785 (9)	0.48555 (9)	0.0632 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0489 (7)	0.0561 (9)	0.0535 (9)	0.0029 (6)	0.0146 (6)	0.0022 (7)
C2	0.0636 (9)	0.0662 (11)	0.0666 (11)	0.0026 (8)	0.0182 (8)	0.0136 (9)
C3	0.0774 (12)	0.0564 (11)	0.0939 (15)	-0.0017 (9)	0.0278 (11)	0.0091 (10)
C4	0.0742 (11)	0.0606 (11)	0.0849 (14)	-0.0079 (9)	0.0183 (10)	-0.0079 (10)

supplementary materials

C5	0.0627 (9)	0.0696 (11)	0.0595 (10)	-0.0025 (8)	0.0081 (7)	-0.0072 (9)
C6	0.0476 (7)	0.0574 (9)	0.0519 (8)	0.0043 (6)	0.0122 (6)	0.0002 (7)
C7	0.0499 (7)	0.0552 (9)	0.0439 (8)	0.0047 (6)	0.0104 (6)	-0.0002 (6)
C8	0.0529 (8)	0.0556 (9)	0.0459 (8)	0.0041 (6)	0.0144 (6)	0.0012 (7)
C9	0.0883 (13)	0.0815 (13)	0.0488 (9)	0.0011 (10)	0.0035 (9)	0.0088 (9)
C10	0.0770 (11)	0.0675 (11)	0.0492 (9)	0.0038 (9)	0.0043 (8)	0.0050 (8)
C11	0.0624 (9)	0.0597 (10)	0.0462 (8)	0.0004 (7)	0.0155 (7)	-0.0025 (7)
C12	0.0722 (10)	0.0549 (10)	0.0682 (11)	0.0016 (8)	0.0226 (8)	-0.0044 (8)
C13	0.0902 (13)	0.0688 (12)	0.0804 (13)	0.0093 (10)	0.0189 (11)	0.0097 (10)
N1	0.0588 (7)	0.0631 (9)	0.0467 (7)	0.0022 (6)	0.0109 (6)	0.0036 (6)
O1	0.1481 (14)	0.0728 (9)	0.0498 (7)	-0.0162 (9)	0.0050 (8)	-0.0092 (6)
O2	0.0805 (8)	0.0518 (7)	0.0536 (7)	-0.0007 (5)	0.0113 (6)	-0.0014 (5)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.370 (2)	C9—N1	1.459 (2)
C1—C2	1.400 (2)	C9—H9A	0.9600
C1—C6	1.402 (2)	C9—H9B	0.9600
C2—C3	1.354 (3)	C9—H9C	0.9600
C2—H2	0.9300	C10—H10A	0.9600
C3—C4	1.394 (3)	C10—H10B	0.9600
C3—H3	0.9300	C10—H10C	0.9600
C4—C5	1.356 (3)	C11—O1	1.198 (2)
C4—H4	0.9300	C11—O2	1.319 (2)
C5—C6	1.404 (2)	C12—O2	1.443 (2)
C5—H5	0.9300	C12—C13	1.478 (3)
C6—C7	1.412 (2)	C12—H12A	0.9700
C7—C8	1.377 (2)	C12—H12B	0.9700
C7—C10	1.488 (2)	C13—H13A	0.9600
C8—N1	1.394 (2)	C13—H13B	0.9600
C8—C11	1.462 (2)	C13—H13C	0.9600
N1—C1—C2	130.38 (16)	H9A—C9—H9C	109.5
N1—C1—C6	108.26 (14)	H9B—C9—H9C	109.5
C2—C1—C6	121.33 (16)	C7—C10—H10A	109.5
C3—C2—C1	117.57 (17)	C7—C10—H10B	109.5
C3—C2—H2	121.2	H10A—C10—H10B	109.5
C1—C2—H2	121.2	C7—C10—H10C	109.5
C2—C3—C4	122.06 (18)	H10A—C10—H10C	109.5
C2—C3—H3	119.0	H10B—C10—H10C	109.5
C4—C3—H3	119.0	O1—C11—O2	122.77 (16)
C5—C4—C3	121.00 (18)	O1—C11—C8	125.30 (16)
C5—C4—H4	119.5	O2—C11—C8	111.93 (14)
C3—C4—H4	119.5	O2—C12—C13	107.02 (15)
C4—C5—C6	119.02 (17)	O2—C12—H12A	110.3
C4—C5—H5	120.5	C13—C12—H12A	110.3
C6—C5—H5	120.5	O2—C12—H12B	110.3
C1—C6—C5	119.01 (15)	C13—C12—H12B	110.3
C1—C6—C7	107.83 (14)	H12A—C12—H12B	108.6
C5—C6—C7	133.16 (15)	C12—C13—H13A	109.5

C8—C7—C6	106.51 (13)	C12—C13—H13B	109.5
C8—C7—C10	129.70 (16)	H13A—C13—H13B	109.5
C6—C7—C10	123.78 (14)	C12—C13—H13C	109.5
C7—C8—N1	109.60 (14)	H13A—C13—H13C	109.5
C7—C8—C11	129.14 (14)	H13B—C13—H13C	109.5
N1—C8—C11	121.25 (14)	C1—N1—C8	107.79 (13)
N1—C9—H9A	109.5	C1—N1—C9	123.47 (14)
N1—C9—H9B	109.5	C8—N1—C9	128.70 (15)
H9A—C9—H9B	109.5	C11—O2—C12	116.72 (13)
N1—C9—H9C	109.5		
N1—C1—C2—C3	−177.99 (16)	C6—C7—C8—C11	−179.58 (14)
C6—C1—C2—C3	0.1 (2)	C10—C7—C8—C11	−0.7 (3)
C1—C2—C3—C4	0.0 (3)	C7—C8—C11—O1	173.58 (17)
C2—C3—C4—C5	−0.5 (3)	N1—C8—C11—O1	−5.4 (3)
C3—C4—C5—C6	0.9 (3)	C7—C8—C11—O2	−6.4 (2)
N1—C1—C6—C5	178.77 (13)	N1—C8—C11—O2	174.65 (12)
C2—C1—C6—C5	0.3 (2)	C2—C1—N1—C8	178.04 (15)
N1—C1—C6—C7	−0.08 (16)	C6—C1—N1—C8	−0.24 (16)
C2—C1—C6—C7	−178.54 (14)	C2—C1—N1—C9	−0.1 (3)
C4—C5—C6—C1	−0.8 (2)	C6—C1—N1—C9	−178.40 (14)
C4—C5—C6—C7	177.72 (16)	C7—C8—N1—C1	0.48 (16)
C1—C6—C7—C8	0.37 (16)	C11—C8—N1—C1	179.62 (13)
C5—C6—C7—C8	−178.25 (16)	C7—C8—N1—C9	178.52 (15)
C1—C6—C7—C10	−178.61 (14)	C11—C8—N1—C9	−2.3 (2)
C5—C6—C7—C10	2.8 (3)	O1—C11—O2—C12	−1.2 (2)
C6—C7—C8—N1	−0.52 (15)	C8—C11—O2—C12	178.80 (13)
C10—C7—C8—N1	178.38 (15)	C13—C12—O2—C11	−173.84 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C10—H10A···O2	0.96	2.49	2.858 (2)	103
C2—H2···O1 ⁱ	0.93	2.53	3.401 (2)	156
C12—H12A···Cg1 ⁱⁱ	0.97	2.76	3.646 (2)	152

Symmetry codes: (i) $-x+2, y-1/2, -z+3/2$; (ii) $-x+2, -y, -z+1$.

supplementary materials

Fig. 1




Fig. 2

