

Aqua[2,6-bis(2-pyridylamino)pyridine]-sulfatonickel(II) monohydrate

Lu Lu,* Jun Wang, Bin Xie and Li-Ke Zou

Department of Chemistry and Pharmaceutical Engineering, Sichuan University of Science and Engineering, Zigong, Sichuan 643000, People's Republic of China
Correspondence e-mail: lulusczg@126.com


Received 23 October 2008; accepted 28 October 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.003$ Å;
 R factor = 0.021; wR factor = 0.056; data-to-parameter ratio = 11.8.

The Ni atom in the title complex, $[Ni(SO_4)(C_{15}H_{13}N_5)(H_2O)] \cdot H_2O$, has a distorted trigonal-bipyramidal coordination formed by the tridentate 2,6-bis(2-pyridylamino)pyridine (tpdaH₂) ligand, one sulfate and one coordinated water molecule. The tpdaH₂ ligand is three-coordinated, with the N atom of the central pyridine ring in the equatorial position [Ni—N = 1.9961 (14) Å] and the N atoms of the peripheral pyridine rings in the axial positions [Ni—N = 1.9668 (15) and 1.9895 (15) Å]. The remaining equatorial positions are occupied by the O atoms of the sulfate ligand and the water molecule. The H atoms of both NH groups of the tpdaH₂ ligand are involved in hydrogen bonds with the O atoms of the uncoordinated water molecule and the sulfate group which link the complex molecules, forming an infinite three-dimensional network.

Related literature

For the properties of transition metal complexes with poly-pyridylamine ligands, see: Wang *et al.* (1999). For the tri-pyridylamine ligand, see: Jing *et al.* (2000). For metal–metal interactions, see: Cotton *et al.* (1998); Yang *et al.* (1997).



Experimental

Crystal data

$[Ni(SO_4)(C_{15}H_{13}N_5)(H_2O)] \cdot H_2O$	$V = 1703.3$ (3) Å ³
$M_r = 454.11$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 7.3536$ (8) Å	$\mu = 1.31$ mm ⁻¹
$b = 18.026$ (2) Å	$T = 298$ (2) K
$c = 12.9125$ (14) Å	$0.22 \times 0.16 \times 0.12$ mm
$\beta = 95.634$ (2)°	

Data collection

Bruker APEXII area-detector	8650 measured reflections
diffractometer	3068 independent reflections
Absorption correction: multi-scan	2821 reflections with $I > 2\sigma(I)$
(SADABS; Bruker, 2004)	$R_{\text{int}} = 0.014$
	$T_{\min} = 0.761$, $T_{\max} = 0.859$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of
$wR(F^2) = 0.056$	independent and constrained
$S = 1.06$	refinement
3068 reflections	$\Delta\rho_{\max} = 0.24$ e Å ⁻³
259 parameters	$\Delta\rho_{\min} = -0.32$ e Å ⁻³
8 restraints	

Table 1
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N2—H2B···O6 ⁱ	0.866 (9)	2.079 (10)	2.9433 (19)	176 (2)
N4—H4B···O4 ⁱⁱ	0.862 (9)	2.050 (10)	2.8967 (18)	167.1 (18)
O5—H5A···O6	0.86	2.08	2.9112 (18)	163
O5—H5B···O4 ⁱⁱⁱ	0.85	1.98	2.8192 (19)	169
O6—H6A···O2 ^{iv}	0.85	1.91	2.7531 (18)	173
O6—H6B···O3 ⁱⁱⁱ	0.85	1.97	2.785 (2)	162

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x + 1, -y, -z$; (iii) $x - 1, y, z$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996) and *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors are grateful to Sichuan University of Science and Engineering for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2397).

References

- Bruker (2004). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
Cotton, F. A., Daniels, L. M., Murillo, C. A. & Wang, X. (1998). *Chem. Commun.* pp. 39–40.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.

- Jing, B.-W., Wu, T., Zhang, M.-W. & Shen, T. (2000). *Chem. J. Chin. Univ.* **21**, 395–400.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Wang, Z., Xiong, R.-G., Naggar, E., Foxman, B. M. & Lin, W.-B. (1999). *Inorg. Chim. Acta*, **288**, 215–219.
Yang, M. H., Lin, T. W., Chou, C. C., Lee, H. C., Chang, H. C., Lee, G. H., Leung, M. K. & Peng, S. M. (1997). *Chem. Commun.* pp. 39–40.

supplementary materials

Acta Cryst. (2008). E64, m1490-m1491 [doi:10.1107/S1600536808035101]

Aqua[2,6-bis(2-pyridylamino)pyridine]sulfatonickel(II) monohydrate

L. Lu, J. Wang, B. Xie and L.-K. Zou

Comment

Transition metal complexes with polypyridylamine ligands, possessing diverse structures and special optical and electro-magnetic properties (Wang *et al.*, 1999), have aroused great interest among researchers. Tri-pyridylamine ligand usually exhibits donor as well as acceptor properties and can be used as a popular chelating ligand (Jing *et al.*, 2000). In recent years great efforts have been taken to synthesize and characterize metal chain complexes which can be used to study the metal-metal interactions (Yang *et al.*, 1997; Cotton *et al.*, 1998). Herein we report the synthesis and crystal structure of the title complex with tri-pyridylamine ligand.

The Ni1 atom in the title complex has a distorted trigonal-bipyramidal coordination formed by the tridentate tpdaH₂ ligand, one sulfate and one coordinated water molecule. (Fig. 1). The tpdaH₂ ligand is tri-coordinated, with the peripheral N1 and N5 atoms in the axial positions [Ni1—N1 = 1.9895 (15) Å, Ni1—N5 = 1.9668 (15) Å and N1—Ni1—N5 = 169.26 (6)°] and the central N3 atom in the equatorial plane of the bipyramid [Ni1—N3 = 1.9961 (14) Å]. The remaining equatorial positions are occupied by one sulfate and one coordinated water molecule. Selected geometric parameters have been listed in tabel 1.

The three pyridine rings of the tpdaH₂ ligand are not coplanar. The dihedral angles between the planes of the central pyridine ring and two peripheral rings are 15.0 (7) and 22.7 (3)° respectively. In the title complex the two H atoms of both NH groups of tpdaH₂ act as active H atoms in forming inter-molecular classical hydrogen bonds (Table 2). The inter-molecular hydrogen bonds function greatly in linking the complex to be a infinite three-dimensional network.

Experimental

Tripyridylamine (0.031 g, 0.12 mmol), NiSO₄ (0.26 g, 0.13 mmol), were added in a solvent of acetonitrile, the mixture was heated for six hours under reflux. during the process stirring and influx were required. The resultant was then filtered to give a pure solution which was infiltrated by diethyl ether freely in a closed vessel, three weeks later some single crystals of the size suitable for X-Ray diffraction analysis.

Refinement

Carbon H atoms were positioned geometrically and treated as riding on their parent atoms, with C—H distances of 0.93 Å (pyridine ring) with $U_{\text{iso}}(\text{H})$ 1.2 $U_{\text{eq}}(\text{C})$. The amine H atoms were located in difference maps and freely refined with $U_{\text{iso}}(\text{H})$ 1.2 $U_{\text{eq}}(\text{N})$. The water H atoms were located in different map and, in the first stage of refinement, refined with the O—H and H—H distances restraints to 0.85 Å and 1.39 Å respectively and with $U_{\text{iso}}(\text{H})$ 1.5 $U_{\text{eq}}(\text{O})$. In the last cycle, they were treated as riding on their parent O atoms.

supplementary materials

Figures




Fig. 1. View of compound (I) with the atom-labeling scheme. Displacement ellipsoids are drawn at the 30% probability level. Only H atoms attached to water have been represented as small spheres of arbitrary radii. H bond is shown as dashed line.

Aqua[2,6-bis(2-pyridylamino)pyridine]sulfatonickel(II) monohydrate

Crystal data

$[\text{Ni}(\text{SO}_4)(\text{C}_{15}\text{H}_{13}\text{N}_5)(\text{H}_2\text{O})]\cdot\text{H}_2\text{O}$	$F_{000} = 936$
$M_r = 454.11$	$D_x = 1.771 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 7.3536 (8) \text{ \AA}$	Cell parameters from 3068 reflections
$b = 18.026 (2) \text{ \AA}$	$\theta = 2.0\text{--}25.2^\circ$
$c = 12.9125 (14) \text{ \AA}$	$\mu = 1.31 \text{ mm}^{-1}$
$\beta = 95.634 (2)^\circ$	$T = 298 (2) \text{ K}$
$V = 1703.3 (3) \text{ \AA}^3$	Block, green
$Z = 4$	$0.22 \times 0.16 \times 0.12 \text{ mm}$

Data collection

Bruker APEXII area-detector diffractometer	3068 independent reflections
Radiation source: fine-focus sealed tube	2821 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.014$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 25.2^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2004)	$h = -8\text{--}8$
$T_{\text{min}} = 0.762$, $T_{\text{max}} = 0.859$	$k = -20\text{--}21$
8650 measured reflections	$l = -15\text{--}13$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.056$	$w = 1/[s^2(F_{\text{o}}^2) + (0.0291P)^2 + 0.6456P]$

$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} < 0.001$
3068 reflections	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
259 parameters	$\Delta\rho_{\text{min}} = -0.32 \text{ e \AA}^{-3}$
8 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.26748 (3)	0.067232 (10)	0.233759 (14)	0.02550 (8)
N1	0.3089 (2)	0.06540 (7)	0.38831 (11)	0.0315 (3)
N2	0.2360 (2)	-0.06140 (8)	0.39796 (11)	0.0372 (4)
H2B	0.215 (3)	-0.0957 (9)	0.4422 (13)	0.045*
N3	0.28575 (18)	-0.04276 (7)	0.22027 (10)	0.0263 (3)
N4	0.3172 (2)	-0.03370 (8)	0.03833 (10)	0.0308 (3)
H4B	0.348 (3)	-0.0593 (9)	-0.0137 (11)	0.037*
N5	0.1828 (2)	0.07800 (7)	0.08526 (11)	0.0302 (3)
C1	0.3337 (3)	0.13148 (10)	0.43903 (14)	0.0397 (4)
H1	0.3680	0.1724	0.4015	0.048*
C2	0.3110 (3)	0.14091 (11)	0.54147 (15)	0.0464 (5)
H2	0.3247	0.1874	0.5725	0.056*
C3	0.2670 (3)	0.07959 (13)	0.59813 (15)	0.0525 (5)
H3	0.2530	0.0842	0.6687	0.063*
C4	0.2442 (3)	0.01220 (12)	0.55020 (14)	0.0468 (5)
H4	0.2156	-0.0295	0.5878	0.056*
C5	0.2643 (2)	0.00678 (9)	0.44353 (13)	0.0314 (4)
C6	0.2794 (2)	-0.08911 (9)	0.30303 (12)	0.0288 (3)
C7	0.3110 (3)	-0.16439 (9)	0.29762 (13)	0.0355 (4)
H7	0.2963	-0.1949	0.3543	0.043*
C8	0.3645 (3)	-0.19360 (9)	0.20725 (14)	0.0362 (4)
H8	0.3949	-0.2436	0.2037	0.043*
C9	0.3728 (2)	-0.14842 (9)	0.12190 (13)	0.0327 (4)
H9	0.4099	-0.1672	0.0603	0.039*
C10	0.3249 (2)	-0.07446 (8)	0.12950 (12)	0.0268 (3)

supplementary materials

C11	0.2342 (2)	0.03302 (9)	0.01132 (12)	0.0274 (3)
C12	0.2087 (3)	0.05246 (10)	-0.09421 (13)	0.0353 (4)
H12	0.2458	0.0206	-0.1448	0.042*
C13	0.1283 (3)	0.11901 (10)	-0.12178 (14)	0.0389 (4)
H13	0.1129	0.1334	-0.1913	0.047*
C14	0.0701 (2)	0.16493 (10)	-0.04534 (14)	0.0370 (4)
H14	0.0126	0.2098	-0.0628	0.044*
C15	0.0987 (2)	0.14286 (9)	0.05574 (14)	0.0351 (4)
H15	0.0590	0.1735	0.1069	0.042*
S1	0.55296 (6)	0.18214 (2)	0.18169 (3)	0.02779 (10)
O1	0.37943 (18)	0.16682 (6)	0.22894 (10)	0.0409 (3)
O2	0.5111 (2)	0.22657 (8)	0.08910 (10)	0.0496 (4)
O3	0.6762 (2)	0.22117 (8)	0.25801 (11)	0.0553 (4)
O4	0.63271 (18)	0.11012 (7)	0.15417 (10)	0.0417 (3)
O5	-0.02550 (18)	0.09435 (8)	0.27157 (10)	0.0466 (3)
H5A	-0.0394	0.1213	0.3252	0.070*
H5B	-0.1208	0.1008	0.2292	0.070*
O6	-0.1511 (2)	0.17329 (7)	0.44790 (10)	0.0473 (3)
H6A	-0.0976	0.2059	0.4872	0.071*
H6B	-0.2151	0.1947	0.3983	0.071*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.03692 (13)	0.01854 (12)	0.02107 (12)	0.00086 (8)	0.00302 (8)	-0.00001 (7)
N1	0.0354 (8)	0.0303 (8)	0.0283 (7)	0.0020 (6)	-0.0002 (6)	-0.0009 (6)
N2	0.0549 (10)	0.0298 (8)	0.0288 (8)	-0.0026 (7)	0.0136 (7)	0.0031 (6)
N3	0.0305 (7)	0.0233 (7)	0.0256 (7)	-0.0004 (5)	0.0051 (5)	0.0005 (5)
N4	0.0429 (9)	0.0260 (7)	0.0247 (7)	0.0043 (6)	0.0097 (6)	0.0005 (6)
N5	0.0371 (8)	0.0250 (7)	0.0289 (7)	0.0033 (6)	0.0048 (6)	0.0019 (6)
C1	0.0503 (11)	0.0323 (9)	0.0346 (9)	0.0003 (8)	-0.0053 (8)	-0.0044 (7)
C2	0.0556 (13)	0.0447 (11)	0.0373 (10)	0.0055 (9)	-0.0041 (9)	-0.0141 (9)
C3	0.0634 (14)	0.0641 (14)	0.0313 (10)	0.0005 (11)	0.0108 (9)	-0.0123 (10)
C4	0.0602 (13)	0.0507 (12)	0.0314 (10)	-0.0041 (10)	0.0140 (9)	0.0006 (8)
C5	0.0315 (9)	0.0335 (9)	0.0294 (8)	0.0029 (7)	0.0039 (7)	0.0002 (7)
C6	0.0316 (9)	0.0278 (8)	0.0275 (8)	-0.0018 (7)	0.0054 (6)	0.0018 (7)
C7	0.0469 (11)	0.0272 (9)	0.0327 (9)	-0.0019 (7)	0.0051 (8)	0.0063 (7)
C8	0.0451 (11)	0.0221 (8)	0.0413 (10)	0.0027 (7)	0.0032 (8)	0.0010 (7)
C9	0.0392 (10)	0.0270 (8)	0.0327 (9)	0.0022 (7)	0.0071 (7)	-0.0024 (7)
C10	0.0266 (8)	0.0259 (8)	0.0280 (8)	-0.0010 (6)	0.0039 (6)	0.0002 (6)
C11	0.0285 (8)	0.0256 (8)	0.0283 (8)	-0.0029 (6)	0.0041 (6)	0.0014 (6)
C12	0.0443 (11)	0.0342 (9)	0.0278 (8)	0.0003 (8)	0.0057 (7)	0.0002 (7)
C13	0.0459 (11)	0.0401 (10)	0.0299 (9)	0.0004 (8)	-0.0001 (8)	0.0079 (8)
C14	0.0385 (10)	0.0299 (9)	0.0414 (10)	0.0032 (7)	-0.0016 (8)	0.0071 (8)
C15	0.0401 (10)	0.0286 (9)	0.0366 (9)	0.0049 (7)	0.0037 (7)	0.0005 (7)
S1	0.0340 (2)	0.0242 (2)	0.02483 (19)	0.00013 (16)	0.00140 (16)	-0.00106 (15)
O1	0.0525 (8)	0.0268 (6)	0.0467 (7)	-0.0059 (5)	0.0217 (6)	-0.0056 (5)
O2	0.0642 (9)	0.0480 (8)	0.0378 (7)	0.0110 (7)	0.0114 (6)	0.0162 (6)

O3	0.0541 (9)	0.0483 (9)	0.0593 (9)	-0.0030 (7)	-0.0153 (7)	-0.0187 (7)
O4	0.0489 (8)	0.0356 (7)	0.0401 (7)	0.0116 (6)	0.0020 (6)	-0.0076 (5)
O5	0.0364 (7)	0.0672 (9)	0.0361 (7)	0.0131 (7)	0.0035 (5)	-0.0052 (6)
O6	0.0628 (9)	0.0392 (7)	0.0391 (7)	-0.0064 (6)	0.0004 (6)	-0.0034 (6)

Geometric parameters (\AA , $^\circ$)

Ni1—N5	1.9665 (14)	C4—H4	0.9300
Ni1—O1	1.9784 (12)	C6—C7	1.380 (2)
Ni1—N1	1.9892 (14)	C7—C8	1.373 (3)
Ni1—N3	1.9961 (14)	C7—H7	0.9300
Ni1—O5	2.3077 (13)	C8—C9	1.377 (2)
N1—C5	1.334 (2)	C8—H8	0.9300
N1—C1	1.363 (2)	C9—C10	1.385 (2)
N2—C5	1.370 (2)	C9—H9	0.9300
N2—C6	1.389 (2)	C11—C12	1.402 (2)
N2—H2B	0.866 (9)	C12—C13	1.369 (2)
N3—C10	1.360 (2)	C12—H12	0.9300
N3—C6	1.361 (2)	C13—C14	1.388 (3)
N4—C11	1.378 (2)	C13—H13	0.9300
N4—C10	1.384 (2)	C14—C15	1.361 (2)
N4—H4B	0.862 (9)	C14—H14	0.9300
N5—C11	1.335 (2)	C15—H15	0.9300
N5—C15	1.360 (2)	S1—O2	1.4467 (13)
C1—C2	1.360 (3)	S1—O3	1.4531 (14)
C1—H1	0.9300	S1—O4	1.4821 (12)
C2—C3	1.381 (3)	S1—O1	1.4932 (13)
C2—H2	0.9300	O5—H5A	0.8598
C3—C4	1.366 (3)	O5—H5B	0.8532
C3—H3	0.9300	O6—H6A	0.8477
C4—C5	1.403 (2)	O6—H6B	0.8488
N5—Ni1—O1	88.43 (6)	N3—C6—N2	120.04 (15)
N5—Ni1—N1	169.25 (6)	C7—C6—N2	116.92 (15)
O1—Ni1—N1	91.34 (6)	C8—C7—C6	118.96 (16)
N5—Ni1—N3	91.75 (5)	C8—C7—H7	120.5
O1—Ni1—N3	150.08 (5)	C6—C7—H7	120.5
N1—Ni1—N3	93.79 (5)	C7—C8—C9	119.51 (16)
N5—Ni1—O5	88.44 (5)	C7—C8—H8	120.2
O1—Ni1—O5	102.39 (5)	C9—C8—H8	120.2
N1—Ni1—O5	81.11 (5)	C8—C9—C10	118.76 (16)
N3—Ni1—O5	107.52 (5)	C8—C9—H9	120.6
C5—N1—C1	117.63 (15)	C10—C9—H9	120.6
C5—N1—Ni1	121.88 (11)	N3—C10—N4	121.05 (14)
C1—N1—Ni1	117.88 (11)	N3—C10—C9	122.84 (15)
C5—N2—C6	131.54 (15)	N4—C10—C9	116.11 (14)
C5—N2—H2B	112.7 (14)	N5—C11—N4	119.90 (14)
C6—N2—H2B	113.3 (14)	N5—C11—C12	121.54 (15)
C10—N3—C6	116.45 (14)	N4—C11—C12	118.55 (15)
C10—N3—Ni1	120.95 (10)	C13—C12—C11	119.02 (16)

supplementary materials

C6—N3—Ni1	122.24 (11)	C13—C12—H12	120.5
C11—N4—C10	131.19 (14)	C11—C12—H12	120.5
C11—N4—H4B	114.3 (13)	C12—C13—C14	119.56 (16)
C10—N4—H4B	112.8 (12)	C12—C13—H13	120.2
C11—N5—C15	118.33 (14)	C14—C13—H13	120.2
C11—N5—Ni1	123.55 (11)	C15—C14—C13	118.56 (16)
C15—N5—Ni1	116.74 (11)	C15—C14—H14	120.7
C2—C1—N1	123.59 (18)	C13—C14—H14	120.7
C2—C1—H1	118.2	N5—C15—C14	122.94 (16)
N1—C1—H1	118.2	N5—C15—H15	118.5
C1—C2—C3	118.20 (18)	C14—C15—H15	118.5
C1—C2—H2	120.9	O2—S1—O3	111.12 (9)
C3—C2—H2	120.9	O2—S1—O4	110.11 (8)
C4—C3—C2	119.76 (18)	O3—S1—O4	110.59 (8)
C4—C3—H3	120.1	O2—S1—O1	108.60 (8)
C2—C3—H3	120.1	O3—S1—O1	108.26 (9)
C3—C4—C5	119.10 (19)	O4—S1—O1	108.06 (7)
C3—C4—H4	120.4	S1—O1—Ni1	123.76 (7)
C5—C4—H4	120.4	Ni1—O5—H5A	118.5
N1—C5—N2	121.08 (15)	Ni1—O5—H5B	128.2
N1—C5—C4	121.67 (16)	H5A—O5—H5B	106.5
N2—C5—C4	117.25 (16)	H6A—O6—H6B	109.1
N3—C6—C7	123.03 (15)		

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N2—H2B···O6 ⁱ	0.866 (9)	2.079 (10)	2.9433 (19)	176 (2)
N4—H4B···O4 ⁱⁱ	0.862 (9)	2.050 (10)	2.8967 (18)	167.1 (18)
O5—H5A···O6	0.86	2.08	2.9112 (18)	163
O5—H5B···O4 ⁱⁱⁱ	0.85	1.98	2.8192 (19)	169
O6—H6A···O2 ^{iv}	0.85	1.91	2.7531 (18)	173
O6—H6B···O3 ⁱⁱⁱ	0.85	1.97	2.785 (2)	162

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+1, -y, -z$; (iii) $x-1, y, z$; (iv) $x-1/2, -y+1/2, z+1/2$.

Fig. 1

