

Bis{2-[(2-cyanophenyl)iminomethyl]-phenolato}copper(II)

Rong Xia, Hai-Jun Xu* and Han Wang

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: xuhj@seu.edu.cn


Received 5 August 2008; accepted 20 October 2008

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.050; wR factor = 0.125; data-to-parameter ratio = 16.4.

In the title mononuclear copper(II) complex, $[\text{Cu}(\text{C}_{14}\text{H}_9\text{N}_2\text{O})_2]$, the Cu^{II} atom, situated on an inversion centre, shows a slightly distorted square-planar geometry and is coordinated by the N and O atoms from two deprotonated symmetry-related Schiff base ligands. The $\text{Cu}-\text{N}$ and $\text{Cu}-\text{O}$ bond lengths are 2.009 (2) and 1.888 (2) \AA , respectively. The dihedral angle between the cyanophenyl rings and phenolate rings is 42.28 (13) $^\circ$.

Related literature

For Schiff base complexes with copper(II) and nickel(II), see: Gong *et al.* (2008); Kitaura *et al.* (2004); Marganian *et al.* (1995). For bond-length data, see: Jian *et al.* (2004); Ünver (2002). For a related structure, see: Xia *et al.* (2008).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{14}\text{H}_9\text{N}_2\text{O})_2]$	$V = 1148.0 (6)\text{ \AA}^3$
$M_r = 506.00$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.698 (3)\text{ \AA}$	$\mu = 0.99\text{ mm}^{-1}$
$b = 11.403 (3)\text{ \AA}$	$T = 293 (2)\text{ K}$
$c = 10.889 (3)\text{ \AA}$	$0.15 \times 0.10 \times 0.10\text{ mm}$
$\beta = 107.570 (11)^\circ$	

Data collection

Rigaku Mercury2 diffractometer	11307 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	2629 independent reflections
$T_{\min} = 0.909$, $T_{\max} = 1.000$	1952 reflections with $I > 2\sigma(I)$
(expected range = 0.824–0.906)	$R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	160 parameters
$wR(F^2) = 0.125$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\max} = 0.26\text{ e \AA}^{-3}$
2629 reflections	$\Delta\rho_{\min} = -0.43\text{ e \AA}^{-3}$

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

HJX acknowledges a start-up grant from Southeast University, People's Republic of China

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GW2050).

References

- Gong, X.-X., Xia, R. & Xu, H.-J. (2008). *Acta Cryst. E64*, m494.
- Jian, F.-F., Li, L., Sun, P.-P. & Xiao, H.-L. (2004). *Chin. J. Inorg. Chem. 20*, 1295–1298.
- Kitaura, R., Onoyama, G., Sakamoto, H., Matsuda, R., Noro, S. & Kitagawa, S. (2004). *Angew. Chem. Int. Ed.* **43**, 2684–2687.
- Marganian, C. A., Vazir, H., Baidya, N., Olmstead, M. M. & Mascharak, P. K. (1995). *J. Am. Chem. Soc.* **117**, 1584–1594.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Ünver, H. (2002). *J. Mol. Struct. 641*, 35–40.
- Xia, R., Xu, H.-J. & Gong, X.-X. (2008). *Acta Cryst. E64*, o1047.

supplementary materials

Acta Cryst. (2008). E64, m1539 [doi:10.1107/S160053680803417X]

Bis{2-[(2-cyanophenyl)iminomethyl]phenolato}copper(II)

R. Xia, H.-J. Xu and H. Wang

Comment

Metal derivatives of Schiff bases have been studied extensively, and copper(II) and nickel(II) complexes play an important role in both synthetic and structural research. These complexes have received much attention in recent years (Marganian *et al.*, 1995; Kitaura *et al.* 2004). We have reported previously the crystal structures of monomeric Schiff base complexes of Ni^{II} (Gong *et al.*, 2008). As a continuation of our research on the synthesis and structure of transition metal complexes of Schiff base compounds, we here report the results of the reaction of copper(II) with the didentate ligand 2-(2-cyanophenyliminomethyl)phenol, forming the title compound (I).

Fig. 1 shows the molecular structure of the title compound. The copper(II) is coordinated by the two imine N and two phenolate O atoms of the two Schiff base ligands in a slightly distorted square-planar geometry in a *trans* arrangement. 2-(2-cyanophenyliminomethyl)phenol loses a proton from the hydroxyl group and acts as a singly charged bidentate ligand coordinating to Copper(II) through the phenolate O and imine N atoms. The dihedral angle between the C1—C6 and C8—C13 benzene rings is 42.28 (0.13)°. The N1—Cu1—O1 bond angles is 90.78 (9)°. The two equivalent Cu—N and Cu—O distances are 2.009 (2) Å and 1.888 (2) Å, respectively. All these parameters conform to values in other square-planar-coordinated copper(II) compounds (Jian *et al.*, 2004; Ünver 2002).

Experimental

2-(2-Cyanophenyliminomethyl)phenol was prepared according to the literature (Xia *et al.*, 2008). CuCl₂·2H₂O (17. mg, 0.1 mmol) in methanol (5 ml) was added to the solution of 2-(2-cyanophenyliminomethyl)phenol (22.2 mg, 0.1 mmol) in the methanol (5 ml), pH of the mixture was adjusted to 8–9 and stirred for 4 h. The filtrate was kept at room temperature for about two weeks, and green block crystals of (I) for X-ray single-crystal investigation were obtained.

Refinement

All H atoms attached to C atoms were fixed geometrically and treated as riding with C—H = 0.93 Å and *U*_{iso}(H) = 1.2*U*_{eq}(C).

Figures




Fig. 1. Molecular view of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii. [Symmetry code: (i) 2 - *x*, 1 - *y*, -*z*]

supplementary materials

Bis{2-[(2-cyanophenyl)iminomethyl]phenolato}copper(II)

Crystal data

[Cu(C ₁₄ H ₉ N ₂ O) ₂]	$F_{000} = 518$
$M_r = 506.00$	$D_x = 1.464 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 9.698 (3) \text{ \AA}$	Cell parameters from 2373 reflections
$b = 11.403 (3) \text{ \AA}$	$\theta = 3.1\text{--}27.4^\circ$
$c = 10.889 (3) \text{ \AA}$	$\mu = 0.99 \text{ mm}^{-1}$
$\beta = 107.570 (11)^\circ$	$T = 293 (2) \text{ K}$
$V = 1148.0 (6) \text{ \AA}^3$	Block, red
$Z = 2$	$0.15 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Rigaku Mercury2 diffractometer	2629 independent reflections
Radiation source: fine-focus sealed tube	1952 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.057$
Detector resolution: 13.6612 pixels mm ⁻¹	$\theta_{\text{max}} = 27.5^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 3.1^\circ$
ω scans	$h = -12 \rightarrow 12$
Absorption correction: multi-scan (CrystalClear; Rigaku, 2005)	$k = -14 \rightarrow 14$
$T_{\text{min}} = 0.909$, $T_{\text{max}} = 1.000$	$l = -14 \rightarrow 14$
11307 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.050$	H-atom parameters constrained
$wR(F^2) = 0.125$	$w = 1/[\sigma^2(F_o^2) + (0.0516P)^2 + 0.2943P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2629 reflections	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
160 parameters	$\Delta\rho_{\text{min}} = -0.43 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	1.0000	0.5000	0.0000	0.03878 (18)
N1	0.9612 (2)	0.6674 (2)	0.0368 (2)	0.0383 (5)
O1	1.0810 (2)	0.46983 (18)	0.1778 (2)	0.0469 (5)
C8	0.8569 (3)	0.7359 (3)	-0.0563 (3)	0.0400 (6)
C1	1.1420 (3)	0.6686 (3)	0.2473 (3)	0.0416 (7)
C7	1.0377 (3)	0.7214 (3)	0.1397 (3)	0.0416 (7)
H7A	1.0234	0.8018	0.1439	0.050*
C9	0.7154 (3)	0.6925 (3)	-0.1043 (3)	0.0469 (7)
C2	1.1568 (3)	0.5450 (3)	0.2631 (3)	0.0403 (7)
C4	1.3310 (4)	0.5785 (3)	0.4717 (3)	0.0587 (9)
H4A	1.3938	0.5479	0.5471	0.070*
C6	1.2221 (3)	0.7432 (3)	0.3469 (3)	0.0535 (8)
H6A	1.2098	0.8239	0.3373	0.064*
C13	0.8922 (4)	0.8413 (3)	-0.1019 (3)	0.0507 (8)
H13A	0.9852	0.8715	-0.0698	0.061*
C14	0.6811 (3)	0.5801 (3)	-0.0602 (3)	0.0539 (8)
C10	0.6108 (4)	0.7564 (4)	-0.1967 (3)	0.0624 (10)
H10A	0.5162	0.7291	-0.2270	0.075*
C12	0.7880 (5)	0.9012 (3)	-0.1957 (4)	0.0670 (10)
H12A	0.8126	0.9711	-0.2280	0.080*
C5	1.3176 (4)	0.6996 (3)	0.4574 (3)	0.0601 (9)
H5A	1.3720	0.7496	0.5211	0.072*
C3	1.2547 (4)	0.5035 (3)	0.3782 (3)	0.0507 (8)
H3A	1.2678	0.4231	0.3909	0.061*
N2	0.6591 (4)	0.4905 (3)	-0.0242 (4)	0.0716 (9)
C11	0.6491 (5)	0.8599 (4)	-0.2422 (4)	0.0729 (12)
H11A	0.5805	0.9021	-0.3049	0.087*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0424 (3)	0.0348 (3)	0.0347 (3)	0.0037 (2)	0.0050 (2)	-0.0003 (2)

supplementary materials

N1	0.0399 (13)	0.0373 (13)	0.0349 (12)	0.0036 (10)	0.0069 (10)	0.0010 (10)
O1	0.0581 (14)	0.0381 (12)	0.0378 (12)	0.0047 (9)	0.0046 (10)	-0.0004 (8)
C8	0.0428 (16)	0.0400 (16)	0.0340 (15)	0.0077 (12)	0.0066 (12)	-0.0006 (12)
C1	0.0385 (16)	0.0445 (17)	0.0373 (16)	0.0018 (12)	0.0044 (12)	-0.0004 (12)
C7	0.0445 (17)	0.0332 (15)	0.0433 (16)	0.0020 (12)	0.0074 (13)	-0.0001 (12)
C9	0.0429 (18)	0.0526 (19)	0.0407 (17)	0.0068 (14)	0.0059 (13)	-0.0004 (14)
C2	0.0407 (17)	0.0443 (16)	0.0349 (15)	0.0040 (13)	0.0098 (13)	0.0009 (12)
C4	0.050 (2)	0.073 (3)	0.0432 (19)	0.0075 (17)	-0.0005 (15)	0.0027 (16)
C6	0.058 (2)	0.0493 (19)	0.0458 (19)	-0.0028 (15)	0.0038 (15)	-0.0048 (14)
C13	0.057 (2)	0.0476 (18)	0.0479 (19)	0.0047 (15)	0.0162 (15)	0.0054 (14)
C14	0.0386 (18)	0.071 (2)	0.0456 (19)	-0.0032 (16)	0.0030 (14)	-0.0054 (17)
C10	0.046 (2)	0.081 (3)	0.050 (2)	0.0174 (18)	-0.0011 (16)	0.0000 (18)
C12	0.092 (3)	0.051 (2)	0.055 (2)	0.0172 (19)	0.018 (2)	0.0155 (16)
C5	0.061 (2)	0.065 (2)	0.0416 (19)	-0.0074 (17)	-0.0045 (16)	-0.0060 (16)
C3	0.056 (2)	0.0516 (19)	0.0402 (17)	0.0096 (15)	0.0080 (14)	0.0061 (14)
N2	0.064 (2)	0.067 (2)	0.077 (2)	-0.0158 (16)	0.0115 (17)	0.0050 (17)
C11	0.079 (3)	0.073 (3)	0.055 (2)	0.036 (2)	0.003 (2)	0.0149 (19)

Geometric parameters (\AA , $^\circ$)

Cu1—O1 ⁱ	1.888 (2)	C4—C3	1.364 (4)
Cu1—O1	1.888 (2)	C4—C5	1.391 (5)
Cu1—N1 ⁱ	2.009 (2)	C4—H4A	0.9300
Cu1—N1	2.009 (2)	C6—C5	1.371 (4)
N1—C7	1.298 (3)	C6—H6A	0.9300
N1—C8	1.429 (3)	C13—C12	1.381 (4)
O1—C2	1.313 (4)	C13—H13A	0.9300
C8—C13	1.382 (4)	C14—N2	1.137 (4)
C8—C9	1.403 (4)	C10—C11	1.374 (5)
C1—C2	1.421 (4)	C10—H10A	0.9300
C1—C6	1.413 (4)	C12—C11	1.372 (5)
C1—C7	1.429 (4)	C12—H12A	0.9300
C7—H7A	0.9300	C5—H5A	0.9300
C9—C10	1.398 (4)	C3—H3A	0.9300
C9—C14	1.443 (5)	C11—H11A	0.9300
C2—C3	1.407 (4)		
O1 ⁱ —Cu1—O1	180.00 (12)	C3—C4—C5	121.9 (3)
O1 ⁱ —Cu1—N1 ⁱ	90.78 (9)	C3—C4—H4A	119.1
O1—Cu1—N1 ⁱ	89.22 (9)	C5—C4—H4A	119.1
O1 ⁱ —Cu1—N1	89.22 (9)	C5—C6—C1	121.7 (3)
O1—Cu1—N1	90.78 (9)	C5—C6—H6A	119.2
N1 ⁱ —Cu1—N1	180.00 (13)	C1—C6—H6A	119.2
C7—N1—C8	116.8 (2)	C8—C13—C12	119.4 (3)
C7—N1—Cu1	122.03 (19)	C8—C13—H13A	120.3
C8—N1—Cu1	120.84 (18)	C12—C13—H13A	120.3
C2—O1—Cu1	125.24 (19)	N2—C14—C9	177.6 (4)
C13—C8—C9	119.5 (3)	C11—C10—C9	119.4 (3)

supplementary materials

C13—C8—N1	122.1 (3)	C11—C10—H10A	120.3
C9—C8—N1	118.4 (3)	C9—C10—H10A	120.3
C2—C1—C6	119.5 (3)	C11—C12—C13	121.3 (4)
C2—C1—C7	122.5 (3)	C11—C12—H12A	119.3
C6—C1—C7	117.7 (3)	C13—C12—H12A	119.3
N1—C7—C1	125.9 (3)	C6—C5—C4	118.3 (3)
N1—C7—H7A	117.1	C6—C5—H5A	120.9
C1—C7—H7A	117.1	C4—C5—H5A	120.9
C8—C9—C10	120.0 (3)	C4—C3—C2	121.5 (3)
C8—C9—C14	119.1 (3)	C4—C3—H3A	119.3
C10—C9—C14	120.8 (3)	C2—C3—H3A	119.3
O1—C2—C3	119.6 (3)	C12—C11—C10	120.3 (3)
O1—C2—C1	123.2 (3)	C12—C11—H11A	119.9
C3—C2—C1	117.2 (3)	C10—C11—H11A	119.9

Symmetry codes: (i) $-x+2, -y+1, -z$.

supplementary materials

Fig. 1

