

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

2,2-Dimethyl-*N,N'*-bis(4-nitrobenzylidene)propane-1,3-diamineHoong-Kun Fun,<sup>a\*</sup> Hadi Kargar<sup>b‡</sup> and Reza Kia<sup>a</sup><sup>a</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>b</sup>Department of Chemistry, School of Science, Payame Noor University (PNU), Ardakan, Yazd, Iran

Correspondence e-mail: hkfun@usm.my


Received 1 November 2008; accepted 10 November 2008

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.061;  $wR$  factor = 0.152; data-to-parameter ratio = 27.8.

In the title compound,  $\text{C}_{19}\text{H}_{20}\text{N}_4\text{O}_4$ , a potential bidentate Schiff base ligand, each imino ( $\text{C}=\text{N}$ ) functional group is coplanar with the adjacent benzene ring. The two benzene rings form a dihedral angle of  $10.52(6)^\circ$ . Intermolecular  $\text{C}-\text{H}\cdots\text{O}$  contacts link neighbouring molecules into supra-molecular array with an  $R_2^2(32)$  ring motif and a  $\text{C}-\text{H}\cdots\pi$  contact is also present.

## Related literature

For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: Li *et al.* (2005); Bomfim *et al.* (2005); Glidewell *et al.* (2005, 2006); Sun *et al.* (2004); Fun *et al.* (2008a,b).



## Experimental

## Crystal data

$\text{C}_{19}\text{H}_{20}\text{N}_4\text{O}_4$   
 $M_r = 368.39$   
 Monoclinic,  $P2_1/c$   
 $a = 7.8219(1)$  Å  
 $b = 19.9716(4)$  Å

$c = 12.0125(3)$  Å  
 $\beta = 92.831(1)^\circ$   
 $V = 1874.25(6)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.09$  mm<sup>-1</sup>  
 $T = 100(1)$  K

0.45 × 0.19 × 0.08 mm

## Data collection

Bruker SMART APEXII CCD  
 area-detector diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2005)  
 $T_{\min} = 0.959$ ,  $T_{\max} = 0.993$

21845 measured reflections  
 6784 independent reflections  
 4307 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.048$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.152$   
 $S = 1.04$   
 6784 reflections

244 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.36$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.23$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}1-\text{H}1\text{A}\cdots\text{O}4^{\text{i}}$	0.93	2.52	3.4330 (18)	168
$\text{C}17-\text{H}17\text{A}\cdots\text{O}2^{\text{ii}}$	0.93	2.48	3.4063 (18)	171
$\text{C}19-\text{H}19\text{A}\cdots\text{C}g1^{\text{iii}}$	0.96	2.86	3.8058 (16)	171

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ ; (ii)  $-x+2, -y+1, -z+1$ ; (iii)  $x, -y-\frac{1}{2}, z-\frac{3}{2}$ . Cg1 is the centroid of the C12–C17 benzene ring.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for Science Fund grant No. 305/PFIZIK/613312. RK thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HK thanks PNU for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2324).

## References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.  
 Bomfim, J. A. S., Wardell, J. L., Low, J. N., Skakle, J. M. S. & Glidewell, C. (2005). *Acta Cryst.* **C61**, o53–o56.  
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Fun, H.-K., Kargar, H. & Kia, R. (2008a). *Acta Cryst.* **E64**, o1308.  
 Fun, H.-K., Kargar, H. & Kia, R. (2008b). *Acta Cryst.* **E64**, o2273.  
 Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2005). *Acta Cryst.* **E61**, o3551–o3553.  
 Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2006). *Acta Cryst.* **C62**, o1–o4.  
 Li, Y.-G., Zhu, H.-L., Chen, X.-Z. & Song, Y. (2005). *Acta Cryst.* **E61**, o4156–o4157.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.  
 Sun, Y.-X., You, Z.-L. & Zhu, H.-L. (2004). *Acta Cryst.* **E60**, o1707–o1708.

‡ Additional correspondence author, e-mail: hkargar@pnu.ac.ir.

**supplementary materials**

*Acta Cryst.* (2008). E64, o2335 [ doi:10.1107/S1600536808037033 ]

## 2,2-Dimethyl-*N,N'*-bis(4-nitrobenzylidene)propane-1,3-diamine

H.-K. Fun, H. Kargar and R. Kia

### Comment

Schiff bases are one of most prevalent mixed-donor ligands in the field of coordination chemistry. They play an important role in the development of coordination chemistry related to catalysis and enzymatic reactions, magnetism, and supramolecular architectures. Structures of Schiff bases derived from substituted benzaldehydes and closely related to the title compound, (I), have been reported previously (Li *et al.*, 2005; Bomfim *et al.*, 2005; Glidewell *et al.*, 2005, 2006; Sun *et al.*, 2004; Fun *et al.*, 2008*a,b*).

Each imino (C=N) functional group in (I), Fig. 1, is co-planar with the adjacent benzene ring. Intramolecular C—H $\cdots$ O contacts form five-membered rings, producing *S*(5) ring motifs (Bernstein *et al.*, 1995). The two benzene rings form a dihedral angle of 10.52 (6)°. Intermolecular C—H $\cdots$ O contacts link neighbouring molecules into supramolecular array with  $R^2_2(32)$  ring motif, Fig. 2 and Table 1. The crystal structure is further stabilized by weak C—H $\cdots$  $\pi$  interactions, ( Table 1.

### Experimental

The synthetic method has been described earlier (Fun *et al.*, 2008). Single crystals suitable for X-ray diffraction were obtained by evaporation of an ethanol solution of (I) held at room temperature.

### Refinement

All hydrogen atoms were positioned geometrically and refined using a riding model with C—H = 0.93 - 0.97 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2\text{-}1.5U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of (I) with atom labels and 50% displacement ellipsoids for non-H atoms.




Fig. 2. The crystal packing of (I), viewed down the *c*-axis showing the linking of neighbouring molecules via  $R^2_2(32)$  ring motifs. Intermolecular hydrogen bonds are shown as dashed lines.

## 2,2-Dimethyl-*N,N'*-bis(4-nitrobenzylidene)propane-1,3-diamine

### Crystal data

$C_{19}H_{20}N_4O_4$	$F_{000} = 776$
$M_r = 368.39$	$D_x = 1.306 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 7.8219 (1) \text{ \AA}$	Cell parameters from 3284 reflections
$b = 19.9716 (4) \text{ \AA}$	$\theta = 2.7\text{--}28.9^\circ$
$c = 12.0125 (3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 92.831 (1)^\circ$	$T = 100 (1) \text{ K}$
$V = 1874.25 (6) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.45 \times 0.19 \times 0.08 \text{ mm}$

### Data collection

Bruker SMART APEXII CCD area-detector diffractometer	6784 independent reflections
Radiation source: fine-focus sealed tube	4307 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.048$
$T = 100(1) \text{ K}$	$\theta_{\text{max}} = 32.6^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.959$ , $T_{\text{max}} = 0.993$	$k = -30 \rightarrow 24$
21845 measured reflections	$l = -16 \rightarrow 18$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.061$	H-atom parameters constrained
$wR(F^2) = 0.152$	$w = 1/[\sigma^2(F_o^2) + (0.063P)^2 + 0.2454P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
6784 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
244 parameters	$\Delta\rho_{\text{max}} = 0.36 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$
	Extinction correction: none

### Special details

**Experimental.** The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.36590 (14)	0.55914 (6)	0.45685 (10)	0.0335 (3)
O2	1.36634 (15)	0.48567 (6)	0.32431 (10)	0.0376 (3)
O3	0.04816 (14)	0.42804 (6)	1.17606 (9)	0.0317 (3)
O4	0.13096 (14)	0.52456 (5)	1.11560 (10)	0.0313 (3)
N1	0.91963 (15)	0.26572 (6)	0.65265 (11)	0.0229 (3)
N2	0.44695 (14)	0.27677 (6)	0.73309 (11)	0.0232 (3)
N3	1.32746 (15)	0.50399 (7)	0.41710 (12)	0.0263 (3)
N4	0.12292 (14)	0.46308 (6)	1.10932 (10)	0.0240 (3)
C1	1.07594 (17)	0.43692 (7)	0.64720 (12)	0.0214 (3)
H1A	1.0348	0.4512	0.7146	0.026*
C2	1.17251 (17)	0.48038 (7)	0.58500 (12)	0.0222 (3)
H2A	1.1983	0.5234	0.6103	0.027*
C3	1.22894 (16)	0.45767 (7)	0.48441 (12)	0.0212 (3)
C4	1.19437 (18)	0.39416 (8)	0.44352 (13)	0.0254 (3)
H4A	1.2337	0.3806	0.3753	0.031*
C5	1.09965 (17)	0.35135 (7)	0.50692 (13)	0.0245 (3)
H5A	1.0751	0.3083	0.4813	0.029*
C6	1.04054 (16)	0.37232 (7)	0.60928 (12)	0.0195 (3)
C7	0.94133 (16)	0.32672 (7)	0.67822 (12)	0.0205 (3)
H7A	0.8936	0.3432	0.7421	0.025*
C8	0.81845 (17)	0.22410 (7)	0.72454 (12)	0.0220 (3)
H8A	0.7833	0.2506	0.7871	0.026*
H8B	0.8885	0.1874	0.7537	0.026*
C9	0.65824 (17)	0.19558 (7)	0.66089 (12)	0.0211 (3)
C10	0.52727 (18)	0.25158 (7)	0.63434 (13)	0.0228 (3)
H10A	0.5843	0.2883	0.5986	0.027*
H10B	0.4391	0.2346	0.5821	0.027*
C11	0.44070 (16)	0.33980 (7)	0.74535 (12)	0.0211 (3)
H11A	0.4901	0.3673	0.6933	0.025*
C12	0.35718 (16)	0.37059 (7)	0.83998 (12)	0.0192 (3)
C13	0.25521 (16)	0.33267 (7)	0.90909 (12)	0.0214 (3)
H13A	0.2384	0.2873	0.8946	0.026*
C14	0.17916 (17)	0.36202 (7)	0.99872 (12)	0.0222 (3)
H14A	0.1125	0.3369	1.0453	0.027*
C15	0.20531 (16)	0.43004 (7)	1.01696 (12)	0.0203 (3)

## supplementary materials

---

C16	0.30321 (16)	0.46938 (7)	0.94913 (12)	0.0210 (3)
H16A	0.3173	0.5149	0.9627	0.025*
C17	0.37936 (17)	0.43876 (7)	0.86057 (12)	0.0215 (3)
H17A	0.4461	0.4641	0.8143	0.026*
C18	0.58134 (19)	0.14167 (8)	0.73376 (14)	0.0289 (3)
H18A	0.4811	0.1232	0.6959	0.043*
H18B	0.5507	0.1611	0.8031	0.043*
H18C	0.6639	0.1068	0.7481	0.043*
C19	0.70671 (19)	0.16493 (7)	0.54990 (13)	0.0259 (3)
H19A	0.6059	0.1477	0.5111	0.039*
H19B	0.7870	0.1292	0.5638	0.039*
H19C	0.7577	0.1987	0.5052	0.039*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0318 (6)	0.0268 (6)	0.0423 (7)	-0.0069 (5)	0.0043 (5)	0.0051 (5)
O2	0.0377 (6)	0.0447 (7)	0.0314 (7)	-0.0070 (5)	0.0128 (5)	0.0041 (5)
O3	0.0332 (6)	0.0354 (6)	0.0276 (6)	-0.0072 (5)	0.0108 (5)	-0.0019 (5)
O4	0.0381 (6)	0.0230 (6)	0.0334 (7)	0.0044 (4)	0.0082 (5)	-0.0026 (5)
N1	0.0213 (5)	0.0232 (6)	0.0244 (7)	-0.0026 (4)	0.0020 (5)	-0.0003 (5)
N2	0.0211 (5)	0.0225 (6)	0.0261 (7)	0.0004 (4)	0.0023 (5)	-0.0012 (5)
N3	0.0185 (5)	0.0297 (7)	0.0308 (7)	-0.0009 (5)	0.0019 (5)	0.0075 (6)
N4	0.0202 (5)	0.0280 (7)	0.0238 (7)	0.0009 (5)	0.0024 (5)	-0.0016 (5)
C1	0.0197 (6)	0.0234 (7)	0.0211 (7)	0.0017 (5)	0.0010 (5)	-0.0015 (6)
C2	0.0197 (6)	0.0212 (7)	0.0255 (8)	-0.0003 (5)	-0.0016 (5)	-0.0005 (6)
C3	0.0162 (6)	0.0235 (7)	0.0240 (7)	0.0008 (5)	0.0021 (5)	0.0055 (6)
C4	0.0255 (7)	0.0286 (8)	0.0229 (8)	-0.0007 (6)	0.0076 (6)	-0.0018 (6)
C5	0.0254 (7)	0.0228 (7)	0.0257 (8)	-0.0011 (5)	0.0051 (6)	-0.0037 (6)
C6	0.0173 (6)	0.0204 (6)	0.0207 (7)	0.0001 (5)	0.0011 (5)	0.0007 (5)
C7	0.0187 (6)	0.0234 (7)	0.0196 (7)	-0.0003 (5)	0.0018 (5)	-0.0004 (6)
C8	0.0239 (6)	0.0208 (7)	0.0215 (7)	-0.0013 (5)	0.0020 (5)	0.0012 (6)
C9	0.0218 (6)	0.0177 (6)	0.0238 (7)	-0.0015 (5)	0.0013 (5)	-0.0003 (6)
C10	0.0241 (6)	0.0216 (7)	0.0227 (8)	0.0002 (5)	0.0002 (5)	-0.0019 (6)
C11	0.0195 (6)	0.0219 (7)	0.0220 (7)	-0.0001 (5)	0.0018 (5)	0.0010 (6)
C12	0.0170 (6)	0.0200 (6)	0.0204 (7)	0.0021 (5)	-0.0001 (5)	0.0015 (5)
C13	0.0195 (6)	0.0179 (6)	0.0268 (8)	-0.0007 (5)	0.0015 (5)	0.0002 (6)
C14	0.0192 (6)	0.0229 (7)	0.0248 (8)	0.0000 (5)	0.0036 (5)	0.0046 (6)
C15	0.0180 (6)	0.0228 (7)	0.0201 (7)	0.0024 (5)	0.0010 (5)	-0.0002 (5)
C16	0.0207 (6)	0.0180 (6)	0.0242 (7)	0.0002 (5)	0.0007 (5)	0.0003 (6)
C17	0.0207 (6)	0.0211 (7)	0.0229 (7)	-0.0005 (5)	0.0027 (5)	0.0031 (6)
C18	0.0282 (7)	0.0228 (7)	0.0357 (9)	-0.0041 (6)	0.0025 (6)	0.0048 (7)
C19	0.0275 (7)	0.0212 (7)	0.0289 (8)	0.0011 (5)	-0.0008 (6)	-0.0045 (6)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—N3	1.2319 (17)	C8—H8B	0.9700
O2—N3	1.2256 (17)	C9—C18	1.530 (2)
O3—N4	1.2330 (15)	C9—C19	1.532 (2)

O4—N4	1.2316 (16)	C9—C10	1.539 (2)
N1—C7	1.2657 (18)	C10—H10A	0.9700
N1—C8	1.4597 (17)	C10—H10B	0.9700
N2—C11	1.2687 (18)	C11—C12	1.4732 (19)
N2—C10	1.4591 (18)	C11—H11A	0.9300
N3—C3	1.4714 (18)	C12—C17	1.3931 (19)
N4—C15	1.4671 (18)	C12—C13	1.4014 (18)
C1—C6	1.3914 (19)	C13—C14	1.3857 (19)
C1—C2	1.3921 (19)	C13—H13A	0.9300
C1—H1A	0.9300	C14—C15	1.389 (2)
C2—C3	1.383 (2)	C14—H14A	0.9300
C2—H2A	0.9300	C15—C16	1.3890 (18)
C3—C4	1.382 (2)	C16—C17	1.3868 (19)
C4—C5	1.3840 (19)	C16—H16A	0.9300
C4—H4A	0.9300	C17—H17A	0.9300
C5—C6	1.3990 (19)	C18—H18A	0.9600
C5—H5A	0.9300	C18—H18B	0.9600
C6—C7	1.4768 (18)	C18—H18C	0.9600
C7—H7A	0.9300	C19—H19A	0.9600
C8—C9	1.544 (2)	C19—H19B	0.9600
C8—H8A	0.9700	C19—H19C	0.9600
C7—N1—C8	118.34 (12)	C10—C9—C8	110.49 (11)
C11—N2—C10	117.20 (12)	N2—C10—C9	113.01 (12)
O2—N3—O1	123.55 (13)	N2—C10—H10A	109.0
O2—N3—C3	118.19 (13)	C9—C10—H10A	109.0
O1—N3—C3	118.26 (13)	N2—C10—H10B	109.0
O4—N4—O3	123.39 (12)	C9—C10—H10B	109.0
O4—N4—C15	118.14 (12)	H10A—C10—H10B	107.8
O3—N4—C15	118.47 (12)	N2—C11—C12	121.65 (13)
C6—C1—C2	120.46 (13)	N2—C11—H11A	119.2
C6—C1—H1A	119.8	C12—C11—H11A	119.2
C2—C1—H1A	119.8	C17—C12—C13	119.50 (12)
C3—C2—C1	118.00 (13)	C17—C12—C11	119.26 (12)
C3—C2—H2A	121.0	C13—C12—C11	121.24 (12)
C1—C2—H2A	121.0	C14—C13—C12	120.69 (13)
C4—C3—C2	123.10 (13)	C14—C13—H13A	119.7
C4—C3—N3	118.70 (13)	C12—C13—H13A	119.7
C2—C3—N3	118.18 (13)	C13—C14—C15	118.07 (12)
C3—C4—C5	118.16 (13)	C13—C14—H14A	121.0
C3—C4—H4A	120.9	C15—C14—H14A	121.0
C5—C4—H4A	120.9	C16—C15—C14	122.80 (13)
C4—C5—C6	120.52 (14)	C16—C15—N4	117.69 (12)
C4—C5—H5A	119.7	C14—C15—N4	119.48 (12)
C6—C5—H5A	119.7	C17—C16—C15	118.07 (13)
C1—C6—C5	119.75 (13)	C17—C16—H16A	121.0
C1—C6—C7	119.33 (12)	C15—C16—H16A	121.0
C5—C6—C7	120.92 (13)	C16—C17—C12	120.85 (12)
N1—C7—C6	121.69 (13)	C16—C17—H17A	119.6
N1—C7—H7A	119.2	C12—C17—H17A	119.6

## supplementary materials

C6—C7—H7A	119.2	C9—C18—H18A	109.5
N1—C8—C9	111.56 (12)	C9—C18—H18B	109.5
N1—C8—H8A	109.3	H18A—C18—H18B	109.5
C9—C8—H8A	109.3	C9—C18—H18C	109.5
N1—C8—H8B	109.3	H18A—C18—H18C	109.5
C9—C8—H8B	109.3	H18B—C18—H18C	109.5
H8A—C8—H8B	108.0	C9—C19—H19A	109.5
C18—C9—C19	109.76 (12)	C9—C19—H19B	109.5
C18—C9—C10	110.70 (11)	H19A—C19—H19B	109.5
C19—C9—C10	107.51 (12)	C9—C19—H19C	109.5
C18—C9—C8	107.98 (12)	H19A—C19—H19C	109.5
C19—C9—C8	110.41 (11)	H19B—C19—H19C	109.5
C6—C1—C2—C3	1.0 (2)	C11—N2—C10—C9	131.95 (13)
C1—C2—C3—C4	-0.2 (2)	C18—C9—C10—N2	49.59 (16)
C1—C2—C3—N3	178.53 (12)	C19—C9—C10—N2	169.47 (11)
O2—N3—C3—C4	4.2 (2)	C8—C9—C10—N2	-69.98 (14)
O1—N3—C3—C4	-176.02 (13)	C10—N2—C11—C12	177.85 (12)
O2—N3—C3—C2	-174.58 (13)	N2—C11—C12—C17	168.88 (14)
O1—N3—C3—C2	5.17 (19)	N2—C11—C12—C13	-11.5 (2)
C2—C3—C4—C5	-0.4 (2)	C17—C12—C13—C14	-1.2 (2)
N3—C3—C4—C5	-179.16 (13)	C11—C12—C13—C14	179.15 (13)
C3—C4—C5—C6	0.2 (2)	C12—C13—C14—C15	0.8 (2)
C2—C1—C6—C5	-1.2 (2)	C13—C14—C15—C16	0.3 (2)
C2—C1—C6—C7	178.54 (12)	C13—C14—C15—N4	178.38 (13)
C4—C5—C6—C1	0.6 (2)	O4—N4—C15—C16	6.84 (19)
C4—C5—C6—C7	-179.19 (13)	O3—N4—C15—C16	-173.73 (12)
C8—N1—C7—C6	-179.39 (12)	O4—N4—C15—C14	-171.35 (13)
C1—C6—C7—N1	-173.22 (14)	O3—N4—C15—C14	8.09 (19)
C5—C6—C7—N1	6.5 (2)	C14—C15—C16—C17	-0.9 (2)
C7—N1—C8—C9	118.97 (14)	N4—C15—C16—C17	-179.01 (12)
N1—C8—C9—C18	168.26 (11)	C15—C16—C17—C12	0.4 (2)
N1—C8—C9—C19	48.27 (15)	C13—C12—C17—C16	0.6 (2)
N1—C8—C9—C10	-70.53 (14)	C11—C12—C17—C16	-179.76 (13)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1A $\cdots$ O4 <sup>i</sup>	0.93	2.52	3.4330 (18)	168
C17—H17A $\cdots$ O2 <sup>ii</sup>	0.93	2.48	3.4063 (18)	171
C19—H19A $\cdots$ Cg1 <sup>iii</sup>	0.96	2.86	3.8058 (16)	171

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ ; (ii)  $-x+2, -y+1, -z+1$ ; (iii)  $x, -y-1/2, z-3/2$ .



Fig. 1




Fig. 2

