

## Naphthalene-2,3-diylbis[(2-thienyl)-methanone]

S. Thenmozhi,<sup>a</sup> A. SubbiahPandi,<sup>a\*</sup> S. Ranjith,<sup>a</sup> J. Arul Clement<sup>b</sup> and A. K. MohanaKrishnan<sup>b</sup>

<sup>a</sup>Department of Physics, Presidency College (Autonomous), Chennai 600 005, India, and <sup>b</sup>Department of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India

Correspondence e-mail: a\_spandian@yahoo.com


Received 15 October 2008; accepted 19 November 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.112; data-to-parameter ratio = 19.8.

The asymmetric unit of the title compound,  $\text{C}_{20}\text{H}_{12}\text{O}_2\text{S}_2$ , contains two crystallographically independent molecules which differ in the orientations of thienylmethanone units with respect to the naphthalene ring system [dihedral angles of 65.30 (11) and 50.94 (11)° in one molecule, 41.94 (12) and 69.61 (13)° in the other]. The crystal structure is stabilized by  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\pi$  interactions.

### Related literature

For a related structure, see: Labat & Halfpenny (2005). For general background, see: Pellis & West (1968); Cohen *et al.* (1977); Csaszar & Morvay (1983); Lakshmi *et al.* (1985); EI-Maghraby *et al.* (1984); Dzhurayev *et al.* (1992); Gewald *et al.* (1996); Jones *et al.* (1984); Palani *et al.* (2006).



### Experimental

#### Crystal data

$\text{C}_{20}\text{H}_{12}\text{O}_2\text{S}_2$	$V = 1680.69$ (10) Å <sup>3</sup>
$M_r = 348.42$	$Z = 4$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 9.7638$ (3) Å	$\mu = 0.33$ mm <sup>-1</sup>
$b = 11.1418$ (4) Å	$T = 293$ (2) K
$c = 15.4496$ (6) Å	$0.21 \times 0.19 \times 0.16$ mm
$\beta = 90.266$ (1)°	

#### Data collection

Bruker APEXII CCD area-detector diffractometer	21289 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	8580 independent reflections
$T_{\min} = 0.800$ , $T_{\max} = 0.950$	6591 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.024$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.112$	$\Delta\rho_{\text{max}} = 0.44$ e Å <sup>-3</sup>
$S = 1.01$	$\Delta\rho_{\text{min}} = -0.32$ e Å <sup>-3</sup>
8580 reflections	Absolute structure: Flack (1983),
433 parameters	3970 Friedel pairs
1 restraint	Flack parameter: 0.02 (5)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C4}-\text{H4}\cdots\text{O2}^{\text{i}}$	0.93	2.53	3.407 (3)	158
$\text{C13}-\text{H13}\cdots\text{Cg1}^{\text{ii}}$	0.93	2.86	3.737 (2)	157

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + 1$ ; (ii)  $-x + 1, y - \frac{1}{2}, -z + 1$ . Cg1 is the S2/C17-C20 ring centroid.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2003).

ST and ASP thank Dr J. Jothi Kumar, Principal of Presidency College (Autonomous), Chennai, for providing computer and internet facilities. Dr Babu Vargheese, SAIF, IIT, Madras, India, is thanked for his help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2690).

### References

- Bruker (2004). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cohen, V. I., Rist, N. & Duponchel, C. (1977). *J. Pharm. Sci.* **66**, 1322–1334.
- Csaszar, J. & Morvay, J. (1983). *Acta Pharm. Hung.* **53**, 121–128.
- Dzhurayev, A. D., Karimkulov, K. M., Makhsumov, A. G. & Amanov, N. (1992). *Khim. Form. Zh.* **26**, 73–75.
- EI-Maghraby, A. A., Haroun, B. & Mohammed, N. A. (1984). *Egypt. J. Pharm. Sci.* **23**, 327–336.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gewald, K., Schinke, E. & Botcher, H. (1996). *Chem. Ber.* **99**, 99–100.
- Jones, C. D., Jevnikar, M. G., Pike, A. J., Peters, M. K., Black, L. J., Thompson, A. R., Falcone, J. F. & Clemens, J. A. (1984). *J. Med. Chem.* **27**, 1057–1066.
- Labat, G. & Halfpenny, J. (2005). *Acta Cryst.* **E61**, o2813–o2814.
- Lakshmi, V. V., Sridhar, P. & Polasa, H. (1985). *Indian J. Pharm. Sci.* **47**, 202–204.
- Palani, K., Amaladass, P., Mohanakrishnan, A. K. & Ponnuswamy, M. N. (2006). *Acta Cryst.* **E62**, o49–o51.
- Pellis, G. & West, G. B. (1968). *Progress in Medicinal Chemistry*, Vol. 5, pp. 320–324. London: Butterworth & Co. Ltd.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

**supplementary materials**

*Acta Cryst.* (2008). E64, o2432 [ doi:10.1107/S1600536808038701 ]

## Naphthalene-2,3-diylbis[(2-thienyl)methanone]

S. Thenmozhi, A. SubbiahPandi, S. Ranjith, J. A. Clement and A. K. MohanaKrishnan

### Comment

Sulfur containing Schiff bases (Pellis & West, 1968; Cohen *et al.*, 1977; Csaszar & Morvay, 1983; Lakshmi *et al.*, 1985), and their thiophene derivatives (EI-Maghraby *et al.*, 1984; Dzhurayev *et al.*, 1992), possess pharmacological activities such as anti-bacterial, anti-cancer, anti-inflammatory and anti-toxic properties (Gewald *et al.*, 1996). Benzo(*b*)thiophene analogs have been shown to possess interesting estrogenic and antiestrogenic effects. Some of these compounds inhibit estradiol with greater potency than tamoxifen, and inhibition of the growth of DMBA induced mammary tumors by such compounds has been reported (Jones *et al.*, 1984). Some of the thiophene derivatives were screened against gram-positive, gram-negative bacteria and have shown promising anti-bacterial activity (Palani *et al.*, 2006). In view of this biological importance, the crystal structure of the title compound has been determined and the results are presented here.

The asymmetric unit of the title compound contains two crystallographically independent molecules (Fig. 1). The corresponding bond lengths and angles of the two molecules agree with each other, and are comparable to those observed in the structure of 1,5-bis(3-thienyloxy)-3-oxapentane (Labat & Halfpenny, 2005). The two independent molecules differ in the orientations of thienylmethanone units with respect to the naphthalene ring system. The S1/C12-C15 and S2/C17-C20 rings form dihedral angles of 65.30 (11)° and 50.94 (11)°, respectively, with the C1-C10 naphthalene ring system, whereas, the S1'/C12'-C15' and S2'/C17'-C20' rings form dihedral angles of 41.94 (12)° and 69.61 (13)°, respectively, with the C1'-C10' naphthalene ring system.

The crystal structure is stabilized by C–H···O and C–H··· $\pi$  interactions (Table 1 and Fig.2).

### Experimental

A mixture of phthalaldehyde (1 g, 7.46 mmol) and 1,4-di(thiophene-2-yl)butane-1,4-diane (1.86 g 7.46 mmol) was dissolved in ethanol and tertiary-butane oxide (2.08 g, 18.6 mmol), and then allowed to stir for 4 h at room temperature to get the product. The crude product was filtered and then recrystallized in chloroform. Single crystals suitable for X-ray diffraction were obtained by slow evaporation of a chloroform solution at room temperature.

### Refinement

All H atoms were fixed geometrically and allowed to ride on their parent C atoms, with C–H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

## Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. The crystal packing of the title compound, viewed down the *b* axis. Hydrogen bonds are shown as dashed lines.

## Naphthalene-2,3-diylbis[(2-thienyl)methanone]

### Crystal data

$C_{20}H_{12}O_2S_2$

$M_r = 348.42$

Monoclinic,  $P2_1$

Hall symbol: P 2yb

$a = 9.7638$  (3) Å

$b = 11.1418$  (4) Å

$c = 15.4496$  (6) Å

$\beta = 90.266$  (1)°

$V = 1680.69$  (10) Å<sup>3</sup>

$Z = 4$

$F_{000} = 720$

$D_x = 1.377$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 8580 reflections

$\theta = 1.3$ – $28.9$ °

$\mu = 0.33$  mm<sup>-1</sup>

$T = 293$  (2) K

Block, colourless

$0.21 \times 0.19 \times 0.16$  mm

### Data collection

Bruker APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

$\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.800$ ,  $T_{\max} = 0.950$

21289 measured reflections

8580 independent reflections

6591 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\text{max}} = 28.9$ °

$\theta_{\text{min}} = 1.3$ °

$h = -13 \rightarrow 13$

$k = -15 \rightarrow 14$

$l = -20 \rightarrow 17$

Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0499P)^2 + 0.3545P]$
$wR(F^2) = 0.112$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\max} = 0.001$
8580 reflections	$\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$
433 parameters	$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 3970 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.02 (5)

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.06339 (8)	-0.02637 (6)	0.45118 (6)	0.0657 (2)
S2	0.33739 (8)	0.44467 (8)	0.69777 (5)	0.0644 (2)
O1	0.0949 (2)	0.24092 (19)	0.44816 (17)	0.0822 (7)
O2	0.3446 (3)	0.24071 (18)	0.57466 (14)	0.0815 (7)
C1	0.4852 (3)	0.3351 (2)	0.18008 (17)	0.0542 (6)
H1	0.4468	0.2785	0.1429	0.065*
C2	0.5757 (3)	0.4164 (3)	0.14849 (18)	0.0609 (7)
H2	0.5984	0.4148	0.0901	0.073*
C3	0.6350 (3)	0.5021 (3)	0.20256 (18)	0.0571 (6)
H3	0.6966	0.5574	0.1800	0.069*
C4	0.6032 (2)	0.5053 (2)	0.28836 (17)	0.0488 (6)
H4	0.6430	0.5631	0.3239	0.059*
C5	0.5102 (2)	0.42129 (19)	0.32380 (15)	0.0398 (5)
C6	0.4485 (2)	0.3350 (2)	0.26773 (16)	0.0421 (5)
C7	0.3516 (2)	0.2545 (2)	0.30259 (16)	0.0451 (5)
H7	0.3127	0.1966	0.2668	0.054*

## supplementary materials

---

C8	0.3135 (2)	0.25942 (19)	0.38716 (16)	0.0434 (5)
C9	0.3768 (2)	0.3442 (2)	0.44373 (16)	0.0442 (5)
C10	0.4727 (2)	0.4227 (2)	0.41136 (15)	0.0428 (5)
H10	0.5137	0.4779	0.4484	0.051*
C11	0.1943 (3)	0.1879 (2)	0.42024 (19)	0.0508 (6)
C12	0.2010 (2)	0.0578 (2)	0.41656 (16)	0.0448 (5)
C13	0.3071 (2)	-0.0134 (2)	0.39437 (17)	0.0484 (6)
H13	0.3903	0.0156	0.3741	0.058*
C14	0.2781 (3)	-0.1368 (2)	0.4053 (2)	0.0608 (7)
H14	0.3397	-0.1980	0.3929	0.073*
C15	0.1515 (3)	-0.1553 (3)	0.4355 (2)	0.0653 (8)
H15	0.1158	-0.2310	0.4467	0.078*
C16	0.3484 (3)	0.3376 (2)	0.53847 (17)	0.0506 (6)
C17	0.3315 (2)	0.4492 (2)	0.58672 (15)	0.0456 (5)
C18	0.3095 (3)	0.5646 (2)	0.55646 (19)	0.0521 (6)
H18	0.3026	0.5854	0.4983	0.062*
C19	0.2991 (3)	0.6466 (3)	0.6262 (2)	0.0696 (8)
H19	0.2841	0.7284	0.6186	0.084*
C20	0.3127 (3)	0.5957 (3)	0.7037 (2)	0.0722 (9)
H20	0.3091	0.6381	0.7555	0.087*
S1'	0.36488 (7)	0.50071 (8)	0.93775 (5)	0.0670 (2)
S2'	1.03585 (9)	0.72925 (8)	0.94344 (7)	0.0795 (3)
O1'	0.6344 (2)	0.55500 (17)	0.87437 (16)	0.0701 (6)
O2'	0.8753 (3)	0.5098 (2)	0.99329 (14)	0.0745 (6)
C1'	0.7926 (3)	0.0907 (2)	0.7211 (2)	0.0694 (9)
H1'	0.7053	0.0575	0.7197	0.083*
C2'	0.8959 (4)	0.0362 (2)	0.6773 (2)	0.0730 (9)
H2'	0.8777	-0.0319	0.6445	0.088*
C3'	1.0280 (3)	0.0812 (2)	0.6812 (2)	0.0677 (8)
H3'	1.0981	0.0421	0.6521	0.081*
C4'	1.0559 (3)	0.1833 (2)	0.7278 (2)	0.0597 (7)
H4'	1.1449	0.2130	0.7300	0.072*
C5'	0.9510 (3)	0.2433 (2)	0.77216 (17)	0.0469 (6)
C6'	0.8160 (3)	0.1970 (2)	0.76870 (18)	0.0498 (6)
C7'	0.7096 (3)	0.2617 (2)	0.80915 (18)	0.0526 (6)
H7'	0.6207	0.2320	0.8061	0.063*
C8'	0.7328 (2)	0.3666 (2)	0.85265 (17)	0.0450 (5)
C9'	0.8698 (2)	0.40998 (19)	0.85839 (16)	0.0433 (5)
C10'	0.9736 (3)	0.3506 (2)	0.81877 (18)	0.0491 (6)
H10'	1.0620	0.3814	0.8224	0.059*
C11'	0.6203 (3)	0.4464 (2)	0.88117 (17)	0.0490 (6)
C12'	0.4927 (3)	0.3989 (2)	0.91630 (17)	0.0484 (6)
C13'	0.4546 (3)	0.2823 (3)	0.93971 (18)	0.0561 (7)
H13'	0.5098	0.2148	0.9340	0.067*
C14'	0.3192 (3)	0.2820 (3)	0.9735 (2)	0.0713 (9)
H14'	0.2752	0.2126	0.9921	0.086*
C15'	0.2605 (3)	0.3915 (3)	0.9763 (2)	0.0723 (9)
H15'	0.1727	0.4056	0.9970	0.087*
C16'	0.9038 (3)	0.5142 (2)	0.91669 (18)	0.0495 (6)

C17'	0.9723 (2)	0.6167 (2)	0.87875 (19)	0.0511 (6)
C18'	0.9843 (3)	0.6463 (2)	0.7926 (2)	0.0577 (7)
H18'	0.9581	0.5964	0.7472	0.069*
C19'	1.0413 (4)	0.7619 (3)	0.7816 (3)	0.0902 (12)
H19'	1.0548	0.7975	0.7278	0.108*
C20'	1.0736 (4)	0.8144 (3)	0.8562 (3)	0.0952 (14)
H20'	1.1130	0.8902	0.8600	0.114*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0555 (4)	0.0583 (4)	0.0835 (5)	-0.0129 (3)	0.0201 (4)	-0.0068 (4)
S2	0.0706 (5)	0.0755 (5)	0.0471 (4)	-0.0149 (4)	0.0070 (3)	-0.0040 (4)
O1	0.0665 (13)	0.0542 (12)	0.126 (2)	0.0072 (10)	0.0366 (13)	-0.0069 (13)
O2	0.135 (2)	0.0491 (11)	0.0603 (13)	0.0066 (12)	0.0098 (13)	0.0124 (10)
C1	0.0642 (17)	0.0556 (15)	0.0428 (15)	-0.0048 (12)	-0.0067 (12)	-0.0048 (12)
C2	0.0709 (19)	0.0683 (18)	0.0437 (15)	-0.0089 (15)	0.0013 (14)	0.0006 (13)
C3	0.0580 (15)	0.0573 (15)	0.0562 (16)	-0.0080 (13)	0.0039 (13)	0.0054 (13)
C4	0.0466 (13)	0.0469 (12)	0.0527 (15)	-0.0022 (11)	-0.0047 (11)	-0.0053 (11)
C5	0.0392 (11)	0.0353 (10)	0.0448 (13)	0.0058 (9)	-0.0047 (10)	-0.0025 (9)
C6	0.0440 (12)	0.0368 (10)	0.0455 (14)	0.0039 (9)	-0.0051 (10)	-0.0031 (9)
C7	0.0466 (13)	0.0396 (11)	0.0491 (15)	-0.0017 (9)	-0.0061 (11)	-0.0072 (10)
C8	0.0438 (12)	0.0333 (10)	0.0530 (15)	0.0032 (9)	-0.0011 (11)	-0.0019 (10)
C9	0.0478 (13)	0.0389 (11)	0.0460 (14)	0.0070 (10)	-0.0008 (11)	-0.0008 (10)
C10	0.0462 (12)	0.0382 (11)	0.0439 (13)	0.0019 (9)	-0.0068 (10)	-0.0060 (10)
C11	0.0481 (14)	0.0435 (12)	0.0610 (17)	0.0015 (10)	0.0054 (12)	-0.0060 (11)
C12	0.0433 (13)	0.0437 (11)	0.0474 (14)	-0.0074 (10)	-0.0008 (11)	-0.0020 (10)
C13	0.0421 (12)	0.0449 (12)	0.0582 (16)	-0.0006 (10)	-0.0013 (11)	0.0002 (11)
C14	0.0656 (18)	0.0427 (13)	0.074 (2)	-0.0007 (12)	-0.0019 (15)	-0.0039 (13)
C15	0.074 (2)	0.0439 (13)	0.078 (2)	-0.0143 (13)	0.0014 (16)	-0.0052 (14)
C16	0.0563 (15)	0.0479 (13)	0.0477 (15)	-0.0002 (11)	0.0008 (12)	0.0014 (11)
C17	0.0433 (12)	0.0505 (13)	0.0429 (13)	-0.0061 (10)	0.0021 (10)	-0.0061 (11)
C18	0.0522 (15)	0.0505 (13)	0.0536 (16)	-0.0017 (11)	0.0036 (12)	-0.0111 (12)
C19	0.082 (2)	0.0509 (15)	0.076 (2)	-0.0088 (14)	0.0077 (17)	-0.0145 (15)
C20	0.085 (2)	0.0714 (19)	0.061 (2)	-0.0177 (17)	0.0105 (17)	-0.0213 (16)
S1'	0.0518 (4)	0.0751 (5)	0.0741 (5)	0.0064 (4)	0.0049 (3)	0.0109 (4)
S2'	0.0642 (5)	0.0705 (5)	0.1039 (7)	-0.0178 (4)	0.0145 (4)	-0.0433 (5)
O1'	0.0608 (12)	0.0444 (10)	0.1051 (18)	0.0024 (8)	0.0171 (12)	0.0143 (10)
O2'	0.1087 (17)	0.0605 (11)	0.0544 (13)	-0.0031 (12)	0.0069 (12)	-0.0051 (10)
C1'	0.0614 (17)	0.0451 (14)	0.102 (3)	-0.0143 (13)	-0.0104 (17)	-0.0143 (15)
C2'	0.077 (2)	0.0430 (15)	0.099 (3)	-0.0012 (13)	-0.0106 (18)	-0.0220 (15)
C3'	0.0627 (18)	0.0486 (15)	0.092 (2)	0.0101 (13)	-0.0075 (16)	-0.0179 (15)
C4'	0.0483 (14)	0.0480 (14)	0.083 (2)	0.0035 (11)	-0.0085 (14)	-0.0124 (14)
C5'	0.0477 (13)	0.0370 (11)	0.0559 (15)	-0.0005 (10)	-0.0089 (11)	-0.0013 (11)
C6'	0.0494 (13)	0.0375 (11)	0.0624 (16)	-0.0048 (10)	-0.0072 (12)	-0.0010 (11)
C7'	0.0459 (14)	0.0474 (13)	0.0645 (17)	-0.0124 (11)	-0.0019 (12)	0.0012 (12)
C8'	0.0424 (12)	0.0398 (12)	0.0528 (15)	-0.0050 (9)	-0.0026 (11)	0.0064 (10)
C9'	0.0453 (13)	0.0356 (11)	0.0489 (14)	-0.0034 (9)	-0.0055 (11)	0.0015 (9)

## supplementary materials

---

C10'	0.0417 (13)	0.0427 (12)	0.0629 (17)	-0.0068 (10)	-0.0077 (12)	-0.0056 (12)
C11'	0.0474 (13)	0.0449 (12)	0.0547 (15)	-0.0022 (10)	-0.0015 (11)	0.0052 (11)
C12'	0.0455 (13)	0.0548 (13)	0.0447 (14)	-0.0021 (10)	-0.0032 (11)	0.0053 (11)
C13'	0.0575 (16)	0.0607 (15)	0.0502 (16)	-0.0155 (12)	0.0022 (13)	0.0066 (13)
C14'	0.0674 (19)	0.080 (2)	0.067 (2)	-0.0244 (17)	0.0051 (16)	0.0108 (17)
C15'	0.0505 (16)	0.097 (2)	0.070 (2)	-0.0097 (16)	0.0056 (15)	0.0123 (18)
C16'	0.0467 (13)	0.0436 (12)	0.0580 (17)	0.0035 (10)	-0.0070 (12)	-0.0038 (12)
C17'	0.0399 (13)	0.0418 (11)	0.0717 (18)	-0.0020 (10)	0.0014 (12)	-0.0172 (12)
C18'	0.0589 (16)	0.0415 (13)	0.073 (2)	-0.0023 (11)	0.0052 (14)	-0.0020 (12)
C19'	0.102 (3)	0.0523 (18)	0.116 (3)	-0.0066 (17)	0.041 (2)	0.0019 (19)
C20'	0.080 (2)	0.0568 (18)	0.150 (4)	-0.0257 (17)	0.051 (2)	-0.027 (2)

### *Geometric parameters (Å, °)*

S1—C15	1.692 (3)	S1'—C15'	1.697 (3)
S1—C12	1.726 (2)	S1'—C12'	1.720 (3)
S2—C20	1.702 (4)	S2'—C20'	1.690 (4)
S2—C17	1.717 (2)	S2'—C17'	1.718 (2)
O1—C11	1.216 (3)	O1'—C11'	1.222 (3)
O2—C16	1.216 (3)	O2'—C16'	1.218 (3)
C1—C2	1.358 (4)	C1'—C2'	1.361 (5)
C1—C6	1.402 (4)	C1'—C6'	1.412 (4)
C1—H1	0.93	C1'—H1'	0.93
C2—C3	1.393 (4)	C2'—C3'	1.384 (4)
C2—H2	0.93	C2'—H2'	0.93
C3—C4	1.363 (4)	C3'—C4'	1.373 (4)
C3—H3	0.93	C3'—H3'	0.93
C4—C5	1.416 (3)	C4'—C5'	1.404 (4)
C4—H4	0.93	C4'—H4'	0.93
C5—C10	1.403 (3)	C5'—C10'	1.413 (3)
C5—C6	1.426 (3)	C5'—C6'	1.416 (3)
C6—C7	1.412 (3)	C6'—C7'	1.412 (4)
C7—C8	1.361 (4)	C7'—C8'	1.366 (3)
C7—H7	0.93	C7'—H7'	0.93
C8—C9	1.425 (3)	C8'—C9'	1.424 (3)
C8—C11	1.503 (4)	C8'—C11'	1.482 (4)
C9—C10	1.377 (3)	C9'—C10'	1.359 (4)
C9—C16	1.493 (4)	C9'—C16'	1.506 (3)
C10—H10	0.93	C10'—H10'	0.93
C11—C12	1.452 (3)	C11'—C12'	1.461 (4)
C12—C13	1.350 (3)	C12'—C13'	1.398 (4)
C13—C14	1.414 (4)	C13'—C14'	1.424 (4)
C13—H13	0.93	C13'—H13'	0.93
C14—C15	1.339 (4)	C14'—C15'	1.348 (5)
C14—H14	0.93	C14'—H14'	0.93
C15—H15	0.93	C15'—H15'	0.93
C16—C17	1.459 (4)	C16'—C17'	1.448 (4)
C17—C18	1.385 (4)	C17'—C18'	1.377 (4)
C18—C19	1.418 (4)	C18'—C19'	1.414 (4)



C18—H18	0.93	C18'—H18'	0.93
C19—C20	1.330 (5)	C19'—C20'	1.329 (6)
C19—H19	0.93	C19'—H19'	0.93
C20—H20	0.93	C20'—H20'	0.93
C15—S1—C12	91.10 (14)	C15'—S1'—C12'	91.82 (16)
C20—S2—C17	91.19 (15)	C20'—S2'—C17'	91.46 (17)
C2—C1—C6	121.1 (2)	C2'—C1'—C6'	121.0 (3)
C2—C1—H1	119.4	C2'—C1'—H1'	119.5
C6—C1—H1	119.4	C6'—C1'—H1'	119.5
C1—C2—C3	120.7 (3)	C1'—C2'—C3'	120.6 (3)
C1—C2—H2	119.6	C1'—C2'—H2'	119.7
C3—C2—H2	119.6	C3'—C2'—H2'	119.7
C4—C3—C2	120.4 (3)	C4'—C3'—C2'	120.4 (3)
C4—C3—H3	119.8	C4'—C3'—H3'	119.8
C2—C3—H3	119.8	C2'—C3'—H3'	119.8
C3—C4—C5	120.5 (2)	C3'—C4'—C5'	120.4 (3)
C3—C4—H4	119.7	C3'—C4'—H4'	119.8
C5—C4—H4	119.7	C5'—C4'—H4'	119.8
C10—C5—C4	122.4 (2)	C4'—C5'—C10'	122.6 (2)
C10—C5—C6	118.8 (2)	C4'—C5'—C6'	119.3 (2)
C4—C5—C6	118.7 (2)	C10'—C5'—C6'	118.1 (2)
C1—C6—C7	122.9 (2)	C1'—C6'—C7'	122.8 (2)
C1—C6—C5	118.5 (2)	C1'—C6'—C5'	118.2 (3)
C7—C6—C5	118.6 (2)	C7'—C6'—C5'	118.9 (2)
C8—C7—C6	121.8 (2)	C8'—C7'—C6'	122.3 (2)
C8—C7—H7	119.1	C8'—C7'—H7'	118.9
C6—C7—H7	119.1	C6'—C7'—H7'	118.9
C7—C8—C9	119.7 (2)	C7'—C8'—C9'	118.3 (2)
C7—C8—C11	121.4 (2)	C7'—C8'—C11'	122.5 (2)
C9—C8—C11	118.5 (2)	C9'—C8'—C11'	118.4 (2)
C10—C9—C8	119.4 (2)	C10'—C9'—C8'	120.6 (2)
C10—C9—C16	121.1 (2)	C10'—C9'—C16'	118.8 (2)
C8—C9—C16	119.1 (2)	C8'—C9'—C16'	120.2 (2)
C9—C10—C5	121.6 (2)	C9'—C10'—C5'	121.8 (2)
C9—C10—H10	119.2	C9'—C10'—H10'	119.1
C5—C10—H10	119.2	C5'—C10'—H10'	119.1
O1—C11—C12	122.4 (3)	O1'—C11'—C12'	119.2 (2)
O1—C11—C8	118.9 (2)	O1'—C11'—C8'	119.0 (2)
C12—C11—C8	118.7 (2)	C12'—C11'—C8'	121.8 (2)
C13—C12—C11	129.1 (2)	C13'—C12'—C11'	131.4 (3)
C13—C12—S1	111.06 (19)	C13'—C12'—S1'	111.6 (2)
C11—C12—S1	119.6 (2)	C11'—C12'—S1'	116.94 (19)
C12—C13—C14	112.7 (2)	C12'—C13'—C14'	110.2 (3)
C12—C13—H13	123.6	C12'—C13'—H13'	124.9
C14—C13—H13	123.6	C14'—C13'—H13'	124.9
C15—C14—C13	112.1 (3)	C15'—C14'—C13'	113.9 (3)
C15—C14—H14	123.9	C15'—C14'—H14'	123.1
C13—C14—H14	123.9	C13'—C14'—H14'	123.1
C14—C15—S1	113.0 (2)	C14'—C15'—S1'	112.5 (3)

## supplementary materials

---

C14—C15—H15	123.5	C14'—C15'—H15'	123.8
S1—C15—H15	123.5	S1'—C15'—H15'	123.8
O2—C16—C17	121.2 (2)	O2'—C16'—C17'	122.2 (2)
O2—C16—C9	120.0 (2)	O2'—C16'—C9'	120.0 (2)
C17—C16—C9	118.7 (2)	C17'—C16'—C9'	117.8 (2)
C18—C17—C16	129.5 (2)	C18'—C17'—C16'	128.5 (2)
C18—C17—S2	111.64 (19)	C18'—C17'—S2'	110.8 (2)
C16—C17—S2	118.83 (19)	C16'—C17'—S2'	120.4 (2)
C17—C18—C19	110.7 (3)	C17'—C18'—C19'	111.8 (3)
C17—C18—H18	124.6	C17'—C18'—H18'	124.1
C19—C18—H18	124.6	C19'—C18'—H18'	124.1
C20—C19—C18	113.7 (3)	C20'—C19'—C18'	112.8 (4)
C20—C19—H19	123.2	C20'—C19'—H19'	123.6
C18—C19—H19	123.2	C18'—C19'—H19'	123.6
C19—C20—S2	112.8 (2)	C19'—C20'—S2'	113.2 (3)
C19—C20—H20	123.6	C19'—C20'—H20'	123.4
S2—C20—H20	123.6	S2'—C20'—H20'	123.4
C6—C1—C2—C3	0.1 (4)	C6'—C1'—C2'—C3'	2.6 (6)
C1—C2—C3—C4	0.2 (4)	C1'—C2'—C3'—C4'	-1.5 (6)
C2—C3—C4—C5	0.3 (4)	C2'—C3'—C4'—C5'	0.1 (5)
C3—C4—C5—C10	-178.8 (2)	C3'—C4'—C5'—C10'	-178.0 (3)
C3—C4—C5—C6	-1.2 (3)	C3'—C4'—C5'—C6'	0.3 (4)
C2—C1—C6—C7	177.9 (3)	C2'—C1'—C6'—C7'	174.7 (3)
C2—C1—C6—C5	-1.0 (4)	C2'—C1'—C6'—C5'	-2.1 (5)
C10—C5—C6—C1	179.2 (2)	C4'—C5'—C6'—C1'	0.7 (4)
C4—C5—C6—C1	1.5 (3)	C10'—C5'—C6'—C1'	179.1 (3)
C10—C5—C6—C7	0.3 (3)	C4'—C5'—C6'—C7'	-176.3 (2)
C4—C5—C6—C7	-177.5 (2)	C10'—C5'—C6'—C7'	2.1 (4)
C1—C6—C7—C8	-177.3 (2)	C1'—C6'—C7'—C8'	-177.8 (3)
C5—C6—C7—C8	1.6 (3)	C5'—C6'—C7'—C8'	-1.0 (4)
C6—C7—C8—C9	-2.8 (3)	C6'—C7'—C8'—C9'	-1.3 (4)
C6—C7—C8—C11	169.7 (2)	C6'—C7'—C8'—C11'	168.4 (2)
C7—C8—C9—C10	2.0 (3)	C7'—C8'—C9'—C10'	2.4 (4)
C11—C8—C9—C10	-170.7 (2)	C11'—C8'—C9'—C10'	-167.7 (2)
C7—C8—C9—C16	-171.7 (2)	C7'—C8'—C9'—C16'	-170.7 (2)
C11—C8—C9—C16	15.6 (3)	C11'—C8'—C9'—C16'	19.2 (3)
C8—C9—C10—C5	-0.1 (3)	C8'—C9'—C10'—C5'	-1.2 (4)
C16—C9—C10—C5	173.5 (2)	C16'—C9'—C10'—C5'	171.9 (2)
C4—C5—C10—C9	176.6 (2)	C4'—C5'—C10'—C9'	177.3 (3)
C6—C5—C10—C9	-1.0 (3)	C6'—C5'—C10'—C9'	-1.0 (4)
C7—C8—C11—O1	-116.2 (3)	C7'—C8'—C11'—O1'	-139.6 (3)
C9—C8—C11—O1	56.4 (4)	C9'—C8'—C11'—O1'	30.0 (4)
C7—C8—C11—C12	63.0 (3)	C7'—C8'—C11'—C12'	39.7 (4)
C9—C8—C11—C12	-124.5 (3)	C9'—C8'—C11'—C12'	-150.6 (2)
O1—C11—C12—C13	-173.9 (3)	O1'—C11'—C12'—C13'	-172.2 (3)
C8—C11—C12—C13	7.0 (4)	C8'—C11'—C12'—C13'	8.4 (4)
O1—C11—C12—S1	1.0 (4)	O1'—C11'—C12'—S1'	5.0 (4)
C8—C11—C12—S1	-178.11 (18)	C8'—C11'—C12'—S1'	-174.30 (19)
C15—S1—C12—C13	0.2 (2)	C15'—S1'—C12'—C13'	-0.6 (2)

C15—S1—C12—C11	-175.6 (2)	C15'—S1'—C12'—C11'	-178.4 (2)
C11—C12—C13—C14	175.2 (3)	C11'—C12'—C13'—C14'	178.2 (3)
S1—C12—C13—C14	-0.1 (3)	S1'—C12'—C13'—C14'	0.9 (3)
C12—C13—C14—C15	-0.2 (4)	C12'—C13'—C14'—C15'	-0.8 (4)
C13—C14—C15—S1	0.3 (4)	C13'—C14'—C15'—S1'	0.3 (4)
C12—S1—C15—C14	-0.3 (3)	C12'—S1'—C15'—C14'	0.1 (3)
C10—C9—C16—O2	-131.8 (3)	C10'—C9'—C16'—O2'	-117.6 (3)
C8—C9—C16—O2	41.8 (4)	C8'—C9'—C16'—O2'	55.6 (3)
C10—C9—C16—C17	45.5 (3)	C10'—C9'—C16'—C17'	61.4 (3)
C8—C9—C16—C17	-140.8 (2)	C8'—C9'—C16'—C17'	-125.4 (2)
O2—C16—C17—C18	-167.8 (3)	O2'—C16'—C17'—C18'	-164.4 (3)
C9—C16—C17—C18	14.9 (4)	C9'—C16'—C17'—C18'	16.6 (4)
O2—C16—C17—S2	11.8 (4)	O2'—C16'—C17'—S2'	8.2 (4)
C9—C16—C17—S2	-165.52 (19)	C9'—C16'—C17'—S2'	-170.75 (18)
C20—S2—C17—C18	-0.6 (2)	C20'—S2'—C17'—C18'	1.2 (2)
C20—S2—C17—C16	179.7 (2)	C20'—S2'—C17'—C16'	-172.6 (2)
C16—C17—C18—C19	-180.0 (3)	C16'—C17'—C18'—C19'	171.4 (3)
S2—C17—C18—C19	0.4 (3)	S2'—C17'—C18'—C19'	-1.8 (3)
C17—C18—C19—C20	0.1 (4)	C17'—C18'—C19'—C20'	1.6 (4)
C18—C19—C20—S2	-0.6 (4)	C18'—C19'—C20'—S2'	-0.7 (5)
C17—S2—C20—C19	0.7 (3)	C17'—S2'—C20'—C19'	-0.3 (3)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C4—H4 $\cdots$ O2 <sup>i</sup>	0.93	2.53	3.407 (3)	158
C13—H13 $\cdots$ Cg1 <sup>ii</sup>	0.93	2.86	3.737 (2)	157

Symmetry codes: (i)  $-x+1, y+1/2, -z+1$ ; (ii)  $-x+1, y-1/2, -z+1$ .

Fig. 1




Fig. 2

