

(E)-Ethyl 3-(4-fluoroanilino)-2-(4-methoxyphenyl)acrylate

Da-Gui Zheng^{a*} and Zhu-Ping Xiao^b

^aKey Laboratory of Applied Organic Chemistry, Higher Institutions of Jiangxi Province, Shangrao Normal College, Shangrao 334001, Jiangxi, People's Republic of China, and ^bCollege of Chemistry & Chemical Engineering, Jishou University, Jishou 416000, People's Republic of China
Correspondence e-mail: shenyangzhou@163.com


Received 21 October 2008; accepted 21 November 2008

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.050; wR factor = 0.162; data-to-parameter ratio = 13.7.

In the title compound, $\text{C}_{18}\text{H}_{18}\text{FNO}_3$, the dihedral angles between the two benzene rings and the plane through the acrylate group and the fluorophenyl ring are $61.58(8)$ and $13.33(9)^\circ$, respectively. Molecules are linked into ribbons through $\text{C}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, and further linked by $\text{C}-\text{H}\cdots\pi$ interactions, forming a three-dimensional network.

Related literature

For related literature regarding the antimicrobial activity of 3-arylaminoo-2-aryl acrylates, see: Shi *et al.* (2007); Xiao *et al.* (2007, 2008); Xue *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{18}\text{FNO}_3$	$V = 1630.0(6)\text{ \AA}^3$
$M_r = 315.33$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 19.000(4)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 6.0400(12)\text{ \AA}$	$T = 293(2)\text{ K}$
$c = 15.081(3)\text{ \AA}$	$0.30 \times 0.30 \times 0.20\text{ mm}$
$\beta = 109.64(3)^\circ$	

Data collection

Enraf–Nonius CAD-4 diffractometer	3073 measured reflections
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	2943 independent reflections
$T_{\min} = 0.972$, $T_{\max} = 0.981$	1807 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.162$	$\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$
$S = 1.02$	$\Delta\rho_{\min} = -0.16\text{ e \AA}^{-3}$
2943 reflections	
215 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C12—H12 \cdots O1 ⁱ	0.93	2.49	3.401 (3)	167
N1—H1 \cdots O3 ⁱⁱ	0.83 (2)	2.56 (3)	3.229 (3)	138 (2)
C16—H16B \cdots Cg1 ⁱⁱⁱ	0.97	2.99	3.788 (3)	141
C18—H18A \cdots Cg2 ^{iv}	0.96	2.80	3.626 (3)	145

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y+1, -z$; (iii) $x, -y+\frac{1}{2}, z+\frac{1}{2}$; (iv) $-x+1, -y, -z$. Cg1 and Cg2 are the centroids of the C1–C6 and C7–C12 rings, respectively.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This research was financially supported by the Education Department of Jiangxi Province, P. R. China (grant No. 2007-402).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2149).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shi, D.-H., You, Z.-L., Xu, C., Zhang, Q. & Zhu, H.-L. (2007). *Inorg. Chem. Commun.* **10**, 404–406.
- Xiao, Z.-P., Fang, R.-Q., Li, H.-Q., Shi, L., Xue, J.-Y., Zheng, Y. & Zhu, H.-L. (2008). *Eur. J. Med. Chem.* **43**, 1828–1836.
- Xiao, Z.-P., Xue, J.-Y., Tan, S.-H., Li, H.-Q. & Zhu, H.-L. (2007). *Bioorg. Med. Chem.* **15**, 4212–4219.
- Xue, J.-Y., Xiao, Z.-P., Shi, L., Tan, S.-H., Li, H.-Q. & Zhu, H.-L. (2007). *Aust. J. Chem.* **60**, 957–962.

supplementary materials

Acta Cryst. (2008). E64, o2456 [doi:10.1107/S1600536808039184]

(E)-Ethyl 3-(4-fluoroanilino)-2-(4-methoxyphenyl)acrylate

D.-G. Zheng and Z.-P. Xiao

Comment

3-Arylamino-2-aryl acrylates, enamines structurally like Schiff bases, show high antimicrobial activity (Xiao *et al.*, 2007; Xue *et al.*, 2007; Xiao *et al.*, 2008; Shi *et al.* 2007), especially for bacteria. In a continuation of our work on the structural characterization of enamine derivatives, we report here the crystal structure of the title compound, (I) (Fig. 1).

The N1—H group lies approximately in the same planes as the fluorophenyl and acrylate groups (with dihedral angles of 4.7 (2) ° and 8.9 (2) °, respectively), suggesting that one of the p orbitals of N1 is conjugated with the π molecular orbitals of the two moieties, thus shortening both the C1—N1 (1.408 (3) Å) and C13—N1 (1.359 (3) Å) bonds. All other double and single bond lengths fall within normal values (Allen *et al.*, 1987).

Molecules are linked into ribbons running along the b-axis via C—H \cdots O and N—H \cdots O hydrogen bonds (Fig. 2 and Table 1). These ribbons are interconnected via weak C16-H16B \cdots π (centroid of C1 to C6) and C18-H18A \cdots π (centroid of C7 to C12) interactions (Table 1 and Fig. 3).

Experimental

Equimolar quantities (6 mmol) of ethyl 2-(4-methoxyphenyl)-3-oxopropanoate (1.33 g) and 4-fluorobenzenamine (0.67 g) in absolute alcohol (18 ml) were heated at 344–354 K for 2 h. The excess solvent was removed under reduced pressure. The residue was purified by flash chromatography with EtOAc-petroleum ether (1:10) to afford two fractions. The first fraction gave the Z-isomer, and the second fraction, after partial solvent evaporation, furnished colorless blocks of (I) suitable for single-crystal structure determination.

Refinement

The H atom bonded to N1 was located in a difference Fourier map. All other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H = 0.93, 0.96 and 0.97 Å for the aromatic, CH₃ and CH₂ type H atoms, respectively. $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{parent atoms})$ were assigned for amino, aromatic and CH₂ type H-atoms and 1.5 $U_{\text{eq}}(\text{parent atoms})$ for CH₃ type H-atoms.

Figures




Fig. 1. The molecular structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. A partial packing diagram of the title compound showing the ribbons connected by C—H···O and N—H···O hydrogen bonds (indicated by dashed lines), viewed along the b axis.




Fig. 3. The crystal packing of the title compound, showing the linking of the hydrogen bonded ribbons by C—H···π interactions. Dashed lines indicate C—H···O, N—H···O and C—H···π interactions.

(E)-Ethyl 3-(4-fluoroanilino)-2-(4-methoxyphenyl)acrylate

Crystal data

C ₁₈ H ₁₈ FNO ₃	$F_{000} = 664$
$M_r = 315.33$	$D_x = 1.285 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 19.000 (4) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 6.0400 (12) \text{ \AA}$	Cell parameters from 1632 reflections
$c = 15.081 (3) \text{ \AA}$	$\theta = 1.4\text{--}24.7^\circ$
$\beta = 109.64 (3)^\circ$	$\mu = 0.10 \text{ mm}^{-1}$
$V = 1630.0 (6) \text{ \AA}^3$	$T = 293 (2) \text{ K}$
$Z = 4$	Block, colorless
	$0.30 \times 0.30 \times 0.20 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	2943 independent reflections
Radiation source: fine-focus sealed tube	1807 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 293(2) \text{ K}$	$\theta_{\text{max}} = 25.3^\circ$
$\omega/2\theta$ scans	$\theta_{\text{min}} = 1.1^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = -22 \rightarrow 21$
$T_{\text{min}} = 0.972$, $T_{\text{max}} = 0.981$	$k = -7 \rightarrow 0$
3073 measured reflections	$l = 0 \rightarrow 18$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.050$	$w = 1/[\sigma^2(F_o^2) + (0.0863P)^2 + 0.0162P]$

$wR(F^2) = 0.162$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\max} < 0.001$
2943 reflections	$\Delta\rho_{\max} = 0.15 \text{ e \AA}^{-3}$
215 parameters	$\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.044 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.78366 (12)	0.7249 (4)	0.02705 (17)	0.0468 (6)
C2	0.86057 (14)	0.7171 (5)	0.0575 (2)	0.0637 (8)
H2	0.8858	0.6095	0.1005	0.076*
C3	0.90034 (15)	0.8673 (6)	0.0246 (2)	0.0733 (9)
H3	0.9523	0.8617	0.0451	0.088*
C4	0.86292 (16)	1.0237 (5)	-0.0380 (2)	0.0682 (9)
C5	0.78687 (17)	1.0339 (5)	-0.0708 (2)	0.0721 (9)
H5	0.7620	1.1403	-0.1148	0.086*
C6	0.74779 (15)	0.8829 (5)	-0.0373 (2)	0.0640 (8)
H6	0.6958	0.8882	-0.0588	0.077*
C7	0.64691 (12)	0.3267 (4)	0.14607 (16)	0.0431 (6)
C8	0.59236 (13)	0.1688 (4)	0.10497 (18)	0.0494 (7)
H8	0.6063	0.0338	0.0862	0.059*
C9	0.51803 (13)	0.2090 (5)	0.09161 (18)	0.0513 (7)
H9	0.4825	0.1013	0.0639	0.062*
C10	0.49607 (13)	0.4075 (4)	0.11902 (17)	0.0470 (6)
C11	0.54911 (13)	0.5661 (4)	0.16114 (19)	0.0523 (7)
H11	0.5350	0.7002	0.1804	0.063*
C12	0.62356 (13)	0.5233 (4)	0.17434 (19)	0.0510 (7)
H12	0.6591	0.6302	0.2032	0.061*
C13	0.76733 (13)	0.4059 (4)	0.12087 (17)	0.0465 (6)
H13	0.8176	0.3699	0.1356	0.056*
C14	0.72695 (12)	0.2824 (4)	0.16064 (16)	0.0433 (6)

supplementary materials

C15	0.76226 (13)	0.0959 (4)	0.22121 (18)	0.0458 (6)
C16	0.87264 (15)	-0.1081 (5)	0.2948 (2)	0.0669 (8)
H16A	0.8550	-0.2514	0.2671	0.080*
H16B	0.8630	-0.0960	0.3538	0.080*
C17	0.95460 (15)	-0.0844 (6)	0.3113 (3)	0.0887 (11)
H17A	0.9637	-0.1045	0.2530	0.133*
H17B	0.9817	-0.1941	0.3558	0.133*
H17C	0.9709	0.0606	0.3358	0.133*
C18	0.39590 (15)	0.6241 (5)	0.1368 (2)	0.0751 (9)
H18A	0.4220	0.6374	0.2032	0.113*
H18B	0.3432	0.6130	0.1255	0.113*
H18C	0.4057	0.7521	0.1051	0.113*
F1	0.90279 (11)	1.1721 (4)	-0.07006 (17)	0.1125 (8)
H1	0.6954 (14)	0.606 (4)	0.0454 (19)	0.059 (8)*
N1	0.74101 (12)	0.5785 (4)	0.06106 (16)	0.0567 (6)
O1	0.73060 (10)	-0.0227 (3)	0.26022 (14)	0.0667 (6)
O2	0.83483 (9)	0.0669 (3)	0.23152 (13)	0.0584 (5)
O3	0.42082 (9)	0.4306 (3)	0.10206 (13)	0.0618 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0401 (13)	0.0536 (15)	0.0482 (14)	-0.0053 (12)	0.0170 (11)	0.0019 (13)
C2	0.0465 (15)	0.076 (2)	0.0642 (17)	-0.0017 (15)	0.0135 (13)	0.0221 (16)
C3	0.0434 (15)	0.093 (2)	0.082 (2)	-0.0056 (16)	0.0193 (15)	0.026 (2)
C4	0.0596 (18)	0.073 (2)	0.082 (2)	-0.0085 (16)	0.0371 (16)	0.0172 (18)
C5	0.074 (2)	0.065 (2)	0.085 (2)	0.0109 (16)	0.0370 (17)	0.0304 (18)
C6	0.0450 (15)	0.0681 (19)	0.078 (2)	0.0063 (14)	0.0193 (14)	0.0157 (17)
C7	0.0403 (13)	0.0433 (14)	0.0443 (14)	-0.0067 (11)	0.0125 (10)	0.0005 (12)
C8	0.0456 (14)	0.0464 (15)	0.0548 (15)	-0.0018 (12)	0.0150 (12)	-0.0075 (13)
C9	0.0412 (13)	0.0540 (17)	0.0528 (15)	-0.0131 (12)	0.0081 (11)	-0.0127 (13)
C10	0.0361 (12)	0.0550 (16)	0.0482 (14)	-0.0008 (12)	0.0118 (11)	0.0038 (13)
C11	0.0490 (15)	0.0417 (14)	0.0668 (17)	-0.0045 (12)	0.0202 (13)	-0.0085 (13)
C12	0.0449 (14)	0.0364 (14)	0.0694 (18)	-0.0074 (11)	0.0164 (13)	-0.0033 (13)
C13	0.0364 (12)	0.0493 (15)	0.0502 (15)	-0.0021 (12)	0.0097 (11)	0.0004 (13)
C14	0.0410 (13)	0.0406 (14)	0.0471 (14)	-0.0060 (11)	0.0133 (11)	-0.0046 (12)
C15	0.0441 (13)	0.0422 (14)	0.0521 (15)	-0.0065 (12)	0.0174 (12)	-0.0088 (13)
C16	0.0616 (17)	0.0523 (17)	0.076 (2)	0.0087 (15)	0.0087 (15)	0.0069 (16)
C17	0.0565 (18)	0.093 (3)	0.104 (3)	0.0198 (18)	0.0101 (18)	0.004 (2)
C18	0.0521 (16)	0.091 (2)	0.085 (2)	0.0079 (16)	0.0267 (16)	-0.0201 (19)
F1	0.0896 (13)	0.1078 (17)	0.154 (2)	-0.0112 (12)	0.0592 (14)	0.0556 (15)
N1	0.0359 (12)	0.0658 (15)	0.0665 (15)	0.0010 (12)	0.0146 (11)	0.0177 (13)
O1	0.0589 (11)	0.0540 (12)	0.0889 (15)	-0.0014 (10)	0.0272 (11)	0.0181 (11)
O2	0.0439 (10)	0.0583 (12)	0.0718 (13)	0.0047 (9)	0.0177 (9)	0.0084 (10)
O3	0.0399 (10)	0.0733 (13)	0.0714 (13)	0.0012 (9)	0.0177 (9)	-0.0061 (11)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.368 (4)	C11—C12	1.385 (3)
-------	-----------	---------	-----------

C1—C2	1.377 (3)	C11—H11	0.9300
C1—N1	1.408 (3)	C12—H12	0.9300
C2—C3	1.375 (4)	C13—C14	1.347 (3)
C2—H2	0.9300	C13—N1	1.359 (3)
C3—C4	1.355 (4)	C13—H13	0.9300
C3—H3	0.9300	C14—C15	1.463 (3)
C4—C5	1.362 (4)	C15—O1	1.208 (3)
C4—F1	1.363 (3)	C15—O2	1.346 (3)
C5—C6	1.375 (4)	C16—O2	1.444 (3)
C5—H5	0.9300	C16—C17	1.499 (4)
C6—H6	0.9300	C16—H16A	0.9700
C7—C12	1.384 (3)	C16—H16B	0.9700
C7—C8	1.392 (3)	C17—H17A	0.9600
C7—C14	1.486 (3)	C17—H17B	0.9600
C8—C9	1.379 (3)	C17—H17C	0.9600
C8—H8	0.9300	C18—O3	1.425 (3)
C9—C10	1.378 (4)	C18—H18A	0.9600
C9—H9	0.9300	C18—H18B	0.9600
C10—O3	1.372 (3)	C18—H18C	0.9600
C10—C11	1.381 (3)	N1—H1	0.83 (2)
C6—C1—C2	118.7 (2)	C7—C12—H12	118.9
C6—C1—N1	119.1 (2)	C11—C12—H12	118.9
C2—C1—N1	122.2 (2)	C14—C13—N1	125.6 (2)
C3—C2—C1	120.5 (3)	C14—C13—H13	117.2
C3—C2—H2	119.8	N1—C13—H13	117.2
C1—C2—H2	119.8	C13—C14—C15	119.6 (2)
C4—C3—C2	119.2 (2)	C13—C14—C7	122.8 (2)
C4—C3—H3	120.4	C15—C14—C7	117.6 (2)
C2—C3—H3	120.4	O1—C15—O2	121.7 (2)
C3—C4—C5	121.9 (3)	O1—C15—C14	124.2 (2)
C3—C4—F1	118.8 (3)	O2—C15—C14	114.2 (2)
C5—C4—F1	119.3 (3)	O2—C16—C17	107.4 (2)
C4—C5—C6	118.3 (3)	O2—C16—H16A	110.2
C4—C5—H5	120.8	C17—C16—H16A	110.2
C6—C5—H5	120.8	O2—C16—H16B	110.2
C1—C6—C5	121.4 (2)	C17—C16—H16B	110.2
C1—C6—H6	119.3	H16A—C16—H16B	108.5
C5—C6—H6	119.3	C16—C17—H17A	109.5
C12—C7—C8	117.2 (2)	C16—C17—H17B	109.5
C12—C7—C14	121.8 (2)	H17A—C17—H17B	109.5
C8—C7—C14	120.9 (2)	C16—C17—H17C	109.5
C9—C8—C7	121.1 (2)	H17A—C17—H17C	109.5
C9—C8—H8	119.4	H17B—C17—H17C	109.5
C7—C8—H8	119.4	O3—C18—H18A	109.5
C10—C9—C8	120.5 (2)	O3—C18—H18B	109.5
C10—C9—H9	119.7	H18A—C18—H18B	109.5
C8—C9—H9	119.7	O3—C18—H18C	109.5
O3—C10—C9	115.8 (2)	H18A—C18—H18C	109.5
O3—C10—C11	124.6 (2)	H18B—C18—H18C	109.5

supplementary materials


C9—C10—C11	119.6 (2)	C13—N1—C1	126.5 (2)
C10—C11—C12	119.2 (2)	C13—N1—H1	117.1 (19)
C10—C11—H11	120.4	C1—N1—H1	116.3 (19)
C12—C11—H11	120.4	C15—O2—C16	115.8 (2)
C7—C12—C11	122.3 (2)	C10—O3—C18	118.0 (2)
C6—C1—C2—C3	0.9 (4)	C10—C11—C12—C7	0.5 (4)
N1—C1—C2—C3	-178.1 (3)	N1—C13—C14—C15	177.8 (2)
C1—C2—C3—C4	0.1 (5)	N1—C13—C14—C7	-2.3 (4)
C2—C3—C4—C5	-1.3 (5)	C12—C7—C14—C13	-62.1 (3)
C2—C3—C4—F1	179.9 (3)	C8—C7—C14—C13	119.3 (3)
C3—C4—C5—C6	1.4 (5)	C12—C7—C14—C15	117.8 (3)
F1—C4—C5—C6	-179.8 (3)	C8—C7—C14—C15	-60.8 (3)
C2—C1—C6—C5	-0.8 (4)	C13—C14—C15—O1	179.5 (2)
N1—C1—C6—C5	178.2 (3)	C7—C14—C15—O1	-0.4 (4)
C4—C5—C6—C1	-0.3 (5)	C13—C14—C15—O2	0.3 (3)
C12—C7—C8—C9	1.2 (4)	C7—C14—C15—O2	-179.6 (2)
C14—C7—C8—C9	179.8 (2)	C14—C13—N1—C1	172.4 (2)
C7—C8—C9—C10	-0.1 (4)	C6—C1—N1—C13	176.2 (3)
C8—C9—C10—O3	179.7 (2)	C2—C1—N1—C13	-4.8 (4)
C8—C9—C10—C11	-0.7 (4)	O1—C15—O2—C16	-2.4 (3)
O3—C10—C11—C12	-179.9 (2)	C14—C15—O2—C16	176.8 (2)
C9—C10—C11—C12	0.5 (4)	C17—C16—O2—C15	-169.0 (2)
C8—C7—C12—C11	-1.3 (4)	C9—C10—O3—C18	174.3 (2)
C14—C7—C12—C11	-180.0 (2)	C11—C10—O3—C18	-5.2 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C12—H12 \cdots O1 ⁱ	0.93	2.49	3.401 (3)	167
N1—H1 \cdots O3 ⁱⁱ	0.83 (2)	2.56 (3)	3.229 (3)	138 (2)
C16—H16B \cdots Cg1 ⁱⁱⁱ	0.97	2.99	3.788 (3)	141
C18—H18A \cdots Cg2 ^{iv}	0.96	2.80	3.626 (3)	145

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y+1, -z$; (iii) $x, -y+1/2, z+1/2$; (iv) $-x+1, -y, -z$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

