

Dichlorido(di-2-pyridylamine)-mercury(II)

Mohammad Yousefi,^a Mohammad Reza Allahgholi Ghasri,^a Amene Heidari^b and Vahid Amani^{a*}

^aIslamic Azad University, Shahr-e-Rey Branch, Tehran, Iran, and ^bDepartment of Chemistry, University of Zabol, Iran
Correspondence e-mail: v_amani2002@yahoo.com


Received 28 November 2008; accepted 30 November 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.012\text{ \AA}$; R factor = 0.046; wR factor = 0.126; data-to-parameter ratio = 21.4.

In the molecule of the title compound, $[\text{HgCl}_2(\text{C}_{10}\text{H}_9\text{N}_3)]$, the Hg^{II} atom is four-coordinated in a distorted tetrahedral configuration by two N atoms from the chelating di-2-pyridylamine ligand and by two Cl atoms. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds link the molecules into centrosymmetric dimers. There is a $\pi-\pi$ contact between the pyridine rings [centroid–centroid distance = 3.896 (5) \AA].

Related literature

For related literature, see: Ahmadi *et al.* (2008); Kalateh, Ebadi *et al.* (2008); Kalateh, Norouzi *et al.* (2008); Khavasi *et al.* (2008); Tadayon Pour *et al.* (2008); Yousefi, Rashidi Vahid *et al.* (2008); Yousefi, Tadayon Pour *et al.* (2008). For related structures, see: Chen *et al.* (2006); Liu *et al.* (2004). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$[\text{HgCl}_2(\text{C}_{10}\text{H}_9\text{N}_3)]$
 $M_r = 442.69$

Triclinic, $P\bar{1}$
 $a = 8.0268 (12)\text{ \AA}$

$b = 8.6127 (11)\text{ \AA}$
 $c = 9.6118 (14)\text{ \AA}$
 $\alpha = 110.606 (11)^\circ$
 $\beta = 98.958 (12)^\circ$
 $\gamma = 96.862 (11)^\circ$
 $V = 603.38 (15)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 13.17\text{ mm}^{-1}$
 $T = 298 (2)\text{ K}$
 $0.24 \times 0.21 \times 0.15\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi scan (*SADABS*; Sheldrick, 1998)
 $T_{\min} = 0.061$, $T_{\max} = 0.142$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.126$
 $S = 1.05$
3214 reflections
150 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 2.43\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -2.08\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Cl1–Hg1	2.3875 (19)	N1–Hg1	2.369 (6)
Cl2–Hg1	2.4579 (19)	N3–Hg1	2.290 (6)
N1–Hg1–Cl1	112.30 (15)	N3–Hg1–Cl2	123.92 (14)
N1–Hg1–Cl2	94.92 (15)	N3–Hg1–N1	82.4 (2)
N3–Hg1–Cl1	112.21 (15)	Cl1–Hg1–Cl2	120.14 (7)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2–H2B \cdots Cl2 ⁱ	0.95 (16)	2.41 (16)	3.345 (6)	169 (13)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to the Islamic Azad University, Shahr-e-Rey Branch, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2588).

References

- Ahmadi, R., Ebadi, A., Kalateh, K., Norouzi, A. & Amani, V. (2008). *Acta Cryst. E64*, m1407.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, W. T., Wang, M. S., Liu, X., Guo, G. C. & Huang, J. S. (2006). *Cryst. Growth Des.* **6**, 2289–2300.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.

metal-organic compounds

- Kalateh, K., Ebadi, A., Ahmadi, R., Amani, V. & Khavasi, H. R. (2008). *Acta Cryst. E*64, m1397–m1398.
- Kalateh, K., Norouzi, A., Ebadi, A., Ahmadi, R. & Amani, V. (2008). *Acta Cryst. E*64, m1583–m1584.
- Khavasi, H. R., Abedi, A., Amani, V., Notash, B. & Safari, N. (2008). *Polyhedron*, **27**, 1848–1854.
- Liu, Q. D., Wang, R. & Wang, S. (2004). *Dalton Trans.* pp. 2073–2079.
- Sheldrick, G. M. (1998). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Tadayon Pour, N., Ebadi, A., Abedi, A., Amani, V. & Khavasi, H. R. (2008). *Acta Cryst. E*64, m1305.
- Yousefi, M., Rashidi Vahid, R., Amani, V., Arab Chamjangali, M. & Khavasi, H. R. (2008). *Acta Cryst. E*64, m1339–m1340.
- Yousefi, M., Tadayon Pour, N., Amani, V. & Khavasi, H. R. (2008). *Acta Cryst. E*64, m1259.

supplementary materials

Acta Cryst. (2009). E65, m9-m10 [doi:10.1107/S1600536808040294]

Dichlorido(di-2-pyridylamine)mercury(II)

M. Yousefi, M. R. Allahgholi Ghasri, A. Heidari and V. Amani

Comment

Recently, we reported the syntheses and crystal structures of $[\text{Hg}(\text{NH(py})_2\text{Br}_2]$, (II), (Kalateh, Norouzi *et al.*, 2008), $[\text{Hg}(4,4'\text{-dmbpy})\text{I}_2]$, (III), (Yousefi, Tadayon Pour *et al.*, 2008), $[\text{Hg}(5,5'\text{-dmbpy})\text{I}_2]$, (IV), (Tadayon Pour *et al.*, 2008), $[\text{Hg}(\text{dmphen})\text{I}_2]$, (V), (Yousefi, Rashidi Vahid *et al.*, 2008), $\{\text{[HgCl(dm4bt)}\]_2(\mu\text{-Cl})_2\}$, (VI), (Khavasi *et al.*, 2008), $[\text{Hg}(6\text{-mbpy})\text{Cl}_2]$, (VII), (Ahmadi *et al.*, 2008) and $[\{\text{HgBr}(4,4'\text{-dmbpy})\}_2(\mu\text{-Br})_2]$, (VIII), (Kalateh, Ebadi *et al.*, 2008) [where $\text{NH(py})_2$ is di-2-pyridylamine, 4,4'-dmbpy is 4,4'-dimethyl-2,2'-bipyridine, 5,5'-dmbpy is 5,5'-dimethyl-2,2'-bipyridine, 6-mbpy is 6-methyl-2,2'-bipyridine, dmphen is 4,7-diphenyl-1,10-phenanthroline and dm4bt is 2,2'-dimethyl-4,4'-bithiazole].

There are several Hg^{II} complexes, with formula, $[\text{Hg}(\text{N—N})\text{Cl}_2]$, such as $[\text{Hg}(\text{bipy})\text{Cl}_2]$, (IX), and $[\text{Hg}(\text{bipy})\text{Cl}_2][\text{HgCl}_2]$, (X), (Chen *et al.*, 2006) and $[\text{Hg}(\text{dpdmbip})\text{Cl}_2]\text{CH}_2\text{Cl}_2$, (XI), (Liu *et al.*, 2004) [where bipy is 2,2'-bipyridine and dpdmbip is 4,4'-diphenyl-6,6'-dimethyl-2,2'-bipyrimidine] have been synthesized and characterized by single-crystal X-ray diffraction methods. We report herein the synthesis and crystal structure of the title compound, (I).

In the title compound, (Fig. 1), the Hg^{II} atom is four-coordinated in a distorted tetrahedral configuration by two N atoms from di-2-pyridylamine and two Cl atoms. The Hg-Cl and Hg-N bond lengths (Allen *et al.*, 1987) and angles (Table 1) are within normal ranges.

In the crystal structure, intermolecular N-H \cdots Cl hydrogen bonds (Table 2) link the molecules into centrosymmetric dimers, in which they may be effective in the stabilization of the crystal structure (Fig. 2). The $\pi\text{-}\pi$ contact between the pyridine rings, Cg2 \cdots Cg3ⁱ [symmetry code: (i) -x, 1 - y, 1 - z, where Cg2 and Cg3 are centroids of the rings A (N1/C1-C5) and B (N3/C6-C10), respectively] may further stabilize the structure, with centroid-centroid distance of 3.896 (5) \AA .

Experimental

For the preparation of the title compound, (I), a solution of di-2-pyridylamine (0.25 g, 1.43 mmol) in methanol (20 ml) was added to a solution of HgCl_2 (0.39 g, 1.43 mmol) in acetonitrile (20 ml) and the resulting colorless solution was stirred for 20 min at 313 K. This solution was left to evaporate slowly at room temperature. After one week, colorless block crystals of the title compound were isolated (yield; 0.47 g, 74.3%).

Refinement

H_2B atom (for NH) was located in difference synthesis and refined isotropically [$\text{N-H} = 0.95$ (14) \AA and $\text{U}_{\text{iso}}(\text{H}) = 0.10$ (4) \AA^2]. The remaining H atoms were positioned geometrically, with $\text{C-H} = 0.93$ \AA for aromatic H and constrained to ride on their parent atoms, with $\text{U}_{\text{iso}}(\text{H}) = 1.2\text{U}_{\text{eq}}(\text{C})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 40% probability level.




Fig. 2. A packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

Dichlorido(di-2-pyridylamine)mercury(II)

Crystal data

[HgCl ₂ (C ₁₀ H ₉ N ₃)]	Z = 2
M _r = 442.69	F ₀₀₀ = 408
Triclinic, P $\bar{1}$	D _x = 2.437 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation
a = 8.0268 (12) Å	λ = 0.71073 Å
b = 8.6127 (11) Å	Cell parameters from 1652 reflections
c = 9.6118 (14) Å	θ = 2.6–29.2°
α = 110.606 (11)°	μ = 13.17 mm ⁻¹
β = 98.958 (12)°	T = 298 (2) K
γ = 96.862 (11)°	Block, colorless
V = 603.38 (15) Å ³	0.24 × 0.21 × 0.15 mm

Data collection

Bruker SMART CCD area-detector diffractometer	3214 independent reflections
Radiation source: fine-focus sealed tube	2806 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.069$
T = 298(2) K	$\theta_{\text{max}} = 29.2^\circ$
ϕ and ω scans	$\theta_{\text{min}} = 2.6^\circ$
Absorption correction: multi scan (SADABS; Sheldrick, 1998)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.061$, $T_{\text{max}} = 0.142$	$k = -11 \rightarrow 11$
7105 measured reflections	$l = -13 \rightarrow 12$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.046$	$w = 1/[\sigma^2(F_o^2) + (0.075P)^2 + 0.8992P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.126$	$(\Delta/\sigma)_{\max} = 0.026$
$S = 1.05$	$\Delta\rho_{\max} = 2.43 \text{ e \AA}^{-3}$
3214 reflections	$\Delta\rho_{\min} = -2.08 \text{ e \AA}^{-3}$
150 parameters	Extinction correction: SHELXTL (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.048 (3)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.21151 (4)	0.24646 (4)	0.63832 (4)	0.06055 (18)
Cl1	0.4456 (3)	0.1830 (3)	0.7818 (3)	0.0683 (5)
Cl2	-0.0856 (2)	0.1450 (2)	0.6448 (2)	0.0553 (4)
N1	0.1944 (8)	0.5364 (8)	0.7382 (7)	0.0497 (12)
N2	0.2047 (8)	0.5833 (7)	0.5110 (6)	0.0459 (11)
N3	0.2876 (7)	0.3143 (7)	0.4439 (7)	0.0447 (11)
C1	0.1758 (12)	0.5982 (12)	0.8836 (9)	0.0636 (19)
H1	0.1862	0.5294	0.9393	0.076*
C2	0.1428 (12)	0.7553 (13)	0.9537 (9)	0.069 (2)
H2	0.1285	0.7920	1.0537	0.083*
H2B	0.186 (18)	0.662 (18)	0.464 (16)	0.10 (4)*
C3	0.1312 (11)	0.8591 (11)	0.8713 (9)	0.0624 (18)
H3	0.1115	0.9683	0.9157	0.075*
C4	0.1493 (9)	0.7975 (9)	0.7231 (8)	0.0514 (14)
H4	0.1379	0.8635	0.6649	0.062*

supplementary materials

C5	0.1848 (7)	0.6352 (7)	0.6597 (7)	0.0398 (11)
C6	0.2674 (7)	0.4531 (7)	0.4151 (7)	0.0395 (10)
C7	0.3045 (9)	0.4711 (10)	0.2836 (7)	0.0499 (13)
H7	0.2849	0.5656	0.2627	0.060*
C8	0.3715 (12)	0.3449 (13)	0.1842 (10)	0.067 (2)
H8	0.4006	0.3559	0.0977	0.080*
C9	0.3935 (9)	0.2066 (10)	0.2156 (9)	0.0587 (18)
H9	0.4365	0.1208	0.1498	0.070*
C10	0.3524 (9)	0.1931 (9)	0.3444 (10)	0.0556 (16)
H10	0.3694	0.0976	0.3648	0.067*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.0618 (2)	0.0660 (2)	0.0781 (3)	0.02555 (14)	0.02487 (14)	0.04774 (18)
Cl1	0.0653 (11)	0.0710 (11)	0.0816 (13)	0.0210 (9)	0.0091 (9)	0.0445 (10)
Cl2	0.0623 (9)	0.0551 (8)	0.0628 (9)	0.0170 (7)	0.0277 (7)	0.0314 (7)
N1	0.051 (3)	0.058 (3)	0.050 (3)	0.019 (2)	0.014 (2)	0.027 (2)
N2	0.059 (3)	0.043 (2)	0.045 (3)	0.016 (2)	0.015 (2)	0.024 (2)
N3	0.043 (2)	0.041 (2)	0.056 (3)	0.006 (2)	0.013 (2)	0.023 (2)
C1	0.072 (5)	0.078 (5)	0.051 (4)	0.020 (4)	0.017 (3)	0.033 (4)
C2	0.071 (5)	0.084 (6)	0.047 (4)	0.019 (4)	0.012 (3)	0.017 (4)
C3	0.066 (4)	0.061 (4)	0.052 (4)	0.013 (4)	0.015 (3)	0.009 (3)
C4	0.053 (3)	0.045 (3)	0.052 (3)	0.007 (3)	0.008 (3)	0.014 (3)
C5	0.035 (2)	0.043 (3)	0.043 (3)	0.006 (2)	0.007 (2)	0.019 (2)
C6	0.037 (2)	0.040 (3)	0.042 (3)	0.005 (2)	0.008 (2)	0.017 (2)
C7	0.052 (3)	0.060 (4)	0.046 (3)	0.017 (3)	0.015 (2)	0.026 (3)
C8	0.064 (4)	0.088 (6)	0.057 (4)	0.013 (4)	0.025 (3)	0.032 (4)
C9	0.045 (3)	0.054 (4)	0.063 (4)	0.008 (3)	0.018 (3)	0.003 (3)
C10	0.049 (3)	0.047 (3)	0.072 (4)	0.011 (3)	0.020 (3)	0.021 (3)

Geometric parameters (\AA , $^\circ$)

Cl1—Hg1	2.3875 (19)	C5—N1	1.323 (8)
Cl2—Hg1	2.4579 (19)	C5—N2	1.381 (8)
N1—Hg1	2.369 (6)	C6—N3	1.341 (8)
N2—H2B	0.95 (14)	C6—N2	1.383 (8)
N3—Hg1	2.290 (6)	C6—C7	1.398 (9)
C1—N1	1.349 (10)	C7—C8	1.397 (11)
C1—C2	1.363 (13)	C7—H7	0.9300
C1—H1	0.9300	C8—C9	1.352 (14)
C2—C3	1.390 (14)	C8—H8	0.9300
C2—H2	0.9300	C9—C10	1.369 (12)
C3—C4	1.372 (11)	C9—H9	0.9300
C3—H3	0.9300	C10—N3	1.362 (9)
C4—C5	1.399 (9)	C10—H10	0.9300
C4—H4	0.9300		
N1—Hg1—Cl1	112.30 (15)	N2—C6—C7	116.6 (5)

N1—Hg1—Cl2	94.92 (15)	C8—C7—C6	119.0 (7)
N3—Hg1—Cl1	112.21 (15)	C8—C7—H7	120.5
N3—Hg1—Cl2	123.92 (14)	C6—C7—H7	120.5
N3—Hg1—N1	82.4 (2)	C9—C8—C7	119.0 (7)
Cl1—Hg1—Cl2	120.14 (7)	C9—C8—H8	120.6
N1—C1—C2	123.8 (8)	C7—C8—H8	120.4
N1—C1—H1	118.1	C8—C9—C10	119.9 (7)
C2—C1—H1	118.1	C8—C9—H9	120.0
C1—C2—C3	117.9 (8)	C10—C9—H9	120.1
C1—C2—H2	121.0	N3—C10—C9	122.5 (7)
C3—C2—H2	121.0	N3—C10—H10	118.7
C4—C3—C2	118.7 (8)	C9—C10—H10	118.8
C4—C3—H3	120.6	C5—N1—C1	118.4 (7)
C2—C3—H3	120.7	C5—N1—Hg1	125.5 (4)
C3—C4—C5	119.9 (7)	C1—N1—Hg1	115.7 (5)
C3—C4—H4	120.1	C6—N2—C5	136.0 (5)
C5—C4—H4	120.0	C6—N2—H2B	109 (8)
N1—C5—N2	122.3 (6)	C5—N2—H2B	115 (8)
N1—C5—C4	121.1 (6)	C6—N3—C10	118.2 (6)
N2—C5—C4	116.6 (6)	C6—N3—Hg1	127.3 (4)
N3—C6—N2	122.1 (6)	C10—N3—Hg1	114.4 (5)
N3—C6—C7	121.3 (6)		
C1—N1—Hg1—N3	−169.7 (6)	C6—C7—C8—C9	2.0 (12)
C5—N1—Hg1—N3	17.3 (5)	C7—C8—C9—C10	−0.9 (13)
C1—N1—Hg1—Cl1	−58.7 (6)	C8—C9—C10—N3	0.6 (12)
C1—N1—Hg1—Cl2	66.7 (6)	N2—C5—N1—C1	179.1 (7)
C5—N1—Hg1—Cl1	128.2 (5)	C4—C5—N1—C1	−2.6 (10)
C5—N1—Hg1—Cl2	−106.3 (5)	N2—C5—N1—Hg1	−8.1 (9)
C6—N3—Hg1—N1	−15.5 (5)	C4—C5—N1—Hg1	170.3 (5)
C10—N3—Hg1—N1	168.0 (5)	C2—C1—N1—C5	1.9 (13)
C6—N3—Hg1—Cl1	−126.5 (5)	C2—C1—N1—Hg1	−171.6 (7)
C10—N3—Hg1—Cl1	57.0 (5)	N3—C6—N2—C5	17.7 (11)
C6—N3—Hg1—Cl2	75.3 (5)	C7—C6—N2—C5	−164.1 (7)
C10—N3—Hg1—Cl2	−101.2 (5)	N1—C5—N2—C6	−15.3 (11)
N1—C1—C2—C3	−1.4 (15)	C4—C5—N2—C6	166.3 (7)
C1—C2—C3—C4	1.5 (14)	N2—C6—N3—C10	−179.2 (6)
C2—C3—C4—C5	−2.3 (12)	C7—C6—N3—C10	2.7 (9)
C3—C4—C5—N1	2.8 (10)	N2—C6—N3—Hg1	4.5 (8)
C3—C4—C5—N2	−178.7 (7)	C7—C6—N3—Hg1	−173.7 (5)
N3—C6—C7—C8	−3.0 (10)	C9—C10—N3—C6	−1.5 (10)
N2—C6—C7—C8	178.7 (7)	C9—C10—N3—Hg1	175.3 (6)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2B···Cl2 ⁱ	0.95 (16)	2.41 (16)	3.345 (6)	169 (13)

Symmetry codes: (i) $-x, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

