

Triaqua(chlorido(1,10-phenanthroline- $\kappa^2 N,N'$)cobalt(II) chloride monohydrate

Yaru Liu,^{a*} Yutian Pan,^b Junshan Sun^c and Chuan Zhang^c

^aSchool of Science, North University of China, Taiyuan 030051, People's Republic of China, ^bSchool of Mechatronic Engineering, North University of China, Taiyuan 030051, People's Republic of China, and ^cDepartment of Materials Science and Chemical Engineering, Taishan University, 271021 Taian, Shandong, People's Republic of China

Correspondence e-mail: xiangyz_2008@163.com


Received 14 November 2008; accepted 15 December 2008

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.013\text{ \AA}$; R factor = 0.071; wR factor = 0.177; data-to-parameter ratio = 13.8.

In the title compound, $[\text{CoCl}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_3]\text{Cl}\cdot\text{H}_2\text{O}$, the Co^{II} ion is coordinated by two N atoms from the 1,10-phenanthroline ligand [$\text{Co}-\text{N} = 2.125(6)$ and $2.146(6)\text{ \AA}$], one chloride ligand [$\text{Co}-\text{Cl} = 2.459(2)\text{ \AA}$] and three water molecules [$\text{Co}-\text{O} = 2.070(5)$ – $2.105(5)\text{ \AA}$] in a distorted octahedral geometry. Intermolecular $\text{O}-\text{H}\cdots\text{Cl}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds form an extensive three-dimensional hydrogen-bonding network, which consolidates the crystal packing.

Related literature

For related crystal structures, see: Liu *et al.* (2006); Zhang *et al.* (1999).



Experimental

Crystal data

$[\text{CoCl}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_3]\text{Cl}\cdot\text{H}_2\text{O}$

$M_r = 382.10$

Monoclinic, $P2_1/c$
 $a = 7.646(4)\text{ \AA}$
 $b = 12.426(6)\text{ \AA}$
 $c = 16.987(8)\text{ \AA}$
 $\beta = 103.54(2)^\circ$
 $V = 1569.1(13)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.45\text{ mm}^{-1}$
 $T = 273(2)\text{ K}$
 $0.37 \times 0.25 \times 0.19\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{min} = 0.616$, $T_{max} = 0.770$

6533 measured reflections
2733 independent reflections
1564 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.080$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.177$
 $S = 1.04$
2733 reflections
198 parameters

13 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.67\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.56\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A \cdots O4	0.85	1.93	2.693 (9)	149
O1—H1B \cdots Cl2	0.85	2.55	3.258 (6)	141
O2—H2A \cdots Cl1 ⁱ	0.85	2.31	3.139 (6)	166
O2—H2B \cdots Cl2 ⁱ	0.85	2.29	3.119 (6)	164
O3—H3A \cdots Cl2 ⁱⁱ	0.85	2.33	3.114 (6)	153
O3—H3B \cdots Cl1 ⁱⁱ	0.85	2.30	3.128 (6)	166
O4—H4A \cdots Cl2 ⁱⁱⁱ	0.85	2.34	3.185 (8)	170
O4—H4B \cdots Cl1 ⁱⁱ	0.85	2.58	3.278 (8)	141

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $-x + 2, -y + 1, -z + 2$; (iii) $-x + 2, y + \frac{1}{2}, -z + \frac{5}{2}$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2481).

References

- Liu, J. T., Fan, S. D. & Li, D. Q. (2006). *Acta Cryst. E62*, m2165–m2166.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Zhang, C., Yu, K., Wu, D. & Zhao, C. (1999). *Acta Cryst. C55*, 1815–1817.

supplementary materials

Acta Cryst. (2009). E65, m114 [doi:10.1107/S1600536808042591]

Triaquaclorido(1,10-phenanthroline- κ^2N,N')cobalt(II) chloride monohydrate

Y. Liu, Y. Pan, J. Sun and C. Zhang

Comment

The structures of some tetraqua(1,10-phenanthroline) metal ionic complexes with different anions have been reported (Liu *et al.*, 2006; Zhang *et al.*, 1999). In our study in this field, we select 1,10-phenanthroline as the co-ligand to continue our exploration of the Co complexes. Herein we report the structure of the title compound (I), which was characterized by elemental analyses and X-ray crystallography diffraction.

In (I) (Fig. 1), each Co^{II} ion is coordinated by two N atoms from 1,10-phenanthroline ligand [$\text{Co}-\text{N}$ 2.125 (6), 2.146 (6) Å], one chlorine anion [$\text{Co}-\text{Cl}$ 2.459 (2) Å] and three water molecules [$\text{Co}-\text{O}$ 2.070 (5)–2.105 (5) Å] in a distorted octahedral geometry. Two aqua O atoms and two N atoms from 1,10-phenanthroline ligand define the special position on a mirror plane. One aqua group and chlorine anion fill the axial apical positions. The intermolecular O—H···Cl and O—H···O hydrogen bonds (Table 1) form an extensive three-dimensional hydrogen-bonding network, which consolidate the crystal packing.

Experimental

A mixture of $\text{Cu}(\text{Cl})_2 \cdot 3\text{H}_2\text{O}$ (168 mg, 1 mmol) and 1,10-phenanthroline (185 mg, 1 mmol) in methanol (30 ml) was placed in a Teflon-lined stainless steel Parr bomb that was heated at 423 K for 72 h. The bomb was then cooled down to the room temperature, the solution was filtered. The solvent was removed from the filtrate under vacuum, and the solid residue was recrystallized from diethyl ether; blue crystals suitable for X-Ray diffraction study were obtained. Yield, 0.724 g, 81%. m.p. 566 K. Analysis, calculated for $\text{C}_{12}\text{H}_{16}\text{Cl}_2\text{CoN}_2\text{O}_4$: C 37.72, H 4.22, N 7.33; found: C 37.53, H 4.68, N 7.17%. The elemental analyses were performed with a Perkin Elmer PE2400II instrument.

Refinement

Water H atoms were located in a difference map and were isotropically refined with an O—H strict distance restraint of 0.85/%A. H atoms bound to C atoms were placed in calculated positions ($\text{C}-\text{H} = 0.93/\text{\AA}$) and refined in the riding-model approximation, $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The structure of the title complex, showing 30% probability displacement ellipsoids and the atom-numbering scheme. Dashed lines denote hydrogen bonds.

supplementary materials

Triaquaclorido(1,10-phenanthroline- κ^2 N,N')cobalt(II) chloride monohydrate

Crystal data

[CoCl(C ₁₂ H ₈ N ₂)(H ₂ O) ₃]Cl·H ₂ O	$F_{000} = 780$
$M_r = 382.10$	$D_x = 1.617 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.646 (4) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 12.426 (6) \text{ \AA}$	Cell parameters from 580 reflections
$c = 16.987 (8) \text{ \AA}$	$\theta = 3.0\text{--}20.1^\circ$
$\beta = 103.54 (2)^\circ$	$\mu = 1.45 \text{ mm}^{-1}$
$V = 1569.1 (13) \text{ \AA}^3$	$T = 273 (2) \text{ K}$
$Z = 4$	Block, blue
	$0.37 \times 0.25 \times 0.19 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	2733 independent reflections
Radiation source: fine-focus sealed tube	1564 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.080$
$T = 273(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 5$
$T_{\text{min}} = 0.616$, $T_{\text{max}} = 0.770$	$k = -10 \rightarrow 14$
6533 measured reflections	$l = -17 \rightarrow 20$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.071$	H-atom parameters constrained
$wR(F^2) = 0.177$	$w = 1/[\sigma^2(F_o^2) + (0.0517P)^2 + 4.0479P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2733 reflections	$\Delta\rho_{\text{max}} = 0.67 \text{ e \AA}^{-3}$
198 parameters	$\Delta\rho_{\text{min}} = -0.56 \text{ e \AA}^{-3}$
13 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.75359 (13)	0.66886 (9)	0.99070 (6)	0.0342 (3)
Cl1	0.6997 (3)	0.49910 (16)	0.91423 (11)	0.0431 (5)
Cl2	0.8145 (3)	0.34056 (18)	1.16347 (12)	0.0536 (6)
N1	0.6918 (8)	0.7697 (5)	0.8867 (4)	0.0360 (15)
N2	0.7887 (8)	0.8255 (5)	1.0452 (4)	0.0405 (16)
O1	0.8190 (8)	0.5892 (4)	1.1032 (3)	0.0478 (15)
H1A	0.9003	0.5920	1.1469	0.15 (6)*
H1B	0.7797	0.5251	1.0958	0.10 (4)*
O2	0.4845 (7)	0.6729 (4)	0.9965 (3)	0.0459 (13)
H2A	0.4527	0.6214	1.0231	0.06 (3)*
H2B	0.4185	0.6755	0.9488	0.05 (3)*
O3	1.0230 (7)	0.6657 (4)	0.9883 (3)	0.0456 (14)
H3A	1.0372	0.6788	0.9411	0.05 (3)*
H3B	1.1121	0.6289	1.0137	0.09 (4)*
O4	1.1261 (10)	0.6517 (6)	1.2062 (4)	0.075 (2)
H4A	1.1566	0.7013	1.2412	0.06 (3)*
H4B	1.2171	0.6296	1.1901	0.16 (7)*
C1	0.6476 (11)	0.7393 (8)	0.8103 (5)	0.049 (2)
H1	0.6394	0.6664	0.7979	0.059*
C2	0.6123 (11)	0.8160 (7)	0.7468 (5)	0.050 (2)
H2	0.5810	0.7933	0.6931	0.060*
C3	0.6243 (11)	0.9222 (7)	0.7645 (5)	0.052 (2)
H3	0.6017	0.9726	0.7228	0.062*
C4	0.6706 (11)	0.9573 (7)	0.8456 (5)	0.046 (2)
C5	0.7032 (10)	0.8787 (6)	0.9056 (5)	0.037 (2)
C6	0.7525 (9)	0.9047 (6)	0.9885 (5)	0.0338 (17)
C7	0.7636 (10)	1.0150 (7)	1.0107 (5)	0.044 (2)
C8	0.8142 (12)	1.0398 (8)	1.0931 (6)	0.062 (3)
H8	0.8232	1.1111	1.1102	0.075*
C9	0.8501 (13)	0.9583 (8)	1.1481 (6)	0.062 (3)
H9	0.8841	0.9740	1.2031	0.074*
C10	0.8363 (10)	0.8527 (7)	1.1225 (5)	0.049 (2)
H10	0.8616	0.7985	1.1612	0.058*

supplementary materials

C11	0.6849 (12)	1.0667 (7)	0.8682 (6)	0.058 (3)
H11	0.6645	1.1199	0.8285	0.070*
C12	0.7284 (13)	1.0944 (7)	0.9475 (6)	0.063 (3)
H12	0.7357	1.1669	0.9614	0.075*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0399 (6)	0.0288 (6)	0.0335 (6)	-0.0003 (5)	0.0080 (4)	0.0012 (5)
Cl1	0.0576 (13)	0.0299 (11)	0.0394 (12)	-0.0009 (9)	0.0068 (9)	-0.0034 (10)
Cl2	0.0673 (14)	0.0495 (14)	0.0443 (12)	0.0020 (11)	0.0133 (10)	0.0060 (12)
N1	0.040 (4)	0.033 (4)	0.036 (4)	0.000 (3)	0.011 (3)	0.002 (3)
N2	0.043 (4)	0.034 (4)	0.045 (4)	-0.001 (3)	0.009 (3)	-0.003 (4)
O1	0.068 (4)	0.040 (4)	0.034 (3)	-0.006 (3)	0.009 (3)	0.002 (3)
O2	0.050 (3)	0.042 (4)	0.048 (3)	0.000 (3)	0.016 (3)	0.004 (3)
O3	0.043 (3)	0.055 (4)	0.042 (3)	0.006 (3)	0.015 (2)	0.007 (3)
O4	0.080 (5)	0.080 (5)	0.058 (4)	0.008 (4)	0.001 (4)	-0.028 (4)
C1	0.058 (6)	0.059 (6)	0.030 (5)	0.005 (4)	0.010 (4)	0.004 (4)
C2	0.061 (6)	0.055 (6)	0.032 (4)	0.003 (4)	0.008 (4)	0.004 (4)
C3	0.062 (6)	0.040 (6)	0.052 (6)	0.003 (4)	0.012 (4)	0.005 (5)
C4	0.046 (5)	0.044 (6)	0.047 (5)	0.005 (4)	0.009 (4)	0.006 (4)
C5	0.036 (4)	0.025 (4)	0.055 (6)	0.000 (3)	0.018 (4)	-0.009 (4)
C6	0.038 (4)	0.022 (4)	0.042 (5)	-0.001 (3)	0.010 (3)	-0.001 (4)
C7	0.043 (5)	0.032 (5)	0.057 (6)	0.001 (4)	0.013 (4)	-0.005 (4)
C8	0.078 (7)	0.034 (6)	0.074 (7)	-0.002 (5)	0.014 (5)	-0.023 (5)
C9	0.080 (7)	0.045 (6)	0.059 (6)	-0.001 (5)	0.016 (5)	-0.021 (5)
C10	0.053 (5)	0.054 (6)	0.038 (5)	0.004 (4)	0.009 (4)	-0.004 (5)
C11	0.072 (7)	0.038 (6)	0.062 (7)	0.009 (5)	0.012 (5)	0.020 (5)
C12	0.075 (7)	0.030 (5)	0.082 (7)	0.002 (5)	0.013 (5)	0.004 (6)

Geometric parameters (\AA , $^\circ$)

Co1—O3	2.071 (5)	C1—H1	0.9300
Co1—O2	2.084 (5)	C2—C3	1.35 (1)
Co1—O1	2.106 (5)	C2—H2	0.9301
Co1—N1	2.127 (6)	C3—C4	1.41 (1)
Co1—N2	2.145 (7)	C3—H3	0.9300
Co1—Cl1	2.461 (2)	C4—C5	1.39 (1)
N1—C1	1.317 (9)	C4—C11	1.41 (1)
N1—C5	1.390 (9)	C5—C6	1.41 (1)
N2—C10	1.321 (9)	C6—C7	1.42 (1)
N2—C6	1.361 (9)	C7—C8	1.40 (1)
O1—H1A	0.8500	C7—C12	1.44 (1)
O1—H1B	0.8501	C8—C9	1.36 (1)
O2—H2A	0.8500	C8—H8	0.9301
O2—H2B	0.8498	C9—C10	1.38 (1)
O3—H3A	0.8500	C9—H9	0.9301
O3—H3B	0.8500	C10—H10	0.9300
O4—H4A	0.8500	C11—C12	1.36 (1)

O4—H4B	0.8500	C11—H11	0.9300
C1—C2	1.42 (1)	C12—H12	0.9300
O3—Co1—O2	178.4 (2)	C1—C2—H2	120.2
O3—Co1—O1	89.0 (2)	C2—C3—C4	120.5 (8)
O2—Co1—O1	89.7 (2)	C2—C3—H3	119.7
O3—Co1—N1	91.3 (2)	C4—C3—H3	119.7
O2—Co1—N1	89.8 (2)	C5—C4—C3	117.4 (8)
O1—Co1—N1	171.9 (2)	C5—C4—C11	119.3 (8)
O3—Co1—N2	90.1 (2)	C3—C4—C11	123.3 (8)
O2—Co1—N2	89.0 (2)	N1—C5—C4	121.6 (8)
O1—Co1—N2	93.2 (2)	N1—C5—C6	116.3 (7)
N1—Co1—N2	78.8 (2)	C4—C5—C6	122.0 (7)
C1—N1—C5	119.6 (7)	N2—C6—C5	120.3 (7)
C1—N1—Co1	127.3 (6)	N2—C6—C7	121.4 (7)
C5—N1—Co1	113.1 (5)	C5—C6—C7	118.3 (7)
C10—N2—C6	118.8 (7)	C8—C7—C6	117.7 (8)
C10—N2—Co1	129.7 (6)	C8—C7—C12	123.8 (8)
C6—N2—Co1	111.5 (5)	C6—C7—C12	118.4 (8)
Co1—O1—H1A	138.0	C9—C8—C7	119.2 (8)
Co1—O1—H1B	107.8	C9—C8—H8	120.4
H1A—O1—H1B	109.2	C7—C8—H8	120.4
Co1—O2—H2A	114.5	C8—C9—C10	120.2 (9)
Co1—O2—H2B	109.1	C8—C9—H9	119.9
H2A—O2—H2B	110.8	C10—C9—H9	119.9
Co1—O3—H3A	111.3	N2—C10—C9	122.7 (9)
Co1—O3—H3B	133.1	N2—C10—H10	118.7
H3A—O3—H3B	108.6	C9—C10—H10	118.7
H4A—O4—H4B	110.4	C12—C11—C4	120.0 (8)
N1—C1—C2	121.2 (8)	C12—C11—H11	120.0
N1—C1—H1	119.4	C4—C11—H11	120.0
C2—C1—H1	119.4	C11—C12—C7	121.9 (9)
C3—C2—C1	119.7 (8)	C11—C12—H12	119.1
C3—C2—H2	120.2	C7—C12—H12	119.1

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1A···O4	0.85	1.93	2.693 (9)	149
O1—H1B···Cl2	0.85	2.55	3.258 (6)	141
O2—H2A···Cl1 ⁱ	0.85	2.31	3.139 (6)	166
O2—H2B···Cl2 ⁱ	0.85	2.29	3.119 (6)	164
O3—H3A···Cl2 ⁱⁱ	0.85	2.33	3.114 (6)	153
O3—H3B···Cl1 ⁱⁱ	0.85	2.30	3.128 (6)	166
O4—H4A···Cl2 ⁱⁱⁱ	0.85	2.34	3.185 (8)	170
O4—H4B···Cl1 ⁱⁱ	0.85	2.58	3.278 (8)	141

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $-x+2, -y+1, -z+2$; (iii) $-x+2, y+1/2, -z+5/2$.

supplementary materials

Fig. 1

