

**[*N'*-(4-Methoxy-2-oxidobenzylidene)-4-nitrobenzohydrazidato- $\kappa^3 O, N, O'$ ]-  
(pyridine- $\kappa N$ )copper(II)**

Nooraziah Mohd Lair, Hapipah Mohd Ali and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 15 December 2008; accepted 16 December 2008

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(C-C) = 0.003$  Å;  
 $R$  factor = 0.028;  $wR$  factor = 0.087; data-to-parameter ratio = 14.5.

The pyridine-coordinated Cu<sup>II</sup> atom in the title Schiff base complex, [Cu(C<sub>15</sub>H<sub>11</sub>N<sub>3</sub>O<sub>5</sub>)(C<sub>5</sub>H<sub>5</sub>N)], is *O,N,O'*-chelated by the doubly deprotonated Schiff base ligand. The metal centre is in a square-planar coordination geometry.

### Related literature

For the pyridine adducts of copper derivatives of similar ligands, see: Ali *et al.* (2004); Chen & Liu (2004); Das & Pal (2005); Fariati *et al.* (2002); Lu & Liu (2005); Lu *et al.* (2003).



### Experimental

#### Crystal data

[Cu(C<sub>15</sub>H<sub>11</sub>N<sub>3</sub>O<sub>5</sub>)(C<sub>5</sub>H<sub>5</sub>N)]  
 $M_r = 455.91$   
Triclinic,  $P\bar{1}$   
 $a = 6.3529$  (1) Å  
 $b = 9.8409$  (2) Å  
 $c = 15.1303$  (3) Å  
 $\alpha = 98.063$  (1)°  
 $\beta = 92.011$  (1)°

$\gamma = 107.088$  (1)°  
 $V = 892.31$  (3) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 1.27$  mm<sup>-1</sup>  
 $T = 100$  (2) K  
 $0.40 \times 0.10 \times 0.05$  mm

#### Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.807$ ,  $T_{\max} = 1.000$   
(expected range = 0.757–0.939)

6268 measured reflections  
3942 independent reflections  
3605 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.013$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$   
 $wR(F^2) = 0.087$   
 $S = 1.09$   
3942 reflections

272 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.45$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.32$  e Å<sup>-3</sup>

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2009).

We thank the University of Malaya for funding this study (Science Fund grants 12–02–03–2031, 12–02–03–2051).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2838).

### References

- Ali, H., Khamis, N. A., Basirun, W. J. & Yamin, B. M. (2004). *Acta Cryst.* **E60**, m982–m983.  
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Chen, W.-H. & Liu, S.-X. (2004). *Chin. J. Inorg. Chem.* **20**, 668–670.  
Das, S. & Pal, S. (2005). *J. Mol. Struct.* **753**, 68–79.  
Fariati, F., Caruso, U., Centore, R., Marcolli, W., De Maria, A., Panunzi, B., Roviello, A. & Tuzi, A. (2002). *Inorg. Chem.* **41**, 6597–6603.  
Lu, W.-G., Feng, X.-L., Liu, H.-W. & Wang, S.-L. (2003). *Chin. J. Inorg. Chem.* **19**, 206–210.  
Lu, W.-G. & Liu, H.-W. (2005). *Chin. J. Struct. Chem.* **24**, 1078–1082.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2009). *pubCIF*. In preparation.

**supplementary materials**

*Acta Cryst.* (2009). E65, m121 [ doi:10.1107/S1600536808042803 ]

**[N'-(4-Methoxy-2-oxidobenzylidene)4-nitrobenzohydrazidato- $\kappa^3O,N,O'$ ](pyridine- $\kappa N$ )copper(II)**

**N. Mohd Lair, H. Mohd Ali and S. W. Ng**

**Comment**

(type here to add)

**Experimental**

N'-2-Hydroxy-3-methoxybenzylidene)-nitrobenzohydrazide (0.30 g, 1 mmol) and copper acetate (0.20 g, 1 mmol) were heated in a ethanol (50 ml) for 2 hours. The solvent was removed and the resulting compound recrystallized from pyridine.

**Refinement**

Hydrogen atoms were placed at calculated positions ( $C_{\text{aromatic-H}} 0.95 \text{ \AA}$ ,  $C_{\text{methyl-H}} 0.98 \text{ \AA}$ ) and were treated as riding on their parent carbon atoms, with  $U(\text{H})$  set to  $1.2U_{\text{eq}}(C_{\text{aromatic}})$  or  $1.5U_{\text{eq}}(C_{\text{methyl}})$ .

**Figures**




Fig. 1. Displacement ellipsoid plot (Barbour, 2001) of  $\text{Cu}(\text{C}_5\text{H}_5\text{N})(\text{C}_{15}\text{H}_{11}\text{N}_3\text{O}_5)$  at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

**[N'-(4-Methoxy-2-oxidobenzylidene)-4-nitrobenzohydrazidato-  $\kappa^3O,N,O'$ ](pyridine- $\kappa N$ )copper(II)**

*Crystal data*

$[\text{Cu}(\text{C}_{15}\text{H}_{11}\text{N}_3\text{O}_5)(\text{C}_5\text{H}_5\text{N})]$

$M_r = 455.91$

Triclinic,  $PT$

Hall symbol:  $-P 1$

$a = 6.3529 (1) \text{ \AA}$

$b = 9.8409 (2) \text{ \AA}$

$c = 15.1303 (3) \text{ \AA}$

$\alpha = 98.063 (1)^\circ$

$\beta = 92.011 (1)^\circ$

$\gamma = 107.088 (1)^\circ$

$V = 892.31 (3) \text{ \AA}^3$

$Z = 2$

$F_{000} = 466$

$D_x = 1.697 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3716 reflections

$\theta = 2.4\text{--}29.2^\circ$

$\mu = 1.27 \text{ mm}^{-1}$

$T = 100 (2) \text{ K}$

Block, brown

$0.40 \times 0.10 \times 0.05 \text{ mm}$

# supplementary materials

---

## Data collection

Bruker SMART APEX diffractometer	3942 independent reflections
Radiation source: fine-focus sealed tube	3605 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.013$
$T = 100(2)$ K	$\theta_{\text{max}} = 27.5^\circ$
$\omega$ scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -6 \rightarrow 8$
$T_{\text{min}} = 0.807$ , $T_{\text{max}} = 1.000$	$k = -12 \rightarrow 12$
6268 measured reflections	$l = -19 \rightarrow 19$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.087$	$w = 1/[\sigma^2(F_o^2) + (0.0494P)^2 + 0.535P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
3942 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
272 parameters	$\Delta\rho_{\text{max}} = 0.45 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.31 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	1.14506 (4)	0.70062 (2)	0.728016 (14)	0.01312 (9)
N1	0.9381 (3)	0.64556 (17)	0.62386 (11)	0.0150 (3)
N2	1.0273 (3)	0.68298 (18)	0.54424 (10)	0.0159 (3)
N3	1.7555 (3)	0.93172 (18)	0.27544 (11)	0.0189 (4)
N4	1.3792 (3)	0.78081 (17)	0.82931 (11)	0.0145 (3)
O1	0.9305 (2)	0.61658 (15)	0.80405 (9)	0.0163 (3)
O2	0.2533 (2)	0.39329 (16)	0.90700 (9)	0.0199 (3)
O3	1.3444 (2)	0.77637 (15)	0.64095 (9)	0.0159 (3)
O4	1.6632 (3)	0.91481 (17)	0.20033 (9)	0.0247 (3)
O5	1.9560 (3)	0.9856 (2)	0.29243 (11)	0.0311 (4)
C1	0.7183 (3)	0.5521 (2)	0.78294 (13)	0.0139 (4)
C2	0.5869 (3)	0.5009 (2)	0.85092 (13)	0.0152 (4)
H2	0.6540	0.5120	0.9097	0.018*
C3	0.3615 (3)	0.4346 (2)	0.83442 (13)	0.0149 (4)
C4	0.2562 (3)	0.4147 (2)	0.74829 (13)	0.0151 (4)
H4	0.1016	0.3693	0.7371	0.018*
C5	0.3837 (3)	0.4630 (2)	0.68048 (13)	0.0146 (4)

H5	0.3142	0.4495	0.6219	0.018*
C6	0.6124 (3)	0.5313 (2)	0.69468 (12)	0.0140 (4)
C7	0.7290 (3)	0.5785 (2)	0.61964 (12)	0.0149 (4)
H7	0.6471	0.5592	0.5631	0.018*
C8	0.0219 (3)	0.3204 (3)	0.89270 (15)	0.0247 (5)
H8A	-0.0352	0.2939	0.9492	0.037*
H8B	-0.0038	0.2333	0.8484	0.037*
H8C	-0.0540	0.3840	0.8707	0.037*
C9	1.2393 (3)	0.7493 (2)	0.56222 (12)	0.0140 (4)
C10	1.3714 (3)	0.79538 (19)	0.48671 (12)	0.0142 (4)
C11	1.2684 (3)	0.7902 (2)	0.40223 (13)	0.0162 (4)
H11	1.1119	0.7555	0.3927	0.019*
C12	1.3932 (3)	0.8352 (2)	0.33284 (13)	0.0174 (4)
H12	1.3242	0.8321	0.2755	0.021*
C13	1.6204 (3)	0.8849 (2)	0.34864 (12)	0.0149 (4)
C14	1.7277 (3)	0.8905 (2)	0.43096 (13)	0.0162 (4)
H14	1.8844	0.9243	0.4396	0.019*
C15	1.6016 (3)	0.8457 (2)	0.50049 (12)	0.0146 (4)
H15	1.6720	0.8492	0.5576	0.017*
C16	1.3332 (3)	0.7615 (2)	0.91374 (13)	0.0189 (4)
H16	1.1851	0.7146	0.9246	0.023*
C17	1.4910 (3)	0.8068 (2)	0.98487 (13)	0.0219 (4)
H17	1.4524	0.7913	1.0435	0.026*
C18	1.7074 (3)	0.8753 (2)	0.96950 (14)	0.0210 (4)
H18	1.8197	0.9079	1.0175	0.025*
C19	1.7569 (3)	0.8955 (2)	0.88305 (13)	0.0189 (4)
H19	1.9041	0.9416	0.8707	0.023*
C20	1.5893 (3)	0.8476 (2)	0.81504 (13)	0.0161 (4)
H20	1.6239	0.8625	0.7559	0.019*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.01168 (13)	0.01707 (13)	0.00996 (13)	0.00263 (9)	0.00095 (8)	0.00366 (8)
N1	0.0165 (8)	0.0172 (8)	0.0117 (7)	0.0044 (6)	0.0032 (6)	0.0046 (6)
N2	0.0166 (8)	0.0193 (8)	0.0109 (7)	0.0027 (7)	0.0034 (6)	0.0048 (6)
N3	0.0245 (9)	0.0184 (8)	0.0147 (8)	0.0060 (7)	0.0075 (7)	0.0047 (6)
N4	0.0133 (8)	0.0162 (8)	0.0137 (8)	0.0039 (6)	0.0007 (6)	0.0032 (6)
O1	0.0109 (6)	0.0234 (7)	0.0136 (6)	0.0025 (5)	0.0012 (5)	0.0057 (5)
O2	0.0136 (7)	0.0297 (8)	0.0154 (7)	0.0027 (6)	0.0032 (5)	0.0082 (6)
O3	0.0139 (7)	0.0218 (7)	0.0110 (6)	0.0028 (5)	0.0006 (5)	0.0046 (5)
O4	0.0325 (9)	0.0307 (8)	0.0127 (7)	0.0103 (7)	0.0053 (6)	0.0074 (6)
O5	0.0224 (8)	0.0431 (10)	0.0229 (8)	-0.0003 (7)	0.0092 (7)	0.0092 (7)
C1	0.0131 (9)	0.0140 (8)	0.0154 (9)	0.0050 (7)	0.0014 (7)	0.0030 (7)
C2	0.0143 (9)	0.0192 (9)	0.0130 (9)	0.0056 (8)	0.0009 (7)	0.0041 (7)
C3	0.0156 (9)	0.0160 (9)	0.0141 (9)	0.0048 (7)	0.0042 (7)	0.0046 (7)
C4	0.0125 (9)	0.0156 (9)	0.0157 (9)	0.0025 (7)	0.0007 (7)	0.0018 (7)
C5	0.0167 (9)	0.0141 (8)	0.0124 (8)	0.0041 (7)	-0.0001 (7)	0.0011 (7)

## supplementary materials

---

C6	0.0156 (9)	0.0133 (8)	0.0127 (8)	0.0040 (7)	0.0021 (7)	0.0021 (7)
C7	0.0164 (9)	0.0153 (9)	0.0115 (8)	0.0027 (7)	-0.0009 (7)	0.0024 (7)
C8	0.0126 (10)	0.0395 (13)	0.0216 (10)	0.0032 (9)	0.0058 (8)	0.0122 (9)
C9	0.0169 (9)	0.0146 (9)	0.0116 (8)	0.0055 (7)	0.0025 (7)	0.0038 (7)
C10	0.0171 (9)	0.0129 (8)	0.0130 (9)	0.0046 (7)	0.0022 (7)	0.0034 (7)
C11	0.0146 (9)	0.0202 (9)	0.0135 (9)	0.0045 (7)	0.0005 (7)	0.0033 (7)
C12	0.0211 (10)	0.0199 (9)	0.0111 (9)	0.0057 (8)	0.0007 (7)	0.0033 (7)
C13	0.0197 (10)	0.0138 (9)	0.0116 (8)	0.0048 (7)	0.0055 (7)	0.0035 (7)
C14	0.0154 (9)	0.0159 (9)	0.0166 (9)	0.0035 (7)	0.0019 (7)	0.0021 (7)
C15	0.0168 (9)	0.0163 (9)	0.0111 (8)	0.0050 (7)	0.0014 (7)	0.0037 (7)
C16	0.0148 (10)	0.0249 (10)	0.0147 (9)	0.0021 (8)	0.0019 (7)	0.0031 (8)
C17	0.0193 (10)	0.0293 (11)	0.0125 (9)	0.0008 (9)	0.0010 (8)	0.0029 (8)
C18	0.0186 (10)	0.0248 (10)	0.0149 (9)	0.0007 (8)	-0.0034 (8)	0.0013 (8)
C19	0.0143 (9)	0.0197 (10)	0.0199 (10)	0.0008 (8)	0.0009 (8)	0.0033 (8)
C20	0.0156 (9)	0.0176 (9)	0.0143 (9)	0.0033 (7)	0.0026 (7)	0.0036 (7)

### *Geometric parameters (Å, °)*

Cu1—O1	1.8922 (14)	C6—C7	1.435 (3)
Cu1—N1	1.9239 (16)	C7—H7	0.9500
Cu1—O3	1.9320 (14)	C8—H8A	0.9800
Cu1—N4	1.9989 (16)	C8—H8B	0.9800
N1—C7	1.293 (3)	C8—H8C	0.9800
N1—N2	1.399 (2)	C9—C10	1.485 (3)
N2—C9	1.312 (3)	C10—C15	1.397 (3)
N3—O5	1.229 (2)	C10—C11	1.403 (3)
N3—O4	1.227 (2)	C11—C12	1.382 (3)
N3—C13	1.467 (2)	C11—H11	0.9500
N4—C20	1.343 (3)	C12—C13	1.381 (3)
N4—C16	1.347 (2)	C12—H12	0.9500
O1—C1	1.316 (2)	C13—C14	1.385 (3)
O2—C3	1.363 (2)	C14—C15	1.387 (3)
O2—C8	1.427 (2)	C14—H14	0.9500
O3—C9	1.299 (2)	C15—H15	0.9500
C1—C2	1.403 (3)	C16—C17	1.375 (3)
C1—C6	1.434 (3)	C16—H16	0.9500
C2—C3	1.385 (3)	C17—C18	1.386 (3)
C2—H2	0.9500	C17—H17	0.9500
C3—C4	1.405 (3)	C18—C19	1.384 (3)
C4—C5	1.380 (3)	C18—H18	0.9500
C4—H4	0.9500	C19—C20	1.381 (3)
C5—C6	1.404 (3)	C19—H19	0.9500
C5—H5	0.9500	C20—H20	0.9500
O1—Cu1—N1	93.57 (6)	O2—C8—H8B	109.5
O1—Cu1—O3	174.58 (6)	H8A—C8—H8B	109.5
N1—Cu1—O3	81.17 (6)	O2—C8—H8C	109.5
O1—Cu1—N4	92.67 (6)	H8A—C8—H8C	109.5
N1—Cu1—N4	172.51 (7)	H8B—C8—H8C	109.5
O3—Cu1—N4	92.68 (6)	O3—C9—N2	125.19 (17)

C7—N1—N2	117.25 (16)	O3—C9—C10	117.01 (17)
C7—N1—Cu1	127.41 (13)	N2—C9—C10	117.79 (16)
N2—N1—Cu1	115.34 (12)	C15—C10—C11	119.71 (17)
C9—N2—N1	108.05 (15)	C15—C10—C9	119.31 (16)
O5—N3—O4	123.25 (17)	C11—C10—C9	120.98 (18)
O5—N3—C13	118.27 (16)	C12—C11—C10	120.41 (18)
O4—N3—C13	118.47 (17)	C12—C11—H11	119.8
C20—N4—C16	117.86 (17)	C10—C11—H11	119.8
C20—N4—Cu1	121.38 (13)	C11—C12—C13	118.53 (17)
C16—N4—Cu1	120.64 (14)	C11—C12—H12	120.7
C1—O1—Cu1	127.54 (12)	C13—C12—H12	120.7
C3—O2—C8	117.33 (15)	C12—C13—C14	122.62 (18)
C9—O3—Cu1	110.23 (12)	C12—C13—N3	119.26 (17)
O1—C1—C2	118.11 (17)	C14—C13—N3	118.12 (18)
O1—C1—C6	124.11 (17)	C15—C14—C13	118.62 (18)
C2—C1—C6	117.78 (17)	C15—C14—H14	120.7
C3—C2—C1	121.55 (17)	C13—C14—H14	120.7
C3—C2—H2	119.2	C14—C15—C10	120.11 (17)
C1—C2—H2	119.2	C14—C15—H15	119.9
O2—C3—C2	115.33 (17)	C10—C15—H15	119.9
O2—C3—C4	123.65 (18)	N4—C16—C17	122.91 (19)
C2—C3—C4	121.01 (18)	N4—C16—H16	118.5
C5—C4—C3	118.12 (18)	C17—C16—H16	118.5
C5—C4—H4	120.9	C16—C17—C18	118.84 (19)
C3—C4—H4	120.9	C16—C17—H17	120.6
C4—C5—C6	122.53 (17)	C18—C17—H17	120.6
C4—C5—H5	118.7	C19—C18—C17	118.81 (19)
C6—C5—H5	118.7	C19—C18—H18	120.6
C5—C6—C1	119.00 (17)	C17—C18—H18	120.6
C5—C6—C7	117.98 (17)	C20—C19—C18	119.05 (19)
C1—C6—C7	123.02 (17)	C20—C19—H19	120.5
N1—C7—C6	124.30 (17)	C18—C19—H19	120.5
N1—C7—H7	117.9	N4—C20—C19	122.52 (18)
C6—C7—H7	117.9	N4—C20—H20	118.7
O2—C8—H8A	109.5	C19—C20—H20	118.7
O1—Cu1—N1—C7	1.77 (18)	C5—C6—C7—N1	178.35 (18)
O3—Cu1—N1—C7	-179.56 (18)	C1—C6—C7—N1	-1.0 (3)
O1—Cu1—N1—N2	-178.07 (13)	Cu1—O3—C9—N2	1.5 (2)
O3—Cu1—N1—N2	0.60 (12)	Cu1—O3—C9—C10	-177.59 (12)
C7—N1—N2—C9	-179.91 (17)	N1—N2—C9—O3	-1.0 (3)
Cu1—N1—N2—C9	-0.05 (19)	N1—N2—C9—C10	178.08 (15)
O1—Cu1—N4—C20	174.53 (15)	O3—C9—C10—C15	9.0 (3)
O3—Cu1—N4—C20	-4.54 (15)	N2—C9—C10—C15	-170.09 (17)
O1—Cu1—N4—C16	-1.39 (16)	O3—C9—C10—C11	-170.48 (17)
O3—Cu1—N4—C16	179.54 (15)	N2—C9—C10—C11	10.4 (3)
N1—Cu1—O1—C1	-2.34 (16)	C15—C10—C11—C12	-0.4 (3)
N4—Cu1—O1—C1	173.51 (16)	C9—C10—C11—C12	179.15 (17)
N1—Cu1—O3—C9	-1.01 (12)	C10—C11—C12—C13	0.1 (3)
N4—Cu1—O3—C9	-176.72 (13)	C11—C12—C13—C14	0.4 (3)

## supplementary materials

---

Cu1—O1—C1—C2	-178.09 (13)	C11—C12—C13—N3	179.37 (17)
Cu1—O1—C1—C6	1.7 (3)	O5—N3—C13—C12	174.64 (18)
O1—C1—C2—C3	178.63 (17)	O4—N3—C13—C12	-5.6 (3)
C6—C1—C2—C3	-1.1 (3)	O5—N3—C13—C14	-6.3 (3)
C8—O2—C3—C2	-178.07 (18)	O4—N3—C13—C14	173.44 (18)
C8—O2—C3—C4	2.8 (3)	C12—C13—C14—C15	-0.6 (3)
C1—C2—C3—O2	-178.47 (17)	N3—C13—C14—C15	-179.62 (16)
C1—C2—C3—C4	0.7 (3)	C13—C14—C15—C10	0.4 (3)
O2—C3—C4—C5	179.17 (17)	C11—C10—C15—C14	0.1 (3)
C2—C3—C4—C5	0.0 (3)	C9—C10—C15—C14	-179.40 (17)
C3—C4—C5—C6	-0.4 (3)	C20—N4—C16—C17	-0.1 (3)
C4—C5—C6—C1	-0.1 (3)	Cu1—N4—C16—C17	175.93 (16)
C4—C5—C6—C7	-179.44 (17)	N4—C16—C17—C18	0.0 (3)
O1—C1—C6—C5	-178.95 (17)	C16—C17—C18—C19	-0.1 (3)
C2—C1—C6—C5	0.8 (3)	C17—C18—C19—C20	0.4 (3)
O1—C1—C6—C7	0.4 (3)	C16—N4—C20—C19	0.4 (3)
C2—C1—C6—C7	-179.86 (17)	Cu1—N4—C20—C19	-175.60 (15)
N2—N1—C7—C6	179.34 (16)	C18—C19—C20—N4	-0.6 (3)
Cu1—N1—C7—C6	-0.5 (3)		



Fig. 1

