

3-Hydroxy-4-nitrophenyl acetate

 Chao Liu,^{a*} Chunqing Cheng^b and Xiujie Ji^{c*}

^aSchool of Materials Science and Engineering, Hebei University of Technology, Tianjin 300130, People's Republic of China, ^bSchool of Pharmacy, Jiangxi Science and Technology, Normal University, Jiangxi 330013, People's Republic of China, and ^cTianjin Municipal Key Laboratory of Fiber Modification and Functional Fibers, Tianjin Polytechnic University, Tianjin 300160, People's Republic of China
Correspondence e-mail: liuchao_tj@yahoo.com, jxjchem@yahoo.com


Received 21 November 2008; accepted 4 December 2008

Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.028; wR factor = 0.083; data-to-parameter ratio = 11.2.

In the molecule of the title compound, $\text{C}_8\text{H}_7\text{NO}_5$, the acetate group is oriented with respect to the aromatic ring at a dihedral angle of $85.30(3)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond results in the formation of a non-planar six-membered ring, adopting an envelope conformation. In the crystal structure, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules.

Related literature

For general background to phenolic esters as intermediates in organic synthesis, see: Trollsås *et al.* (1996); Svensson *et al.* (1998); Atkinson *et al.* (2005); Hu *et al.* (2001). For a related structure, see: Ji *et al.* (2006). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_8\text{H}_7\text{NO}_5$
 $M_r = 197.15$
 Monoclinic, $P2_1/n$
 $a = 10.881(2)$ Å

$b = 5.0543(10)$ Å
 $c = 15.318(3)$ Å
 $\beta = 93.75(3)^\circ$
 $V = 840.6(3)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.13$ mm⁻¹

$T = 153(2)$ K
 $0.24 \times 0.20 \times 0.16$ mm

Data collection

Bruker SMART diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.969$, $T_{\max} = 0.979$

5058 measured reflections
 1449 independent reflections
 1232 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.083$
 $S = 1.12$
 1449 reflections

129 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.18$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{O4}$	0.82	1.91	2.605(2)	142
$\text{C5}-\text{H5}\cdots\text{O1}^i$	0.93	2.58	3.229(2)	127
$\text{C8}-\text{H8}\cdots\text{O3}^{ii}$	0.93	2.56	3.481(2)	170

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, -y + 2, -z + 2$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2582).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Atkinson, P. J., Bromidge, S. M., Duxon, M. S., Gaster, L. M., Hadley, M. S., Hammond, B., Johnson, C. N., Middlemiss, D. N., North, S. E., Price, G. W., Rami, H. K., Riley, G. J., Scott, C. M., Shaw, T. E., Starr, K. R., Stemp, G., Thewlis, K. M., Thomas, D. R., Thompson, M., Vong, A. K. K. & Watson, J. M. (2005). *Bioorg. Med. Chem. Lett.* **15**, 737–741.
- Bruker (1997). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hu, B., Ellingboe, J., Gunawan, I., Han, S., Largis, E., Li, Z., Malamas, M., Mulvey, R., Oliphant, A., Sum, F.-W., Tillett, J. & Wong, V. (2001). *Bioorg. Med. Chem. Lett.* **11**, 757–760.
- Ji, X. & Li, C. (2006). *Synthesis*, **15**, 2478–2482.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Svensson, M., Helgee, B., Skarp, K. & Andersson, G. (1998). *J. Mater. Chem.* **8**, 353–362.
- Trollsås, M., Orrenius, C., Sahlén, F., Gedde, U. W., Norin, T., Hult, A., Hermann, D., Rudquist, P., Komitov, L., Lagerwall, S. T. & Lindström, J. (1996). *J. Am. Chem. Soc.* **118**, 8542–8548.

supplementary materials

Acta Cryst. (2009). E65, o46 [doi:10.1107/S1600536808040981]

3-Hydroxy-4-nitrophenyl acetate

C. Liu, C. Cheng and X. Ji

Comment

Phenolic esters are useful intermediates in organic synthesis (Trollsås *et al.*, 1996; Svensson *et al.*, 1998; Atkinson *et al.*, 2005; Hu *et al.*, 2001). We have developed a new method for the syntheses of some phenolic esters (Ji *et al.*, 2006). The title compound is one of the products, and we report herein its crystal structure.

In the molecule of the title compound (Fig. 1) the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. The acetate group is oriented with respect to the aromatic ring at a dihedral angle of 85.30 (3)°. The intramolecular O-H...O hydrogen bond (Table 1) results in the formation of a nonplanar six-membered ring (N1/O3/O4/C6/C7/H3), adopting envelope conformation.

In the crystal structure, intermolecular C-H...O hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure.


Experimental


For the preparation of the title compound, 2-nitroresorcin acetate (239 mg, 1.0 mmol) was dissolved in chloroform (20 ml). At 273-278 K, anhydrous AlCl₃ (133.5 mg, 1 mmol) was added to this solution, the reaction was stirred at room temperature for 1 h, and then hydrochloric acid (5 ml, 10%) was added. The reaction mixture was extracted with chloroform and dried with anhydrous sodium sulfate. After concentration, the residue was separated by flash column chromatography and purified by recrystallization from chloroform (yield; 144 mg, 73%, m.p. 360 K). Spectroscopic analysis: IR (KBr, ν , cm⁻¹): 3253, 3083, 2946, 1758, 1530, 1204, 1138, 978, 847. Analysis required for C₈H₇NO₅: C 48.74; H 3.58; N 7.10%. Found: C 48.80; H 3.61; N 7.08%.

Refinement

H atoms were positioned geometrically, with O-H = 0.82 Å (for OH) and C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{O})$, where $x = 1.2$ for aromatic H and $x = 1.5$ for all other H atoms.

Figures

 Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen bond is shown as a dashed line.

 Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

3-Hydroxy-4-nitrophenyl acetate

Crystal data

$C_8H_7NO_5$	$F_{000} = 408$
$M_r = 197.15$	$D_x = 1.558 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 360 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation
$a = 10.881 (2) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 5.0543 (10) \text{ \AA}$	Cell parameters from 2650 reflections
$c = 15.318 (3) \text{ \AA}$	$\theta = 2.2\text{--}27.5^\circ$
$\beta = 93.75 (3)^\circ$	$\mu = 0.13 \text{ mm}^{-1}$
$V = 840.6 (3) \text{ \AA}^3$	$T = 153 (2) \text{ K}$
$Z = 4$	Block, colorless
	$0.24 \times 0.20 \times 0.16 \text{ mm}$

Data collection

Bruker P4 diffractometer	1449 independent reflections
Radiation source: fine-focus sealed tube	1232 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 153(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.969$, $T_{\text{max}} = 0.979$	$k = -6 \rightarrow 5$
5058 measured reflections	$l = -18 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.083$	$w = 1/[\sigma^2(F_o^2) + (0.0475P)^2 + 0.1223P]$
$S = 1.12$	where $P = (F_o^2 + 2F_c^2)/3$
1449 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
129 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.14195 (8)	0.58036 (19)	0.89666 (6)	0.0279 (3)
O2	0.31702 (7)	0.35401 (18)	0.93249 (5)	0.0244 (3)
O3	0.59843 (8)	1.04495 (18)	0.88687 (5)	0.0223 (2)
H3	0.6350	1.1192	0.8486	0.033*
O4	0.67226 (8)	1.08418 (17)	0.72904 (5)	0.0230 (2)
O5	0.63078 (8)	0.77288 (18)	0.63478 (5)	0.0247 (2)
N1	0.61633 (9)	0.8778 (2)	0.70551 (6)	0.0179 (3)
C1	0.13908 (11)	0.2225 (3)	1.00085 (8)	0.0213 (3)
H1A	0.0532	0.2620	1.0041	0.032*
H1B	0.1807	0.2425	1.0577	0.032*
H1C	0.1482	0.0438	0.9810	0.032*
C2	0.19373 (11)	0.4079 (2)	0.93821 (7)	0.0179 (3)
C3	0.38267 (10)	0.5030 (3)	0.87381 (8)	0.0195 (3)
C4	0.38272 (11)	0.4184 (3)	0.78716 (8)	0.0207 (3)
H4	0.3323	0.2804	0.7666	0.025*
C5	0.46001 (11)	0.5463 (2)	0.73292 (8)	0.0194 (3)
H5	0.4633	0.4919	0.6751	0.023*
C6	0.53337 (10)	0.7573 (2)	0.76450 (7)	0.0164 (3)
C7	0.53043 (10)	0.8455 (2)	0.85126 (7)	0.0168 (3)
C8	0.45238 (10)	0.7116 (2)	0.90598 (7)	0.0189 (3)
H8	0.4480	0.7643	0.9639	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0222 (5)	0.0277 (6)	0.0342 (5)	0.0061 (4)	0.0054 (4)	0.0129 (4)
O2	0.0180 (5)	0.0266 (5)	0.0290 (5)	0.0010 (4)	0.0041 (3)	0.0131 (4)
O3	0.0269 (5)	0.0225 (5)	0.0177 (4)	-0.0062 (4)	0.0019 (4)	-0.0023 (3)
O4	0.0233 (5)	0.0224 (5)	0.0232 (5)	-0.0057 (4)	0.0018 (3)	-0.0005 (4)
O5	0.0302 (5)	0.0282 (5)	0.0164 (4)	0.0030 (4)	0.0064 (4)	-0.0034 (4)
N1	0.0165 (5)	0.0200 (6)	0.0171 (5)	0.0032 (4)	0.0004 (4)	0.0005 (4)
C1	0.0219 (6)	0.0208 (7)	0.0216 (6)	-0.0008 (5)	0.0033 (5)	0.0030 (5)

supplementary materials

C2	0.0176 (6)	0.0169 (6)	0.0190 (6)	-0.0003 (5)	0.0003 (5)	-0.0026 (5)
C3	0.0143 (6)	0.0208 (7)	0.0235 (6)	0.0042 (5)	0.0021 (5)	0.0071 (5)
C4	0.0177 (6)	0.0182 (7)	0.0255 (7)	-0.0008 (5)	-0.0033 (5)	0.0010 (5)
C5	0.0204 (6)	0.0191 (7)	0.0180 (6)	0.0029 (5)	-0.0022 (5)	-0.0012 (5)
C6	0.0150 (6)	0.0177 (6)	0.0166 (6)	0.0033 (5)	0.0012 (4)	0.0018 (4)
C7	0.0162 (6)	0.0158 (6)	0.0182 (6)	0.0035 (5)	-0.0016 (4)	-0.0003 (5)
C8	0.0197 (6)	0.0211 (7)	0.0159 (6)	0.0043 (5)	0.0022 (5)	0.0017 (5)

Geometric parameters (Å, °)

O1—C2	1.1982 (15)	C1—H1C	0.9600
O2—C2	1.3773 (15)	C3—C8	1.3716 (18)
O2—C3	1.4032 (14)	C3—C4	1.3944 (17)
O3—C7	1.3447 (15)	C4—C5	1.3807 (17)
O3—H3	0.8200	C4—H4	0.9300
O4—N1	1.2489 (13)	C5—C6	1.3994 (17)
O5—N1	1.2255 (12)	C5—H5	0.9300
N1—C6	1.4521 (15)	C6—C7	1.4041 (16)
C1—C2	1.4923 (16)	C7—C8	1.4054 (17)
C1—H1A	0.9600	C8—H8	0.9300
C1—H1B	0.9600		
C2—O2—C3	118.26 (9)	C4—C3—O2	118.51 (11)
C7—O3—H3	109.5	C5—C4—C3	117.87 (12)
O5—N1—O4	121.88 (10)	C5—C4—H4	121.1
O5—N1—C6	119.32 (10)	C3—C4—H4	121.1
O4—N1—C6	118.79 (9)	C4—C5—C6	120.33 (11)
C2—C1—H1A	109.5	C4—C5—H5	119.8
C2—C1—H1B	109.5	C6—C5—H5	119.8
H1A—C1—H1B	109.5	C5—C6—C7	121.42 (11)
C2—C1—H1C	109.5	C5—C6—N1	117.92 (10)
H1A—C1—H1C	109.5	C7—C6—N1	120.63 (11)
H1B—C1—H1C	109.5	O3—C7—C6	125.16 (10)
O1—C2—O2	122.43 (11)	O3—C7—C8	117.20 (10)
O1—C2—C1	127.24 (11)	C6—C7—C8	117.62 (11)
O2—C2—C1	110.33 (10)	C3—C8—C7	119.85 (11)
C8—C3—C4	122.88 (11)	C3—C8—H8	120.1
C8—C3—O2	118.37 (10)	C7—C8—H8	120.1
C3—O2—C2—O1	-1.83 (17)	O5—N1—C6—C7	169.13 (10)
C3—O2—C2—C1	177.59 (10)	O4—N1—C6—C7	-9.92 (16)
C2—O2—C3—C8	98.78 (13)	C5—C6—C7—O3	179.46 (10)
C2—O2—C3—C4	-86.75 (14)	N1—C6—C7—O3	1.57 (18)
C8—C3—C4—C5	2.32 (18)	C5—C6—C7—C8	0.95 (17)
O2—C3—C4—C5	-171.88 (10)	N1—C6—C7—C8	-176.94 (10)
C3—C4—C5—C6	-1.34 (18)	C4—C3—C8—C7	-1.64 (18)
C4—C5—C6—C7	-0.25 (18)	O2—C3—C8—C7	172.57 (10)
C4—C5—C6—N1	177.70 (10)	O3—C7—C8—C3	-178.67 (10)
O5—N1—C6—C5	-8.83 (15)	C6—C7—C8—C3	-0.03 (17)
O4—N1—C6—C5	172.12 (9)		

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 \cdots O4	0.82	1.91	2.605 (2)	142
C5—H5 \cdots O1 ⁱ	0.93	2.58	3.229 (2)	127
C8—H8 \cdots O3 ⁱⁱ	0.93	2.56	3.481 (2)	170

Symmetry codes: (i) $-x+1/2, y-1/2, -z+3/2$; (ii) $-x+1, -y+2, -z+2$.

Fig. 1




Fig. 2

