

Ethyl 4-anilino-3-nitrobenzoate

Hao-Yuan Li,^a Bo-Nian Liu,^a Shi-Gui Tang^b and Cheng Guo^{a*}

^aCollege of Science, Nanjing University of Technology, Xinmofan Road No. 5, Nanjing 210009, People's Republic of China, and ^bCollege of Life Sciences and Pharmaceutical Engineering, Nanjing University of Technology, Nanjing 210009, People's Republic of China
Correspondence e-mail: guocheng@njut.edu.cn


Received 4 December 2008; accepted 7 December 2008

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.071; wR factor = 0.199; data-to-parameter ratio = 13.2.

In the title compound, $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_4$, the aromatic rings are oriented at a dihedral angle of $78.33(3)^\circ$. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond results in a non-planar six-membered ring with a flattened-boat conformation. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules. $\pi-\pi$ contacts between the phenyl rings and both the phenyl and benzene rings, [centroid–centroid distances = $3.841(3)$ and $3.961(3)\text{ \AA}$] may further stabilize the structure.

Related literature

For bond-length data, see: Allen *et al.* (1987). For ring puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data



$M_r = 286.28$

Monoclinic, $P2_1/n$
 $a = 10.683(2)\text{ \AA}$
 $b = 9.905(2)\text{ \AA}$
 $c = 13.698(3)\text{ \AA}$
 $\beta = 105.05(3)^\circ$
 $V = 1399.7(5)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 294(2)\text{ K}$
 $0.30 \times 0.20 \times 0.20\text{ mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.971$, $T_{\max} = 0.980$
2647 measured reflections

2508 independent reflections
1519 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$
3 standard reflections
frequency: 120 min
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.199$
 $S = 1.00$
2508 reflections

190 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A \cdots O3	0.86	1.98	2.623 (5)	131
N2—H2A \cdots O2 ⁱ	0.86	2.31	2.978 (4)	134

Symmetry code: (i) $x - \frac{1}{2}, -y - \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors thank Dr Shan Liu, Nanjing University of Technology, for useful discussions and the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2595).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, S1–19.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft. The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o91 [doi:10.1107/S1600536808041329]

Ethyl 4-anilino-3-nitrobenzoate

H.-Y. Li, B.-N. Liu, S.-G. Tang and C. Guo

Comment

Some derivatives of benzoic acid are important chemical materials. We report herein the crystal structure of the title compound.

In the molecule of the title compound (Fig 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (C4-C9) and B (C10-C15) are, of course, planar, and they are oriented at a dihedral angle of 78.33 (3) $^{\circ}$. The intramolecular N-H \cdots O hydrogen bond (Table 1) results in a nonplanar six-membered ring C (O3/N1/N2/C6/C7/H2A), having total puckering amplitude, Q_T , of 0.131 (2) Å, flattened-boat conformation [$\phi = 140.37$ (3) $^{\circ}$ and $\theta = 75.09$ (4) $^{\circ}$] (Cremer & Pople, 1975).

In the crystal structure, intermolecular N-H \cdots O hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. The π - π contacts between the phenyl rings and the phenyl and the benzene rings, Cg1—Cg1ⁱ and Cg1—Cg2ⁱⁱ [symmetry codes: (i) 1 - x, 1 - y, 1 - z; (ii) x - 1/2, 1/2 - y, z - 1/2, where Cg1 and Cg2 are centroids of the rings A (C4-C6) and B (C10-C15), respectively] may further stabilize the structure, with centroid-centroid distances of 3.841 (3) Å and 3.961 (3) Å.

Experimental

For the preparation of the title compound, ethyl 4-chloro-3-nitrobenzoate (5.0 g, 0.022 mol) was heated in fresh distilled aniline (10 ml) for 18 h at 393 K, and then ethanol (50 ml) was added, at room temperature. The yellow precipitate was sucked, washed with cold ethanol (2 X 20 ml), and then dried (yield; 4.7 g, 75%). Crystals suitable for X-ray analysis were obtained by slow evaporation of an ethanol solution.

Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH) and C-H = 0.93, 0.97 and 0.96 Å for aromatic, methylene and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Hydrogen bond is shown as dashed line.

supplementary materials




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

Ethyl 4-anilino-3-nitrobenzoate

Crystal data

C ₁₅ H ₁₄ N ₂ O ₄	$F_{000} = 600$
$M_r = 286.28$	$D_x = 1.358 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 10.683 (2) \text{ \AA}$	Cell parameters from 25 reflections
$b = 9.905 (2) \text{ \AA}$	$\theta = 10\text{--}13^\circ$
$c = 13.698 (3) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 105.05 (3)^\circ$	$T = 294 (2) \text{ K}$
$V = 1399.7 (5) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.30 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Enraf-Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.051$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.2^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.2^\circ$
$T = 294(2) \text{ K}$	$h = -12 \rightarrow 12$
$\omega/2\theta$ scans	$k = 0 \rightarrow 11$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = 0 \rightarrow 16$
$T_{\text{min}} = 0.971$, $T_{\text{max}} = 0.980$	3 standard reflections
2647 measured reflections	every 120 min
2508 independent reflections	intensity decay: none
1519 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.071$	H-atom parameters constrained
$wR(F^2) = 0.199$	$w = 1/[\sigma^2(F_o^2) + (0.050P)^2 + 3.4P]$

$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
2508 reflections	$(\Delta/\sigma)_{\max} < 0.001$
190 parameters	$\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections (Sheldrick, 2008). The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.0229 (3)	0.1219 (3)	0.2353 (2)	0.0602 (9)
O2	0.1935 (2)	-0.0076 (3)	0.2291 (2)	0.0605 (8)
O3	-0.0506 (3)	-0.4222 (3)	-0.1083 (3)	0.0705 (10)
O4	0.1353 (3)	-0.3433 (4)	-0.0297 (3)	0.0852 (12)
N1	0.0155 (3)	-0.3404 (4)	-0.0480 (3)	0.0560 (9)
N2	-0.2640 (3)	-0.2968 (4)	-0.1003 (2)	0.0523 (9)
H2A	-0.2321	-0.3625	-0.1271	0.063*
C1	0.0290 (5)	0.2946 (6)	0.3560 (4)	0.0943 (19)
H1A	0.0836	0.3471	0.4092	0.141*
H1B	-0.0328	0.2455	0.3820	0.141*
H1C	-0.0159	0.3535	0.3027	0.141*
C2	0.1084 (4)	0.1999 (5)	0.3166 (4)	0.0683 (14)
H2B	0.1541	0.1399	0.3700	0.082*
H2C	0.1718	0.2486	0.2909	0.082*
C3	0.0774 (4)	0.0183 (4)	0.2003 (3)	0.0489 (10)
C4	-0.0131 (3)	-0.0606 (4)	0.1202 (3)	0.0464 (10)
C5	0.0350 (3)	-0.1607 (4)	0.0723 (3)	0.0452 (10)
H5A	0.1238	-0.1764	0.0897	0.054*
C6	-0.0450 (3)	-0.2399 (4)	-0.0018 (3)	0.0434 (9)
C7	-0.1821 (3)	-0.2191 (4)	-0.0295 (3)	0.0438 (9)
C8	-0.2275 (3)	-0.1155 (4)	0.0203 (3)	0.0486 (10)
H8A	-0.3161	-0.0984	0.0034	0.058*
C9	-0.1479 (3)	-0.0370 (4)	0.0934 (3)	0.0465 (10)
H9A	-0.1827	0.0313	0.1249	0.056*
C10	-0.4030 (3)	-0.2732 (5)	-0.1322 (3)	0.0526 (11)
C11	-0.4831 (4)	-0.3588 (6)	-0.1004 (4)	0.0754 (15)

supplementary materials

H11A	-0.4487	-0.4284	-0.0559	0.090*
C12	-0.6166 (5)	-0.3428 (7)	-0.1342 (5)	0.0907 (19)
H12A	-0.6722	-0.4012	-0.1127	0.109*
C13	-0.6650 (4)	-0.2397 (7)	-0.1995 (5)	0.096 (2)
H13A	-0.7543	-0.2280	-0.2219	0.115*
C14	-0.5848 (4)	-0.1536 (6)	-0.2324 (4)	0.0854 (18)
H14A	-0.6199	-0.0841	-0.2768	0.103*
C15	-0.4501 (4)	-0.1691 (5)	-0.1999 (4)	0.0654 (13)
H15B	-0.3944	-0.1122	-0.2226	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0423 (16)	0.0602 (19)	0.0576 (18)	-0.0073 (14)	-0.0236 (13)	-0.0105 (15)
O2	0.0306 (14)	0.075 (2)	0.0604 (18)	-0.0033 (14)	-0.0167 (12)	0.0023 (16)
O3	0.0482 (17)	0.067 (2)	0.080 (2)	0.0093 (16)	-0.0131 (16)	-0.0220 (18)
O4	0.0283 (15)	0.115 (3)	0.104 (3)	0.0123 (17)	0.0021 (16)	-0.016 (2)
N1	0.0367 (18)	0.069 (2)	0.056 (2)	0.0118 (18)	-0.0003 (16)	-0.0016 (19)
N2	0.0263 (16)	0.066 (2)	0.056 (2)	-0.0001 (15)	-0.0046 (14)	-0.0167 (18)
C1	0.090 (4)	0.092 (4)	0.082 (4)	-0.013 (3)	-0.013 (3)	-0.026 (3)
C2	0.057 (3)	0.068 (3)	0.061 (3)	-0.013 (2)	-0.018 (2)	-0.017 (3)
C3	0.040 (2)	0.053 (3)	0.042 (2)	-0.0083 (19)	-0.0115 (17)	0.009 (2)
C4	0.0278 (18)	0.052 (2)	0.045 (2)	-0.0041 (17)	-0.0165 (16)	0.0055 (19)
C5	0.0262 (18)	0.052 (2)	0.046 (2)	-0.0026 (17)	-0.0117 (16)	0.0101 (19)
C6	0.0299 (18)	0.053 (2)	0.041 (2)	0.0049 (17)	-0.0023 (16)	0.0030 (18)
C7	0.0240 (17)	0.049 (2)	0.048 (2)	-0.0009 (17)	-0.0090 (15)	-0.0001 (19)
C8	0.0233 (17)	0.060 (3)	0.053 (2)	0.0000 (17)	-0.0071 (16)	-0.001 (2)
C9	0.0327 (19)	0.052 (2)	0.047 (2)	-0.0006 (17)	-0.0021 (16)	-0.0051 (19)
C10	0.0226 (18)	0.068 (3)	0.057 (3)	-0.0039 (19)	-0.0082 (17)	-0.023 (2)
C11	0.047 (3)	0.107 (4)	0.072 (3)	-0.015 (3)	0.014 (2)	-0.016 (3)
C12	0.048 (3)	0.118 (5)	0.114 (5)	-0.026 (3)	0.036 (3)	-0.033 (4)
C13	0.024 (2)	0.127 (6)	0.122 (5)	-0.009 (3)	-0.006 (3)	-0.064 (5)
C14	0.039 (3)	0.098 (4)	0.097 (4)	0.016 (3)	-0.022 (3)	-0.030 (3)
C15	0.033 (2)	0.059 (3)	0.089 (3)	0.002 (2)	-0.013 (2)	-0.008 (3)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.463 (5)	C5—C6	1.387 (5)
O1—C3	1.329 (5)	C5—H5A	0.9300
O2—C3	1.227 (4)	C6—C7	1.429 (5)
N1—O3	1.239 (4)	C7—C8	1.387 (5)
N1—O4	1.240 (4)	C8—C9	1.374 (5)
N1—C6	1.422 (5)	C8—H8A	0.9300
N2—C7	1.363 (5)	C9—H9A	0.9300
N2—C10	1.454 (4)	C10—C11	1.354 (6)
N2—H2A	0.8600	C10—C15	1.390 (6)
C1—C2	1.458 (7)	C11—C12	1.390 (7)
C1—H1A	0.9600	C11—H11A	0.9300
C1—H1B	0.9600	C12—C13	1.367 (9)

C1—H1C	0.9600	C12—H12A	0.9300
C2—H2B	0.9700	C13—C14	1.366 (8)
C2—H2C	0.9700	C13—H13A	0.9300
C3—C4	1.483 (5)	C14—C15	1.400 (6)
C4—C5	1.360 (6)	C14—H14A	0.9300
C4—C9	1.410 (5)	C15—H15B	0.9300
C3—O1—C2	115.9 (3)	C5—C6—N1	117.2 (3)
O3—N1—O4	120.0 (4)	C5—C6—C7	120.5 (4)
O3—N1—C6	120.6 (3)	N2—C7—C8	121.7 (3)
O4—N1—C6	119.4 (4)	N2—C7—C6	122.1 (4)
C7—N2—C10	122.7 (3)	C8—C7—C6	116.1 (3)
C7—N2—H2A	118.7	C7—C8—H8A	118.4
C10—N2—H2A	118.7	C9—C8—C7	123.2 (3)
C2—C1—H1A	109.5	C9—C8—H8A	118.4
C2—C1—H1B	109.5	C4—C9—H9A	120.3
C2—C1—H1C	109.5	C8—C9—C4	119.5 (4)
H1A—C1—H1B	109.5	C8—C9—H9A	120.3
H1A—C1—H1C	109.5	C11—C10—C15	122.0 (4)
H1B—C1—H1C	109.5	C11—C10—N2	118.9 (4)
O1—C2—H2B	110.0	C15—C10—N2	119.0 (4)
O1—C2—H2C	110.0	C10—C11—C12	120.1 (6)
C1—C2—O1	108.3 (4)	C10—C11—H11A	120.0
C1—C2—H2B	110.0	C12—C11—H11A	120.0
C1—C2—H2C	110.0	C11—C12—H12A	120.5
H2B—C2—H2C	108.4	C13—C12—C11	118.9 (5)
O1—C3—C4	114.3 (3)	C13—C12—H12A	120.5
O2—C3—O1	123.1 (4)	C12—C13—H13A	119.4
O2—C3—C4	122.6 (4)	C14—C13—C12	121.3 (5)
C5—C4—C9	118.9 (3)	C14—C13—H13A	119.4
C5—C4—C3	119.0 (3)	C13—C14—C15	120.5 (6)
C9—C4—C3	122.1 (4)	C13—C14—H14A	119.8
C4—C5—C6	121.8 (3)	C15—C14—H14A	119.8
C4—C5—H5A	119.1	C10—C15—C14	117.2 (5)
C6—C5—H5A	119.1	C10—C15—H15B	121.4
N1—C6—C7	122.4 (3)	C14—C15—H15B	121.4
C3—O1—C2—C1	-171.2 (4)	N1—C6—C7—N2	1.8 (6)
C2—O1—C3—O2	-4.1 (6)	C5—C6—C7—C8	1.1 (6)
C2—O1—C3—C4	178.3 (4)	N1—C6—C7—C8	-178.6 (4)
O2—C3—C4—C5	-3.9 (6)	N2—C7—C8—C9	178.7 (4)
O1—C3—C4—C5	173.7 (4)	C6—C7—C8—C9	-0.9 (6)
O2—C3—C4—C9	174.8 (4)	C7—C8—C9—C4	0.1 (6)
O1—C3—C4—C9	-7.6 (5)	C5—C4—C9—C8	0.4 (6)
C9—C4—C5—C6	-0.2 (6)	C3—C4—C9—C8	-178.4 (4)
C3—C4—C5—C6	178.6 (4)	C7—N2—C10—C11	-106.2 (5)
C4—C5—C6—N1	179.1 (4)	C7—N2—C10—C15	78.1 (5)
C4—C5—C6—C7	-0.6 (6)	C15—C10—C11—C12	-1.0 (7)
O3—N1—C6—C5	173.7 (4)	N2—C10—C11—C12	-176.6 (4)
O4—N1—C6—C5	-8.2 (6)	C10—C11—C12—C13	-0.1 (8)

supplementary materials


O3—N1—C6—C7	−6.6 (6)	C11—C12—C13—C14	0.5 (9)
O4—N1—C6—C7	171.5 (4)	C12—C13—C14—C15	0.1 (8)
C10—N2—C7—C8	3.1 (6)	C11—C10—C15—C14	1.6 (7)
C10—N2—C7—C6	−177.4 (4)	N2—C10—C15—C14	177.2 (4)
C5—C6—C7—N2	−178.5 (4)	C13—C14—C15—C10	−1.1 (7)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2A···O3	0.86	1.98	2.623 (5)	131
N2—H2A···O2 ⁱ	0.86	2.31	2.978 (4)	134

Symmetry codes: (i) $x-1/2, -y-1/2, z-1/2$.

Fig. 1



supplementary materials

Fig. 2

