

4-Chloro-2-((1*R*)-1-[(*R*)-(2-chlorophenyl)(cyclopentyl)methyl]amino)-propylphenol

Guang-You Zhang,^a Ting Yang,^{a*} Bao-Wang Xu,^b Di-Juan Chen^a and Wan-Hui Wang^c

^aSchool of Chemistry, Jinan University, 250022, People's Republic of China, ^bQilu Pharmaceutical Co Ltd, Shandong Province, 250100, People's Republic of China, and ^cGraduate School of Science and Engineering, Saitama University, Sakura, Saitama 338-8570, Japan

Correspondence e-mail: yangting365@126.com


Received 14 November 2008; accepted 10 December 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.093; data-to-parameter ratio = 15.8.

In the title compound, $C_{21}H_{25}Cl_2NO$, the dihedral angle between the two benzene rings is $33.18(11)^\circ$. The five-membered ring adopts an envelope conformation. There is an intramolecular $O-\text{H}\cdots\text{N}$ hydrogen bond. In the crystal, molecules are linked by weak $N-\text{H}\cdots\text{Cl}$ hydrogen bonds, forming a helical chain along the c axis.

Related literature

For related literature on aminophenols, see: Cimarelli *et al.* (2002); Joshi & Malhotra (2003); Li *et al.* (2004); Puigjaner *et al.* (1999); Watts *et al.* (2005). For the synthesis, see: Yang *et al.* (2005).



Experimental

Crystal data

$C_{21}H_{25}Cl_2NO$	$V = 1972.1(5)\text{ \AA}^3$
$M_r = 378.32$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 10.9802(17)\text{ \AA}$	$\mu = 0.34\text{ mm}^{-1}$
$b = 11.5607(18)\text{ \AA}$	$T = 298(2)\text{ K}$
$c = 15.536(2)\text{ \AA}$	$0.49 \times 0.45 \times 0.38\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	10332 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3647 independent reflections
$R_{\text{int}} = 0.025$	3266 reflections with $I > 2\sigma(I)$
$T_{\text{min}} = 0.852$, $T_{\text{max}} = 0.882$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.093$	$\Delta\rho_{\text{max}} = 0.17\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.20\text{ e \AA}^{-3}$
3647 reflections	Absolute structure: Flack (1983), 1556 Friedel pairs
231 parameters	Flack parameter: 0.06 (6)
1 restraint	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots Cl2 ⁱ	0.848 (19)	2.913 (13)	3.7023 (18)	156 (2)
O1—H1A \cdots N1	0.82	1.93	2.642 (2)	144

Symmetry code: (i) $-x + \frac{1}{2}, -y + 2, z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful to the Natural Science Foundation of Shandong Province, China (grant No. G0231) and the Foundation of the Education Ministry of China for Returned Students (grant No. G0220) for financial support. The X-ray data were collected at Shandong Normal University, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2364).

References

- Bruker (1999). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cimarelli, C., Palmieri, G. & Volpini, E. (2002). *Tetrahedron Asymmetry*, **13**, 2011–2018.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Joshi, S. N. & Malhotra, S. V. (2003). *Tetrahedron Asymmetry*, **14**, 1763–1766.
- Li, Y., He, B., Qin, B., Feng, X. & Zhang, G. (2004). *J. Org. Chem.* **69**, 7910–7913.
- Puigjaner, C., Vidal-Ferran, A., Moyano, A., Pericas, M. A., Rieras, M. A. & Riera, A. (1999). *J. Org. Chem.* **64**, 7902–7911.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Watts, C. C., Thoniyyot, P., Hirayama, L. C., Romano, T. & Singaram, B. (2005). *Tetrahedron Asymmetry*, **16**, 1829–1835.
- Yang, X.-F., Zhang, G.-Y., Zhang, Y., Zhao, J.-Y. & Wang, X.-B. (2005). *Acta Cryst.* **C61**, o262–o264.

supplementary materials

Acta Cryst. (2009). E65, o123 [doi:10.1107/S1600536808042025]

4-Chloro-2-((1*R*)-1-<{[(R)-(2-chlorophenyl)(cyclopentyl)methyl]amino}propyl)phenol

G.-Y. Zhang, T. Yang, B.-W. Xu, D.-J. Chen and W.-H. Wang

Comment

The synthesis of enantiopure aminophenols that have different functionalities is an important subject of research because compounds of this class are widespread in natural products, show pharmacological activity and have recently found application in asymmetric synthesis as chiral bases, auxiliaries and ligands (Cimarelli *et al.*, 2002). Chiral aminophenols which are similar to amino alcohols have attracted wide attention for the reason that they can be used in catalytic asymmetric reactions (Puigjaner *et al.*, 1999; Li *et al.*, 2004; Watts *et al.*, 2005), which is one of the most active areas of research in organic chemistry (Joshi & Malhotra, 2003). The synthesis of new aminoalkylphenols is therefore of interest because of potential as asymmetric catalysts.

As part of our continuing studies of chiral aminophenols, we now report the crystal structure of the title compound, (I), which was initially prepared to test its asymmetric catalytic activity. These compounds were prepared by conventional condensation of (*R*)-1-(2-chlorophenyl)-1-cyclopentylmethanamine with 1-(5-chloro-2-hydroxyphenyl)ethanone, followed by reduction using sodium borohydride in a tetrahydrofuran-ethanol (1:1 v/v) mixture. An X-ray study of the title compound, (I), was carried out and the results are presented here. The molecular structure of (I) is shown in Fig. 1.

The molecule has two chiral centres (C7/C10), which have configuration *R*, *R*, as shown in Fig. 1. In the molecules of (I), the five-membered rings adopts an envelope conformation. The dihedral angle between the benzene rings is 33.18 (11)°. There is an intramolecular O1—H1A···N1 hydrogen bond (Table 1). Phenol atom O1 acts as a hydrogen bond donor to atom N1, with O1···N1 = 2.647 (2) Å, which indicates a comparatively strong intramolecular hydrogen bond (Table 1); this distance is significantly shorter than the sum (3.07 Å) of the van der Waals radii for N and O atoms. The molecules are linked via N1—H1···Cl2 hydrogen bonds. An interesting feature of the structure is that the N1—H1···Cl2 hydrogen-bond gives rise to a spiral chain of molecules along the *c* direction. There are no π-π stacking interactions are present in the structure of (I).

Experimental

The title compound were prepared according to the procedure of Yang *et al.* (2005). (*R*)-1-(2-chlorophenyl)-1-cyclopentylmethanamine (0.9 mmol) and 1-(5-chloro-2-hydroxyphenyl)propan-1-one (0.9 mmol) were dissolved in methanol (10 ml) and reacted at room temperature for 48 h. After removal of the solvent, NaBH₄ (4.5 mmol) was added to the solution in THF/ethanol (1:1 v/v, 20 ml) and stirred at 273 K until the solution became colourless. The solvent was then removed under reduced pressure. Water (10 ml) was added to the residue and 1 N HCl was added dropwise until hydrogen production ceased. The mixture was neutralized with aqueous Na₂CO₃, then extracted with CHCl₃, and the organic layer was dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure. Further purification was carried out by thin-layer silica-gel chromatography (chloroform) to give a colourless solid (yield 80.5%). Crystals of (I) were grown from a n-hexane solution.

supplementary materials

Refinement

The N-bound H atom was located in a Fourier difference map and was refined with a distance restraint of N—H = 0.86 (1) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The O-bound and C-bound H atoms were positioned geometrically (O—H = 0.82 Å and C—H = 0.93–0.98 Å) and were treated as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O, methyl C})$.

Figures




Fig. 1. The asymmetric unit of (I), showing the atom-labelling schemes. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as spheres of arbitrary radii.




Fig. 2. A packing diagram of (I), view down the *b* axis, showing the formation of helical chains through O1—H1A…N1 and N1—H1…Cl2 hydrogen bonds (dashed lines). H atoms not involved in hydrogen bonds have been omitted.

4-Chloro-2-((1*R*)-1-{[(*R*)-(2-chlorophenyl)(cyclopentyl)methyl]amino}propyl)phenol

Crystal data

C ₂₁ H ₂₅ Cl ₂ NO	$F_{000} = 800$
$M_r = 378.32$	$D_x = 1.274 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 10.9802 (17) \text{ \AA}$	Cell parameters from 4568 reflections
$b = 11.5607 (18) \text{ \AA}$	$\theta = 2.2\text{--}25.5^\circ$
$c = 15.536 (2) \text{ \AA}$	$\mu = 0.34 \text{ mm}^{-1}$
$V = 1972.1 (5) \text{ \AA}^3$	$T = 298 (2) \text{ K}$
$Z = 4$	Block, colourless
	$0.49 \times 0.45 \times 0.38 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	3647 independent reflections
Radiation source: fine-focus sealed tube	3266 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan	$h = -12\text{--}13$

(SADABS; Sheldrick, 1996)

$T_{\min} = 0.852$, $T_{\max} = 0.882$

10332 measured reflections

$k = -13 \rightarrow 13$

$l = -18 \rightarrow 15$

Refinement

Refinement on F^2

Hydrogen site location: inferred from neighbouring sites

Least-squares matrix: full

H atoms treated by a mixture of independent and constrained refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$

$$w = 1/[\sigma^2(F_o^2) + (0.0504P)^2 + 0.2043P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$wR(F^2) = 0.093$

$$(\Delta/\sigma)_{\max} = 0.001$$

$S = 1.03$

$$\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$$

3647 reflections

$$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$$

231 parameters

Extinction correction: none

1 restraint

Absolute structure: Flack (1983), 1556 Friedel pairs

Primary atom site location: structure-invariant direct methods

Flack parameter: 0.06 (6)

Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4025 (2)	0.9204 (2)	0.46157 (14)	0.0567 (6)
C2	0.3358 (3)	0.8241 (2)	0.43935 (15)	0.0695 (7)
H2	0.3513	0.7853	0.3881	0.083*
C3	0.2468 (3)	0.7864 (2)	0.49342 (16)	0.0687 (7)
H3	0.2013	0.7216	0.4786	0.082*
C4	0.2227 (2)	0.84331 (19)	0.57058 (15)	0.0576 (6)
C5	0.29329 (17)	0.93830 (17)	0.59471 (12)	0.0442 (5)
C6	0.38307 (18)	0.97536 (19)	0.53893 (13)	0.0476 (5)
H6	0.4312	1.0384	0.5539	0.057*
C7	0.26917 (18)	1.00349 (17)	0.67807 (13)	0.0453 (5)
H7	0.3437	1.0439	0.6955	0.054*
C8	0.1659 (2)	1.0917 (2)	0.66944 (15)	0.0621 (6)

supplementary materials

H8A	0.0917	1.0509	0.6542	0.075*
H8B	0.1528	1.1279	0.7250	0.075*
C9	0.1884 (3)	1.1847 (2)	0.60364 (17)	0.0757 (7)
H9A	0.2633	1.2238	0.6167	0.113*
H9B	0.1225	1.2393	0.6048	0.113*
H9C	0.1937	1.1506	0.5475	0.113*
C10	0.33131 (17)	0.84631 (17)	0.77983 (12)	0.0420 (4)
H10	0.3589	0.7969	0.7324	0.050*
C11	0.27940 (18)	0.76780 (18)	0.84923 (13)	0.0479 (5)
H11	0.2531	0.8162	0.8976	0.057*
C12	0.1718 (2)	0.6922 (2)	0.82169 (16)	0.0626 (6)
H12A	0.0968	0.7366	0.8208	0.075*
H12B	0.1855	0.6594	0.7650	0.075*
C13	0.1665 (3)	0.5979 (3)	0.8899 (2)	0.0886 (9)
H13A	0.1118	0.6201	0.9360	0.106*
H13B	0.1376	0.5261	0.8649	0.106*
C14	0.2942 (2)	0.5833 (2)	0.92374 (19)	0.0737 (7)
H14A	0.3261	0.5080	0.9081	0.088*
H14B	0.2951	0.5901	0.9860	0.088*
C15	0.3705 (2)	0.67875 (19)	0.88315 (15)	0.0563 (5)
H15A	0.4197	0.6482	0.8365	0.068*
H15B	0.4240	0.7135	0.9256	0.068*
C16	0.44189 (17)	0.91271 (16)	0.81295 (12)	0.0405 (4)
C17	0.55907 (18)	0.89656 (17)	0.78285 (13)	0.0466 (5)
C18	0.6565 (2)	0.9581 (2)	0.81494 (16)	0.0596 (6)
H18	0.7342	0.9451	0.7932	0.071*
C19	0.6393 (2)	1.0384 (2)	0.87866 (16)	0.0639 (6)
H19	0.7048	1.0806	0.8999	0.077*
C20	0.5237 (2)	1.0559 (2)	0.91091 (15)	0.0608 (6)
H20	0.5112	1.1094	0.9547	0.073*
C21	0.4275 (2)	0.99458 (18)	0.87846 (14)	0.0522 (5)
H21	0.3501	1.0078	0.9007	0.063*
Cl1	0.58898 (6)	0.79389 (6)	0.70334 (4)	0.0724 (2)
Cl2	0.51106 (6)	0.97523 (8)	0.39044 (4)	0.0820 (2)
N1	0.23430 (14)	0.92140 (16)	0.74620 (11)	0.0469 (4)
H1	0.1982 (19)	0.9534 (18)	0.7882 (11)	0.056*
O1	0.13051 (17)	0.80387 (17)	0.62012 (11)	0.0773 (5)
H1A	0.1323	0.8364	0.6670	0.116*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0571 (12)	0.0642 (14)	0.0489 (12)	0.0175 (12)	-0.0106 (10)	0.0005 (10)
C2	0.100 (2)	0.0584 (14)	0.0502 (13)	0.0194 (14)	-0.0232 (14)	-0.0075 (11)
C3	0.102 (2)	0.0470 (12)	0.0576 (14)	-0.0064 (13)	-0.0373 (14)	0.0001 (12)
C4	0.0675 (14)	0.0481 (11)	0.0570 (13)	-0.0112 (11)	-0.0277 (11)	0.0128 (10)
C5	0.0439 (10)	0.0445 (11)	0.0442 (10)	0.0005 (9)	-0.0147 (8)	0.0040 (8)
C6	0.0458 (11)	0.0477 (11)	0.0491 (11)	0.0033 (9)	-0.0125 (9)	-0.0017 (9)

C7	0.0441 (11)	0.0458 (11)	0.0459 (10)	-0.0009 (8)	-0.0084 (8)	0.0015 (9)
C8	0.0641 (14)	0.0609 (13)	0.0614 (13)	0.0163 (12)	0.0008 (11)	0.0083 (11)
C9	0.100 (2)	0.0566 (14)	0.0705 (15)	0.0221 (15)	0.0023 (15)	0.0104 (12)
C10	0.0412 (10)	0.0434 (10)	0.0414 (10)	0.0026 (8)	-0.0012 (8)	-0.0017 (8)
C11	0.0450 (11)	0.0529 (12)	0.0457 (11)	0.0025 (10)	-0.0024 (8)	0.0038 (9)
C12	0.0482 (12)	0.0691 (14)	0.0706 (15)	-0.0085 (11)	-0.0056 (10)	0.0153 (13)
C13	0.0688 (16)	0.090 (2)	0.107 (2)	-0.0184 (15)	-0.0105 (16)	0.0458 (18)
C14	0.0747 (16)	0.0628 (14)	0.0837 (17)	-0.0079 (14)	-0.0110 (13)	0.0242 (14)
C15	0.0546 (12)	0.0533 (12)	0.0610 (13)	0.0016 (10)	-0.0084 (10)	0.0108 (10)
C16	0.0437 (10)	0.0379 (9)	0.0401 (10)	0.0049 (8)	-0.0029 (8)	0.0018 (8)
C17	0.0470 (11)	0.0439 (10)	0.0488 (11)	0.0027 (9)	-0.0002 (9)	0.0001 (9)
C18	0.0447 (11)	0.0578 (13)	0.0763 (15)	0.0009 (10)	-0.0047 (11)	0.0011 (12)
C19	0.0616 (14)	0.0532 (13)	0.0768 (16)	-0.0082 (11)	-0.0235 (12)	-0.0001 (12)
C20	0.0738 (16)	0.0472 (12)	0.0615 (13)	0.0045 (12)	-0.0138 (12)	-0.0097 (11)
C21	0.0520 (12)	0.0485 (12)	0.0562 (12)	0.0073 (10)	-0.0037 (10)	-0.0068 (9)
Cl1	0.0588 (3)	0.0828 (4)	0.0757 (4)	0.0041 (3)	0.0154 (3)	-0.0274 (3)
Cl2	0.0672 (4)	0.1215 (6)	0.0573 (3)	0.0189 (4)	0.0075 (3)	-0.0007 (4)
N1	0.0392 (9)	0.0552 (10)	0.0463 (9)	0.0065 (8)	-0.0023 (7)	0.0070 (8)
O1	0.0815 (12)	0.0804 (12)	0.0701 (11)	-0.0394 (10)	-0.0257 (10)	0.0165 (10)

Geometric parameters (Å, °)

C1—C6	1.376 (3)	C11—C12	1.531 (3)
C1—C2	1.377 (4)	C11—H11	0.9800
C1—Cl2	1.745 (3)	C12—C13	1.521 (3)
C2—C3	1.360 (4)	C12—H12A	0.9700
C2—H2	0.9300	C12—H12B	0.9700
C3—C4	1.393 (4)	C13—C14	1.507 (4)
C3—H3	0.9300	C13—H13A	0.9700
C4—O1	1.351 (3)	C13—H13B	0.9700
C4—C5	1.395 (3)	C14—C15	1.522 (3)
C5—C6	1.381 (3)	C14—H14A	0.9700
C5—C7	1.522 (3)	C14—H14B	0.9700
C6—H6	0.9300	C15—H15A	0.9700
C7—N1	1.472 (3)	C15—H15B	0.9700
C7—C8	1.531 (3)	C16—C17	1.382 (3)
C7—H7	0.9800	C16—C21	1.399 (3)
C8—C9	1.504 (3)	C17—C18	1.378 (3)
C8—H8A	0.9700	C17—Cl1	1.744 (2)
C8—H8B	0.9700	C18—C19	1.370 (3)
C9—H9A	0.9600	C18—H18	0.9300
C9—H9B	0.9600	C19—C20	1.379 (3)
C9—H9C	0.9600	C19—H19	0.9300
C10—N1	1.470 (2)	C20—C21	1.369 (3)
C10—C11	1.520 (3)	C20—H20	0.9300
C10—C16	1.526 (3)	C21—H21	0.9300
C10—H10	0.9800	N1—H1	0.848 (19)
C11—C15	1.529 (3)	O1—H1A	0.8200
C6—C1—C2	120.7 (2)	C12—C11—H11	108.2

supplementary materials

C6—C1—Cl2	119.43 (19)	C13—C12—C11	104.11 (19)
C2—C1—Cl2	119.89 (19)	C13—C12—H12A	110.9
C3—C2—C1	119.1 (2)	C11—C12—H12A	110.9
C3—C2—H2	120.5	C13—C12—H12B	110.9
C1—C2—H2	120.5	C11—C12—H12B	110.9
C2—C3—C4	121.1 (2)	H12A—C12—H12B	109.0
C2—C3—H3	119.4	C14—C13—C12	106.7 (2)
C4—C3—H3	119.4	C14—C13—H13A	110.4
O1—C4—C3	118.2 (2)	C12—C13—H13A	110.4
O1—C4—C5	121.9 (2)	C14—C13—H13B	110.4
C3—C4—C5	119.8 (2)	C12—C13—H13B	110.4
C6—C5—C4	118.2 (2)	H13A—C13—H13B	108.6
C6—C5—C7	120.31 (17)	C13—C14—C15	106.6 (2)
C4—C5—C7	121.45 (19)	C13—C14—H14A	110.4
C1—C6—C5	121.0 (2)	C15—C14—H14A	110.4
C1—C6—H6	119.5	C13—C14—H14B	110.4
C5—C6—H6	119.5	C15—C14—H14B	110.4
N1—C7—C5	109.75 (16)	H14A—C14—H14B	108.6
N1—C7—C8	107.43 (17)	C14—C15—C11	105.70 (18)
C5—C7—C8	112.60 (16)	C14—C15—H15A	110.6
N1—C7—H7	109.0	C11—C15—H15A	110.6
C5—C7—H7	109.0	C14—C15—H15B	110.6
C8—C7—H7	109.0	C11—C15—H15B	110.6
C9—C8—C7	114.5 (2)	H15A—C15—H15B	108.7
C9—C8—H8A	108.6	C17—C16—C21	116.28 (18)
C7—C8—H8A	108.6	C17—C16—C10	123.98 (17)
C9—C8—H8B	108.6	C21—C16—C10	119.74 (17)
C7—C8—H8B	108.6	C18—C17—C16	122.04 (19)
H8A—C8—H8B	107.6	C18—C17—C11	117.49 (16)
C8—C9—H9A	109.5	C16—C17—C11	120.46 (15)
C8—C9—H9B	109.5	C19—C18—C17	120.3 (2)
H9A—C9—H9B	109.5	C19—C18—H18	119.9
C8—C9—H9C	109.5	C17—C18—H18	119.9
H9A—C9—H9C	109.5	C18—C19—C20	119.2 (2)
H9B—C9—H9C	109.5	C18—C19—H19	120.4
N1—C10—C11	109.45 (15)	C20—C19—H19	120.4
N1—C10—C16	113.53 (16)	C21—C20—C19	120.0 (2)
C11—C10—C16	111.07 (15)	C21—C20—H20	120.0
N1—C10—H10	107.5	C19—C20—H20	120.0
C11—C10—H10	107.5	C20—C21—C16	122.1 (2)
C16—C10—H10	107.5	C20—C21—H21	119.0
C10—C11—C15	113.65 (17)	C16—C21—H21	119.0
C10—C11—C12	115.59 (17)	C10—N1—C7	116.59 (15)
C15—C11—C12	102.53 (18)	C10—N1—H1	108.9 (16)
C10—C11—H11	108.2	C7—N1—H1	113.1 (16)
C15—C11—H11	108.2	C4—O1—H1A	109.5
C6—C1—C2—C3	2.6 (3)	C11—C12—C13—C14	-27.7 (3)
Cl2—C1—C2—C3	-176.33 (18)	C12—C13—C14—C15	6.6 (3)
C1—C2—C3—C4	-0.2 (3)	C13—C14—C15—C11	17.1 (3)

C2—C3—C4—O1	177.9 (2)	C10—C11—C15—C14	-159.2 (2)
C2—C3—C4—C5	-2.2 (3)	C12—C11—C15—C14	-33.7 (2)
O1—C4—C5—C6	-177.83 (19)	N1—C10—C16—C17	-122.7 (2)
C3—C4—C5—C6	2.2 (3)	C11—C10—C16—C17	113.4 (2)
O1—C4—C5—C7	-0.6 (3)	N1—C10—C16—C21	58.1 (2)
C3—C4—C5—C7	179.43 (19)	C11—C10—C16—C21	-65.8 (2)
C2—C1—C6—C5	-2.6 (3)	C21—C16—C17—C18	-0.5 (3)
Cl2—C1—C6—C5	176.38 (15)	C10—C16—C17—C18	-179.69 (18)
C4—C5—C6—C1	0.1 (3)	C21—C16—C17—Cl1	178.20 (14)
C7—C5—C6—C1	-177.13 (18)	C10—C16—C17—Cl1	-1.0 (3)
C6—C5—C7—N1	-145.10 (17)	C16—C17—C18—C19	0.0 (3)
C4—C5—C7—N1	37.7 (2)	Cl1—C17—C18—C19	-178.73 (18)
C6—C5—C7—C8	95.3 (2)	C17—C18—C19—C20	0.7 (4)
C4—C5—C7—C8	-81.9 (2)	C18—C19—C20—C21	-0.9 (4)
N1—C7—C8—C9	178.4 (2)	C19—C20—C21—C16	0.4 (3)
C5—C7—C8—C9	-60.7 (3)	C17—C16—C21—C20	0.3 (3)
N1—C10—C11—C15	175.01 (17)	C10—C16—C21—C20	179.5 (2)
C16—C10—C11—C15	-58.8 (2)	C11—C10—N1—C7	179.95 (17)
N1—C10—C11—C12	56.8 (2)	C16—C10—N1—C7	55.2 (2)
C16—C10—C11—C12	-177.03 (17)	C5—C7—N1—C10	71.5 (2)
C10—C11—C12—C13	161.7 (2)	C8—C7—N1—C10	-165.80 (17)
C15—C11—C12—C13	37.5 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···Cl2 ⁱ	0.848 (19)	2.913 (13)	3.7023 (18)	156 (2)
O1—H1A···N1	0.82	1.93	2.642 (2)	144

Symmetry codes: (i) $-x+1/2, -y+2, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

