

Methyl 5-bromo-6-methylpicolinate

Ya-Ming Wu,^{a*} Chun-Ming Wu^b and Yong Wang^c

^aDepartment of Applied Chemistry, Nanjing College of Chemical Technology, No. 625 Geguan Road, Dachang, Nanjing 210048, People's Republic of China,

^bSynergetica-Changzhou, Ltd., Chunjiang Town, Xinbei District, Changzhou City 213033, Jiangsu Province, People's Republic of China, and ^cSchool of Pharmaceutical and Chemical Engineering, Taizhou University, Linhai 317000, People's Republic of China

Correspondence e-mail: adsny05@163.com


Received 8 December 2008; accepted 10 December 2008

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; R factor = 0.061; wR factor = 0.066; data-to-parameter ratio = 14.6.

The title compound, $\text{C}_8\text{H}_8\text{BrNO}_2$, does not show any significant intermolecular $\pi-\pi$ or $\text{C}-\text{H}\cdots\pi$ interactions in the crystal packing except for one weak $\text{Br}\cdots\text{Br}$ [3.715 (1) \AA] interaction.

Related literature

The title compound is an important intermediate for the construction of novel supported PyOX ligands, see: Oila *et al.* (2005). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_8\text{H}_8\text{BrNO}_2$

$M_r = 230.06$

Monoclinic, $P2_1/c$
 $a = 18.518 (4)\text{ \AA}$
 $b = 4.1040 (8)\text{ \AA}$
 $c = 12.442 (3)\text{ \AA}$
 $\beta = 109.52 (3)^\circ$
 $V = 891.2 (4)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 4.57\text{ mm}^{-1}$
 $T = 293 (2)\text{ K}$
 $0.20 \times 0.10 \times 0.10\text{ mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.462$, $T_{\max} = 0.658$
1602 measured reflections

1602 independent reflections
975 reflections with $I > 2\sigma(I)$
3 standard reflections
every 200 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.066$
 $S = 1.75$
1602 reflections

110 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.53\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.61\text{ e \AA}^{-3}$

Data collection: *CAD-4 Software* (Enraf-Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2083).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Enraf-Nonius (1985). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
Oila, M. J., Tois, J. E. & Koskinen, A. M. P. (2005). *Tetrahedron*, **61**, 10748–10756.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o134 [doi:10.1107/S1600536808042104]

Methyl 5-bromo-6-methylpicolinate

Y.-M. Wu, C.-M. Wu and Y. Wang

Comment

The title compound is one of important intermediates for construction of novel supported PyOX-ligands (Oila *et al.*, 2005). Here we report the crystal structure of the title compound, methyl 5-bromo-6-methylpicolinate (Fig. 1).

In the title compound, the bond lengths and angles are within normal ranges (Allen *et al.*, 1987). The crystal structure is stabilized by a weak Br \cdots Brⁱ interaction at 3.715 (1) Å (Fig. 2; symmetry code as in Fig. 2).

Experimental

The title compound, (I) was prepared by a method reported in literature (Oila *et al.*, 2005) with some modification. The crystals were obtained by dissolving I (0.2 g) in methanol (50 ml) and evaporating the solvent slowly at room temperature for about 3 d.

Refinement

H atoms were positioned geometrically, with O—H = 0.82 and C—H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}/\text{O})$, where $x = 1.2$ for aromatic H and $x = 1.5$ for other H.

Figures




Fig. 1. A drawing of the title molecular structure, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. Br \cdots Br interaction in the title compound. [Symmetry code: (i) -x+1, y+1/2, -z+1/2.]

Methyl 5-bromo-6-methylpicolinate

Crystal data

C₈H₈BrNO₂

$F_{000} = 456$

supplementary materials

$M_r = 230.06$	$D_x = 1.715 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 18.518 (4) \text{ \AA}$	Cell parameters from 25 reflections
$b = 4.1040 (8) \text{ \AA}$	$\theta = 10\text{--}14^\circ$
$c = 12.442 (3) \text{ \AA}$	$\mu = 4.57 \text{ mm}^{-1}$
$\beta = 109.52 (3)^\circ$	$T = 293 (2) \text{ K}$
$V = 891.2 (4) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.20 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Enraf-Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.0000$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.3^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 1.2^\circ$
$T = 293(2) \text{ K}$	$h = -22 \rightarrow 20$
$\omega/2\theta$ scans	$k = 0 \rightarrow 4$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = 0 \rightarrow 14$
$T_{\text{min}} = 0.462$, $T_{\text{max}} = 0.658$	3 standard reflections
1602 measured reflections	every 200 reflections
1602 independent reflections	intensity decay: 1%
975 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.061$	H-atom parameters constrained
$wR(F^2) = 0.066$	$w = 1/[\sigma^2(F_o^2)]$
$S = 1.75$	$(\Delta/\sigma)_{\text{max}} < 0.000$
1602 reflections	$\Delta\rho_{\text{max}} = 0.53 \text{ e \AA}^{-3}$
110 parameters	$\Delta\rho_{\text{min}} = -0.61 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculat-

ing R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.41815 (4)	0.6507 (2)	0.25731 (6)	0.0546 (3)
O1	0.0882 (2)	-0.1451 (13)	0.2125 (3)	0.0691 (16)
O2	0.1349 (2)	-0.0435 (11)	0.3986 (4)	0.0533 (15)
N	0.2635 (3)	0.2009 (15)	0.3797 (4)	0.0450 (17)
C1	0.3893 (3)	0.4149 (17)	0.4811 (4)	0.067 (2)
H1A	0.3707	0.3787	0.5434	0.101*
H1B	0.4059	0.6370	0.4822	0.101*
H1C	0.4317	0.2717	0.4882	0.101*
C2	0.3260 (3)	0.3477 (18)	0.3698 (5)	0.0344 (16)
C3	0.3290 (3)	0.4444 (16)	0.2646 (5)	0.039 (2)
C4	0.2680 (3)	0.3789 (19)	0.1651 (5)	0.059 (2)
H4A	0.2702	0.4358	0.0939	0.071*
C5	0.2053 (3)	0.2298 (18)	0.1755 (5)	0.050 (2)
H5A	0.1632	0.1866	0.1109	0.060*
C6	0.2038 (3)	0.1425 (19)	0.2814 (5)	0.0426 (18)
C7	0.1372 (4)	-0.0335 (18)	0.2958 (6)	0.050 (2)
C8	0.0676 (3)	-0.1896 (19)	0.4127 (5)	0.070 (2)
H8A	0.0724	-0.1875	0.4920	0.104*
H8B	0.0627	-0.4104	0.3857	0.104*
H8C	0.0230	-0.0679	0.3698	0.104*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0482 (4)	0.0590 (5)	0.0589 (5)	-0.0020 (6)	0.0211 (3)	0.0028 (6)
O1	0.048 (3)	0.092 (4)	0.055 (4)	-0.011 (4)	0.001 (3)	-0.013 (4)
O2	0.043 (3)	0.070 (4)	0.046 (3)	-0.007 (3)	0.013 (2)	-0.002 (3)
N	0.029 (3)	0.065 (5)	0.033 (3)	-0.006 (4)	0.000 (3)	0.004 (4)
C1	0.056 (4)	0.087 (7)	0.043 (4)	-0.027 (5)	-0.004 (4)	0.012 (5)
C2	0.026 (4)	0.031 (4)	0.035 (4)	-0.003 (4)	-0.004 (3)	-0.004 (5)
C3	0.034 (4)	0.045 (6)	0.040 (4)	0.002 (4)	0.014 (4)	0.009 (4)
C4	0.049 (4)	0.088 (7)	0.033 (4)	0.004 (6)	0.005 (4)	0.031 (5)
C5	0.038 (4)	0.076 (7)	0.029 (4)	0.001 (4)	0.002 (3)	0.002 (4)
C6	0.025 (4)	0.060 (5)	0.042 (4)	0.004 (5)	0.011 (3)	-0.001 (5)
C7	0.037 (5)	0.060 (6)	0.043 (5)	0.004 (4)	0.000 (4)	-0.010 (5)
C8	0.051 (4)	0.079 (7)	0.089 (5)	-0.011 (5)	0.037 (4)	0.010 (6)

Geometric parameters (\AA , $^\circ$)


Br—Br ⁱ	3.715 (1)	C2—C3	1.387 (6)
Br—C3	1.884 (5)	C3—C4	1.396 (6)
O1—C7	1.218 (6)	C4—C5	1.356 (7)

supplementary materials

O2—C7	1.294 (6)	C4—H4A	0.9300
O2—C8	1.446 (6)	C5—C6	1.375 (7)
N—C2	1.347 (6)	C5—H5A	0.9300
N—C6	1.368 (6)	C6—C7	1.491 (8)
C1—C2	1.510 (6)	C8—H8A	0.9600
C1—H1A	0.9600	C8—H8B	0.9600
C1—H1B	0.9600	C8—H8C	0.9600
C1—H1C	0.9600		
C7—O2—C8	116.6 (5)	C3—C4—H4A	121.0
C2—N—C6	117.3 (5)	C4—C5—C6	120.0 (6)
C2—C1—H1A	109.5	C4—C5—H5A	120.0
C2—C1—H1B	109.5	C6—C5—H5A	120.0
H1A—C1—H1B	109.5	N—C6—C5	122.9 (6)
C2—C1—H1C	109.5	N—C6—C7	115.5 (6)
H1A—C1—H1C	109.5	C5—C6—C7	121.6 (6)
H1B—C1—H1C	109.5	O1—C7—O2	124.6 (7)
N—C2—C3	121.4 (5)	O1—C7—C6	119.4 (7)
N—C2—C1	115.2 (5)	O2—C7—C6	115.9 (6)
C3—C2—C1	123.3 (6)	O2—C8—H8A	109.5
C4—C3—C2	120.4 (5)	O2—C8—H8B	109.5
C4—C3—Br	120.5 (5)	H8A—C8—H8B	109.5
C2—C3—Br	119.1 (5)	O2—C8—H8C	109.5
C5—C4—C3	117.9 (6)	H8A—C8—H8C	109.5
C5—C4—H4A	121.0	H8B—C8—H8C	109.5
C6—N—C2—C3	1.8 (10)	C2—N—C6—C7	177.4 (6)
C6—N—C2—C1	178.4 (6)	C4—C5—C6—N	0.4 (11)
N—C2—C3—C4	-2.7 (11)	C4—C5—C6—C7	-177.5 (7)
C1—C2—C3—C4	-179.0 (6)	C8—O2—C7—O1	-2.1 (11)
N—C2—C3—Br	179.6 (5)	C8—O2—C7—C6	175.0 (6)
C1—C2—C3—Br	3.3 (9)	N—C6—C7—O1	-167.1 (7)
C2—C3—C4—C5	2.3 (11)	C5—C6—C7—O1	11.0 (11)
Br—C3—C4—C5	180.0 (5)	N—C6—C7—O2	15.6 (9)
C3—C4—C5—C6	-1.2 (11)	C5—C6—C7—O2	-166.3 (7)
C2—N—C6—C5	-0.6 (10)		

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

