

2-(4-Chlorophenyl)-3-methyl-N-(5-methylthiazol-2-yl)butanamide

Jing-Li Cheng,^a Jin-Hao Zhao,^a Guo-Nian Zhu^a and Fu-Cheng Lin^{b*}

^aCollege of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310029, People's Republic of China, and ^bInstitute of Biotechnology, Zhejiang University, Hangzhou 310029, People's Republic of China
Correspondence e-mail: fuchenglin@zj.com


Received 19 November 2008; accepted 17 December 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.172; data-to-parameter ratio = 20.3.

In the title compound, $\text{C}_{15}\text{H}_{17}\text{ClN}_2\text{OS}$, the thiazole ring, which is essentially planar with a maximum deviation of 0.044 (3) \AA , makes a dihedral angle of 54.76 (8) $^\circ$ with the benzene ring. In the crystal, adjacent molecules related by twofold rotation symmetry are linked by pairs of $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds.

Related literature

For background, see: Holmstead *et al.* (1978); Forlani (1978). For a related structure, see: Zhao *et al.* (2006).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{17}\text{ClN}_2\text{OS}$
 $M_r = 308.83$
Monoclinic, $C2/c$
 $a = 14.9649 (6)\text{ \AA}$
 $b = 17.6062 (7)\text{ \AA}$
 $c = 12.5606 (5)\text{ \AA}$
 $\beta = 99.9482 (11)^\circ$
 $V = 3259.6 (2)\text{ \AA}^3$

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.36\text{ mm}^{-1}$

$T = 298 (1)\text{ K}$
 $0.41 \times 0.33 \times 0.26\text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.858$, $T_{\max} = 0.911$

15655 measured reflections
3708 independent reflections
2559 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.172$
 $S = 1.01$
3708 reflections

183 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H111…N2 ⁱ	0.86	2.08	2.929 (2)	168

Symmetry code: (i) $-x + 1, y, -z + \frac{3}{2}$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

The authors are grateful for support from the National Natural Science Foundation of China (No. 30700532) and also thank Professor Jian-Ming Gu for help with the analysis of the crystal data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2368).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Forlani, L. (1978). *J. Chem. Soc. Perkin Trans. 1*, pp. 1169–1171.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Holmstead, R. L., Fullmer, D. G. & Ruzo, L. O. (1978). *J. Agric. Food Chem.* **26**, 954–959.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MSC (2004). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Zhao, J.-H., Cheng, J.-L., Huang, Y.-K. & Zhu, G.-N. (2006). *Acta Cryst. E* **62**, o4840–o4841.

supplementary materials

Acta Cryst. (2009). E65, o184 [doi:10.1107/S1600536808043031]

2-(4-Chlorophenyl)-3-methyl-N-(5-methylthiazol-2-yl)butanamide

J.-L. Cheng, J.-H. Zhao, G.-N. Zhu and F.-C. Lin

Comment

2-(4-Chlorophenyl)-3-methylbutanoyl chloride is an intermediate in the synthesis of fenvalerate, an excellent insecticide (Holmstead *et al.*, 1978). 2-Amino-5-methyl-thiazole is another heterocyclic intermediate (Forlani, 1978). As part of our continuing interest in the design and synthesis of new pesticides, we have isolated the title compound, (I), the product of the condensation reaction between 2-(4-chlorophenyl)-3-methylbutanoyl chloride and 5-methyl-2-aminothiazole, as colourless crystals suitable for X-ray analysis.

The molecular structure of (I) is illustrated in Fig. 1. Atoms N2, C10, C11, S1, C9 and N1 are coplanar, the largest deviation being 0.044 (3) Å for N1. As expected, the benzene ring is planar, and atom C11 lies only 0.018 (4) Å from the plane defined by the ring C atoms and itself. The angle between these two rings is 54.76 (8)°, smaller than the angle between the thiazole and benzene rings of the compound 2-(4-chlorophenyl)-3-methyl-N-(thiazol-2-yl) butanamide (Zhao *et al.*, 2006). There are N—H···N interactions in the crystal structure, which lead to the formation of hydrogen-bonded dimers (Figs. 2 and 3).

Experimental

2-Amino-5-methylthiazole (1.14 g, 10 mmol), 4-dimethylaminopyridine (0.12 g), triethylamine (1.31 g) and chloroform (100 ml) were added to a 250 ml round flask. The mixture was stirred and cooled to 273 K, and then 2-(4-chlorophenyl)-3-methylbutanoyl chloride (3.47 g) was added dropwise within 30 min. The mixture was stirred at room temperature for 3 h and then 1% aqueous HCl was added (5 ml). The organic layer was washed with water to a neutral pH and dried over Na₂SO₄. After being filtered and concentrated, the organic residue was purified by silica-gel column chromatography, eluted with ethyl acetate-petroleum ether-formic acid (10:80:1, v/v/v), to give a white solid (yield 85%, 2.5 g), (I). It was then recrystallized from ethyl acetate-petroleum ether (2:1, v/v) to give colourless blocks (m.p. 460–461 K).

Refinement

H atoms were included in calculated positions and refined using a riding model, with C—H distances constrained to 0.96 Å for methyl H atoms, 0.93 Å for aryl H atoms and 0.98 Å for the remainder, with N—H distances constrained to 0.86 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. The molecular structure of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. Molecular packing arrangement in the unit cell.




Fig. 3. View showing the N—H···N hydrogen bonding (dashed lines) [symmetry code: (i) 1 - $x, y, 3/2 - z$].

2-(4-Chlorophenyl)-3-methyl-N-(5-methylthiazol-2-yl)butanamide

Crystal data

C ₁₅ H ₁₇ ClN ₂ OS	$F_{000} = 1296.00$
$M_r = 308.83$	$D_x = 1.258 \text{ Mg m}^{-3}$
Monoclinic, C2/c	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71075 \text{ \AA}$
$a = 14.9649 (6) \text{ \AA}$	Cell parameters from 10539 reflections
$b = 17.6062 (7) \text{ \AA}$	$\theta = 3.3\text{--}27.4^\circ$
$c = 12.5606 (5) \text{ \AA}$	$\mu = 0.36 \text{ mm}^{-1}$
$\beta = 99.9482 (11)^\circ$	$T = 298 (1) \text{ K}$
$V = 3259.6 (2) \text{ \AA}^3$	Block, colorless
$Z = 8$	$0.41 \times 0.33 \times 0.26 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer	2559 reflections with $F^2 > 2\sigma(F^2)$
Detector resolution: 10.00 pixels mm ⁻¹	$R_{\text{int}} = 0.027$
ω scans	$\theta_{\text{max}} = 27.4^\circ$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$h = -19 \rightarrow 19$
$T_{\text{min}} = 0.858, T_{\text{max}} = 0.911$	$k = -22 \rightarrow 22$
15655 measured reflections	$l = -16 \rightarrow 15$
3708 independent reflections	

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.121P)^2]$
	where $P = (F_o^2 + 2F_c^2)/3$
$R[F^2 > 2\sigma(F^2)] = 0.040$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$wR(F^2) = 0.172$	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$

$S = 1.01$ $\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$
 3708 reflections Extinction correction: SHELXL
 183 parameters Extinction coefficient: 0.0028 (7)
 H-atom parameters constrained

Special details

Geometry. ENTER SPECIAL DETAILS OF THE MOLECULAR GEOMETRY

Refinement. Refinement using all reflections. The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R -factor (gt) are based on F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.02408 (6)	0.70387 (4)	0.59960 (9)	0.1206 (3)
S1	0.46203 (3)	0.39223 (3)	0.44117 (3)	0.0559 (2)
O1	0.29216 (10)	0.40983 (10)	0.48366 (10)	0.0696 (4)
N1	0.40090 (10)	0.41256 (9)	0.63283 (11)	0.0496 (3)
N2	0.55713 (11)	0.40575 (9)	0.63079 (12)	0.0530 (4)
C1	0.24241 (12)	0.41401 (12)	0.65622 (13)	0.0534 (4)
C2	0.18766 (12)	0.48677 (12)	0.64116 (13)	0.0531 (4)
C3	0.18371 (13)	0.53424 (12)	0.72811 (17)	0.0611 (5)
C4	0.13313 (17)	0.60024 (12)	0.7171 (2)	0.0739 (6)
C5	0.08679 (16)	0.61988 (13)	0.6158 (2)	0.0756 (6)
C6	0.08973 (17)	0.57396 (14)	0.5282 (2)	0.0790 (6)
C7	0.13902 (13)	0.50774 (13)	0.54031 (16)	0.0667 (5)
C8	0.31263 (12)	0.41250 (11)	0.58189 (13)	0.0506 (4)
C9	0.47375 (12)	0.40501 (10)	0.57886 (13)	0.0452 (4)
C10	0.61571 (13)	0.39512 (12)	0.55850 (16)	0.0583 (5)
C11	0.57888 (13)	0.38625 (12)	0.45425 (16)	0.0563 (5)
C12	0.62499 (18)	0.37260 (17)	0.35901 (19)	0.0818 (7)
C13	0.18427 (14)	0.34156 (12)	0.63811 (17)	0.0646 (5)
C14	0.24379 (18)	0.27066 (14)	0.6583 (2)	0.0850 (7)
C15	0.11223 (17)	0.34038 (16)	0.7103 (2)	0.0856 (7)
H1	0.2751	0.4131	0.7309	0.064*
H3	0.2160	0.5213	0.7957	0.073*
H4	0.1303	0.6309	0.7766	0.089*
H6	0.0583	0.5877	0.4605	0.095*
H7	0.1400	0.4765	0.4808	0.080*
H10	0.6782	0.3942	0.5812	0.070*
H13	0.1535	0.3409	0.5626	0.078*
H111	0.4119	0.4175	0.7020	0.060*
H121	0.6896	0.3732	0.3823	0.098*
H122	0.6081	0.4118	0.3062	0.098*
H123	0.6068	0.3241	0.3276	0.098*
H141	0.2729	0.2694	0.7326	0.102*
H142	0.2890	0.2719	0.6126	0.102*
H143	0.2068	0.2262	0.6422	0.102*

supplementary materials

H151	0.0750	0.2960	0.6946	0.103*
H152	0.0752	0.3850	0.6972	0.103*
H153	0.1412	0.3393	0.7848	0.103*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.1126 (6)	0.0730 (4)	0.1748 (9)	0.0221 (3)	0.0210 (5)	0.0226 (5)
S1	0.0557 (3)	0.0783 (3)	0.0340 (2)	-0.0055 (2)	0.0085 (2)	-0.00354 (19)
O1	0.0527 (8)	0.1174 (13)	0.0374 (7)	0.0055 (7)	0.0038 (5)	-0.0021 (6)
N1	0.0439 (8)	0.0725 (9)	0.0325 (7)	0.0020 (6)	0.0070 (5)	-0.0005 (6)
N2	0.0462 (8)	0.0767 (10)	0.0363 (7)	-0.0003 (6)	0.0074 (6)	-0.0030 (6)
C1	0.0451 (10)	0.0762 (12)	0.0382 (8)	0.0032 (8)	0.0052 (7)	0.0008 (8)
C2	0.0412 (9)	0.0713 (11)	0.0466 (9)	-0.0028 (8)	0.0073 (7)	0.0040 (8)
C3	0.0579 (11)	0.0705 (12)	0.0553 (10)	0.0004 (9)	0.0109 (8)	-0.0006 (9)
C4	0.0711 (15)	0.0687 (14)	0.0854 (17)	-0.0069 (10)	0.0233 (12)	-0.0088 (11)
C5	0.0608 (13)	0.0676 (13)	0.0987 (19)	-0.0022 (10)	0.0142 (12)	0.0143 (13)
C6	0.0658 (14)	0.0892 (17)	0.0771 (15)	0.0059 (11)	-0.0015 (11)	0.0222 (13)
C7	0.0589 (11)	0.0826 (14)	0.0546 (11)	0.0032 (10)	-0.0012 (8)	0.0027 (10)
C8	0.0454 (9)	0.0673 (11)	0.0388 (9)	0.0019 (7)	0.0060 (7)	0.0025 (7)
C9	0.0487 (9)	0.0531 (9)	0.0338 (7)	-0.0006 (6)	0.0071 (6)	0.0013 (6)
C10	0.0463 (10)	0.0830 (14)	0.0472 (10)	-0.0028 (8)	0.0127 (8)	-0.0056 (8)
C11	0.0564 (11)	0.0683 (11)	0.0465 (10)	-0.0072 (8)	0.0159 (8)	-0.0059 (8)
C12	0.0754 (15)	0.118 (2)	0.0586 (12)	-0.0162 (14)	0.0299 (11)	-0.0212 (13)
C13	0.0569 (11)	0.0775 (13)	0.0589 (11)	-0.0026 (9)	0.0083 (9)	0.0043 (10)
C14	0.0848 (17)	0.0730 (15)	0.0990 (19)	0.0032 (12)	0.0214 (14)	0.0015 (13)
C15	0.0702 (15)	0.0908 (17)	0.1022 (19)	-0.0038 (12)	0.0331 (13)	0.0192 (14)

Geometric parameters (\AA , $^\circ$)

Cl1—C5	1.744 (2)	C13—C14	1.529 (3)
S1—C9	1.7227 (17)	C13—C15	1.524 (3)
S1—C11	1.731 (2)	N1—H111	0.860
O1—C8	1.220 (2)	C1—H1	0.980
N1—C8	1.365 (2)	C3—H3	0.930
N1—C9	1.386 (2)	C4—H4	0.930
N2—C9	1.304 (2)	C6—H6	0.930
N2—C10	1.380 (2)	C7—H7	0.930
C1—C2	1.515 (2)	C10—H10	0.930
C1—C8	1.522 (2)	C12—H121	0.960
C1—C13	1.539 (2)	C12—H122	0.960
C2—C3	1.385 (2)	C12—H123	0.960
C2—C7	1.397 (2)	C13—H13	0.980
C3—C4	1.381 (3)	C14—H141	0.960
C4—C5	1.384 (3)	C14—H142	0.960
C5—C6	1.372 (3)	C14—H143	0.960
C6—C7	1.374 (3)	C15—H151	0.960
C10—C11	1.339 (2)	C15—H152	0.960
C11—C12	1.500 (3)	C15—H153	0.960

C9—S1—C11	89.24 (9)	C13—C1—H1	107.7
C8—N1—C9	123.37 (14)	C2—C3—H3	119.1
C9—N2—C10	109.35 (15)	C4—C3—H3	119.1
C2—C1—C8	110.74 (16)	C3—C4—H4	120.6
C2—C1—C13	113.75 (15)	C5—C4—H4	120.6
C8—C1—C13	109.12 (16)	C5—C6—H6	119.9
C1—C2—C3	120.52 (15)	C7—C6—H6	119.9
C1—C2—C7	121.50 (17)	C2—C7—H7	119.7
C3—C2—C7	117.97 (19)	C6—C7—H7	119.7
C2—C3—C4	121.75 (19)	N2—C10—H10	121.4
C3—C4—C5	118.8 (2)	C11—C10—H10	121.4
C11—C5—C4	119.5 (2)	C11—C12—H121	109.5
C11—C5—C6	119.8 (2)	C11—C12—H122	109.5
C4—C5—C6	120.6 (2)	C11—C12—H123	109.5
C5—C6—C7	120.1 (2)	H121—C12—H122	109.5
C2—C7—C6	120.7 (2)	H121—C12—H123	109.5
O1—C8—N1	121.86 (17)	H122—C12—H123	109.5
O1—C8—C1	122.81 (15)	C1—C13—H13	108.2
N1—C8—C1	115.30 (14)	C14—C13—H13	108.2
S1—C9—N1	123.45 (12)	C15—C13—H13	108.2
S1—C9—N2	115.22 (14)	C13—C14—H141	109.5
N1—C9—N2	121.32 (15)	C13—C14—H142	109.5
N2—C10—C11	117.29 (18)	C13—C14—H143	109.5
S1—C11—C10	108.91 (16)	H141—C14—H142	109.5
S1—C11—C12	122.03 (14)	H141—C14—H143	109.5
C10—C11—C12	129.06 (19)	H142—C14—H143	109.5
C1—C13—C14	110.73 (18)	C13—C15—H151	109.5
C1—C13—C15	111.30 (18)	C13—C15—H152	109.5
C14—C13—C15	110.0 (2)	C13—C15—H153	109.5
C8—N1—H111	118.3	H151—C15—H152	109.5
C9—N1—H111	118.3	H151—C15—H153	109.5
C2—C1—H1	107.7	H152—C15—H153	109.5
C8—C1—H1	107.7		
C9—S1—C11—C10	-0.73 (16)	C13—C1—C2—C7	-65.9 (2)
C9—S1—C11—C12	178.9 (2)	C8—C1—C13—C14	57.6 (2)
C11—S1—C9—N1	-178.04 (16)	C8—C1—C13—C15	-179.67 (16)
C11—S1—C9—N2	0.70 (15)	C13—C1—C8—O1	60.1 (2)
C8—N1—C9—S1	-1.7 (2)	C13—C1—C8—N1	-118.44 (17)
C8—N1—C9—N2	179.66 (17)	C1—C2—C3—C4	-178.9 (2)
C9—N1—C8—O1	-4.1 (2)	C1—C2—C7—C6	-180.0 (2)
C9—N1—C8—C1	174.50 (16)	C3—C2—C7—C6	0.7 (3)
C9—N2—C10—C11	-0.2 (2)	C7—C2—C3—C4	0.4 (3)
C10—N2—C9—S1	-0.4 (2)	C2—C3—C4—C5	-1.3 (3)
C10—N2—C9—N1	178.33 (16)	C3—C4—C5—Cl1	-178.89 (19)
C2—C1—C8—O1	-65.9 (2)	C3—C4—C5—C6	1.1 (3)
C2—C1—C8—N1	115.61 (17)	C11—C5—C6—C7	179.97 (13)
C8—C1—C2—C3	-123.34 (19)	C4—C5—C6—C7	-0.0 (3)
C8—C1—C2—C7	57.4 (2)	C5—C6—C7—C2	-0.9 (3)

supplementary materials


C2—C1—C13—C14	−178.17 (17)	N2—C10—C11—S1	0.7 (2)
C2—C1—C13—C15	−55.5 (2)	N2—C10—C11—C12	−178.9 (2)
C13—C1—C2—C3	113.3 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H111…N2 ⁱ	0.86	2.08	2.929 (2)	168

Symmetry codes: (i) $-x+1, y, -z+3/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

