

3-(2-Aminoethyl)-2-(4-chloroanilino)-quinazolin-4(3H)-one methanol 0.75-solvate

Xu-Hong Yang,^{a*} Xiao-Bao Chen^b and Si-Xuan Zhou^c

^aDepartment of Chemistry and Life Science, Xianning College, Xianning 4371000, Hubei, People's Republic of China, ^bDepartment of Medicinal Chemistry, Yunyang Medical College, Shiyuan 442000, Hubei, People's Republic of China, and ^cSchool of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, Hunan, People's Republic of China

Correspondence e-mail: yxh502@163.com


Received 10 November 2008; accepted 2 December 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in solvent or counterion; R factor = 0.045; wR factor = 0.132; data-to-parameter ratio = 14.2.

In the asymmetric unit of the title compound, $\text{C}_{16}\text{H}_{15}\text{ClN}_4\text{O} \cdot 0.75\text{CH}_3\text{OH}$, there are two independent quinazolin-4(3H)-one molecules and one and a half methanol molecules. One of the methanol molecules is disordered over two positions with equal occupancies. The dihedral angles between the quinazolinone ring system and the chlorobenzene ring in the two quinazolin-4(3H)-one molecules are essentially the same, at 39.83 (1) and 39.84 (1)°. Intramolecular $\text{N}-\text{H} \cdots \text{N}$ and $\text{O}-\text{H} \cdots \text{O}$, and intermolecular $\text{N}-\text{H} \cdots \text{O}$ and $\text{N}-\text{H} \cdots \text{N}$ hydrogen bonds are observed. In addition, $\pi-\pi$ stacking interactions, with centroid-to-centroid distances of 3.654 (1), 3.766 (1) and 3.767 (1) Å, and weak $\text{C}-\text{H} \cdots \pi$ interactions, are observed.

Related literature

For the biological activity of quinazolin-4(3H)-one derivatives, see: Bartroli *et al.* (1998); Kung *et al.* (1999); Malamas & Millen (1991); Mannschreck *et al.* (1984); Matsuno *et al.* (2002); Palmer *et al.* (1997); Pandeya *et al.* (1999); Shiba *et al.* (1997); Tsou *et al.* (2001). For the synthesis of the title compound, see: Hu *et al.* (2006); Yang *et al.* (2008).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{15}\text{ClN}_4\text{O} \cdot 0.75\text{CH}_3\text{O}$
 $M_r = 338.81$
 Monoclinic, $P2_1/n$
 $a = 13.380$ (3) Å
 $b = 12.048$ (2) Å
 $c = 21.105$ (4) Å
 $\beta = 104.49$ (3)°
 $V = 3293.8$ (11) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.25$ mm⁻¹
 $T = 293$ (2) K
 $0.23 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: none
 18726 measured reflections
 6451 independent reflections
 4529 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.132$
 $S = 1.07$
 6451 reflections
 453 parameters
 9 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.40$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N5}-\text{H5A} \cdots \text{N8}$	0.863 (9)	1.933 (11)	2.787 (3)	170 (2)
$\text{O4}-\text{H4C} \cdots \text{O1}$	0.82	2.15	2.889 (6)	151
$\text{O3}-\text{H3A} \cdots \text{O2}$	0.835 (10)	1.917 (11)	2.744 (3)	170 (2)
$\text{N1}-\text{H1A} \cdots \text{N4}$	0.877 (9)	1.904 (11)	2.760 (3)	164 (2)
$\text{N8}-\text{H8B} \cdots \text{O4}^i$	0.851 (10)	2.467 (14)	3.284 (8)	161 (2)
$\text{N8}-\text{H8A} \cdots \text{N2}^{ii}$	0.861 (10)	2.440 (17)	3.160 (2)	142 (2)
$\text{N4}-\text{H4B} \cdots \text{O3}^{iii}$	0.858 (10)	2.097 (11)	2.941 (3)	168 (2)
$\text{N4}-\text{H4A} \cdots \text{N6}^{iv}$	0.853 (10)	2.491 (17)	3.223 (3)	144 (2)
$\text{C5}-\text{H5} \cdots \text{Cg4}^{iv}$	0.93	2.81	3.375 (2)	120
$\text{C17}-\text{H17} \cdots \text{Cg1}^{ii}$	0.93	2.84	3.459 (2)	125
$\text{C32}-\text{H32A} \cdots \text{Cg2}^{ii}$	0.97	2.98	3.824 (3)	146

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y, z$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x - 1, y, z$. Cg1 , Cg2 and Cg4 are the centroids of the $\text{N2/C7/N3/C8}-\text{C10}$, $\text{C1}-\text{C6}$ and $\text{N6/C23/N7/C24}-\text{C26}$ rings, respectively.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors gratefully acknowledge financial support of this work by Xianning College, and acknowledge the Sophisticated Analytical Instrument Facility, Central China Normal University, Wuhan, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2361).

References

- Bartroli, J., Turmo, E., Alguero, M., Boncompte, E., Vericat, M. L., Conte, L., Ramis, J., Merlos, M., Garcia-Rafanell, J. & Forn, J. (1998). *J. Med. Chem.* **41**, 1869–1882.
- Bruker (2000). *SMART and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hu, Y.-G., Zheng, A.-H. & Li, G.-H. (2006). *Acta Cryst.* **E62**, o1457–o1459.
- Kung, P. P., Casper, M. D., Cook, K. L., Wilson-Lingardo, L., Risen, L. M., Vickers, T. A., Ranken, R., Blyn, L. B., Wyatt, J. R., Cook, P. & Decker, D. J. (1999). *J. Med. Chem.* **42**, 4705–4713.
- Malamas, M. S. & Millen, J. (1991). *J. Med. Chem.* **34**, 1492–1503.
- Mannschreck, A., Koller, H., Stuhler, G., Davis, M. A. & Traber, J. (1984). *Eur. J. Med. Chem.* **19**, 381–383.
- Matsuno, K., Ichimura, M., Nakajima, T., Tahara, K., Fujiwara, S., Kase, H., Ushiki, J., Giese, N. A., Pandey, A., Scarborough, R. M., Lokker, N. A., Yu, J. C., Irie, J., Tsukuda, E., Ide, S., Oda, S. & Nomoto, Y. (2002). *J. Med. Chem.* **45**, 3057–3066.
- Palmer, B. D., Trumpp-Kallmeyer, S., Fry, D. W., Nelson, J. M., Showalter, H. D. H. & Denny, W. A. (1997). *J. Med. Chem.* **40**, 1519–1529.
- Pandeya, S. N., Sriram, D., Nath, G. & Clercq, E. De. (1999). *Pharm. Acta Helv.* **74**, 11–17.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shiba, S. A., El-Khamry, A. A., Shaban, M. & Atia, K. S. (1997). *Pharmazie*, **52**, 189–194.
- Tsou, H. R., Mamuya, N., Johnson, B. D., Reich, M. F., Guber, B. C., Ye, F., Nilakantan, R., Shen, R., Discifani, C., DeBlanc, R., Davis, R., Koehn, F. E., Greenberger, L. M., Wang, Y. F. & Wissner, A. (2001). *J. Med. Chem.* **44**, 2719–2734.
- Yang, X.-H., Wu, M.-H., Sun, S.-F., Ding, M.-W., Xie, J.-L. & Xia, Q.-H. (2008). *J. Heterocycl. Chem.* **45**, 1365–1369.

supplementary materials

Acta Cryst. (2009). E65, o185-o186 [doi:10.1107/S160053680804049X]

3-(2-Aminoethyl)-2-(4-chloroanilino)quinazolin-4(3H)-one methanol 0.75-solvate

X.-H. Yang, X.-B. Chen and S.-X. Zhou

Comment

Quinazoline-4(3H)-one derivatives have broad biological properties. Some of these activities include antimicrobial (Pandeya *et al.*, 1999; Shiba *et al.*, 1997), antidiabetic (Malamas & Millen, 1991), anticonvulsant (Mannschreck *et al.*, 1984), antibacterial (Kung *et al.*, 1999), antifungal (Bartoli *et al.*, 1998), protein tyrosine kinase inhibitors (Palmer *et al.*, 1997), EGFR inhibitors (Tsou *et al.*, 2001) and PDGFR phosphorylation inhibitors (Matsuno *et al.*, 2002). We have recently focused on the synthesis of heterocyclic compounds using an aza-Wittig reaction. We have reported the synthesis of the title compound (Yang *et al.*, 2008). We present here the crystal structure of the title compound, (I) (Fig. 1), which can be used as a precursor for obtaining bioactive molecules.

In the crystal structure, there are two quinazolin-4(3H)-one molecules in the asymmetric unit. The quinazoline heterocycle and the adjacent chlorobenzene ring are not planar, but inclined at 39.83 (1)°. Significant intramolecular N—H···N and O—H···O and intermolecular N—H···O and N—H···N hydrogen bonds contribute strongly to the stability of the molecular configuration (Fig. 2 and Table 1). The crystal structure (Fig. 2) is stabilized by weak intermolecular C—H··· π hydrogen bonds (Table 1) and by π – π stacking interactions with centroid–centroid separations of 3.654 (1), 3.767 (1) and 3.766 (1) Å for Cg1···Cg5ⁱ, Cg2···Cg6ⁱⁱ and Cg3···Cg5ⁱ, respectively, where Cg1, Cg2, Cg3, Cg5 and Cg6 are the centroids of rings N2/C7/N3/C8–C10, C1–C6, C9–C14, C17–C22 and C25–C30, respectively [symmetry codes: (i) 1/2 - x, -1/2 + y, 1/2 - z, (ii) 3/2 - x, -1/2 + y, 1/2 - z].

Experimental

The title compound was prepared according to the literature method (Hu *et al.*, 2006; Yang *et al.*, 2008). Single crystals suitable for X-ray diffraction were obtained from a methanol–dichloromethane (1:1 v/v) solution at room temperature.

Refinement

One of the methanol solvent molecules was found to be disordered over two positions. Their final occupancies were set to be 0.5:0.5. H atoms bonded to C were placed at calculated positions, with C—H distances of 0.97 and 0.93 Å for H atoms bonded to *sp*³ and *sp*² C atoms, respectively. They were refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. H atoms bonded to N and O atoms were found in a difference map and then refined with distance restraints of N—H = 0.86 (2) Å and O—H = 0.82 (2) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ and $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. View of the molecular structure of (I), showing the atom labelling scheme and with displacement ellipsoids drawn at the 50% probability level.




Fig. 2. A partial view of the crystal packing of (I), showing the formation of N—H...N, O—H...O and N—H...O hydrogen-bonds (dashed lines).

3-(2-Aminoethyl)-2-(4-chloroanilino)quinazolin-4(3H)-one methanol 0.75-solvate

Crystal data

$C_{16}H_{15}ClN_4O \cdot 0.75CH_4O$

$M_r = 338.81$

Monoclinic, $P2_1/n$

$a = 13.380$ (3) Å

$b = 12.048$ (2) Å

$c = 21.105$ (4) Å

$\beta = 104.49$ (3)°

$V = 3293.8$ (11) Å³

$Z = 8$

$F_{000} = 1420$

$D_x = 1.366$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 5048 reflections

$\theta = 2.2$ – 25.8 °

$\mu = 0.25$ mm⁻¹

$T = 293$ (2) K

Block, colourless

$0.23 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

φ and ω scans

Absorption correction: none

18726 measured reflections

6451 independent reflections

4529 reflections with $I > 2\sigma(I)$

$R_{int} = 0.018$

$\theta_{max} = 26.0$ °

$\theta_{min} = 2.0$ °

$h = -16 \rightarrow 16$

$k = -13 \rightarrow 14$

$l = -25 \rightarrow 23$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.132$	$w = 1/[\sigma^2(F_o^2) + (0.0651P)^2 + 0.4596P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
6451 reflections	$(\Delta/\sigma)_{\max} = 0.001$
453 parameters	$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
9 restraints	$\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	-0.08984 (15)	0.47979 (16)	0.12395 (9)	0.0528 (5)	
H1	-0.0347	0.4622	0.1063	0.063*	
C2	-0.18874 (15)	0.48069 (16)	0.08437 (10)	0.0556 (5)	
H2	-0.2003	0.4636	0.0402	0.067*	
C3	-0.27035 (14)	0.50686 (16)	0.11023 (10)	0.0534 (5)	
C4	-0.25485 (14)	0.53302 (15)	0.17510 (10)	0.0530 (5)	
H4	-0.3104	0.5514	0.1921	0.064*	
C5	-0.15634 (14)	0.53178 (14)	0.21477 (9)	0.0489 (4)	
H5	-0.1456	0.5491	0.2589	0.059*	
C6	-0.07211 (14)	0.50494 (14)	0.18997 (9)	0.0458 (4)	
C7	0.11714 (13)	0.52077 (14)	0.22075 (9)	0.0462 (4)	
C8	0.30176 (14)	0.52546 (15)	0.26820 (10)	0.0534 (5)	
C9	0.30837 (13)	0.57442 (15)	0.20688 (10)	0.0499 (4)	
C10	0.21855 (13)	0.59011 (14)	0.15719 (9)	0.0481 (4)	
C11	0.22667 (16)	0.63532 (17)	0.09793 (10)	0.0593 (5)	
H11	0.1675	0.6467	0.0644	0.071*	

supplementary materials

C12	0.32126 (18)	0.66313 (18)	0.08869 (12)	0.0688 (6)
H12	0.3256	0.6935	0.0490	0.083*
C13	0.41067 (18)	0.64669 (19)	0.13779 (13)	0.0729 (6)
H13	0.4745	0.6655	0.1309	0.087*
C14	0.40457 (15)	0.60298 (17)	0.19601 (12)	0.0642 (6)
H14	0.4645	0.5918	0.2289	0.077*
C15	0.19411 (16)	0.42811 (17)	0.32874 (10)	0.0621 (5)
H15A	0.2586	0.3890	0.3457	0.074*
H15B	0.1405	0.3729	0.3142	0.074*
C16	0.1693 (2)	0.4950 (2)	0.38362 (11)	0.0721 (6)
H16A	0.1837	0.4505	0.4232	0.087*
H16B	0.2133	0.5600	0.3920	0.087*
C17	1.20193 (15)	0.78133 (15)	0.30649 (10)	0.0524 (5)
H17	1.1943	0.7627	0.2628	0.063*
C18	1.29879 (15)	0.78222 (16)	0.34860 (10)	0.0560 (5)
H18	1.3565	0.7649	0.3335	0.067*
C19	1.30922 (15)	0.80899 (17)	0.41320 (11)	0.0572 (5)
C20	1.22461 (15)	0.83549 (17)	0.43566 (10)	0.0599 (5)
H20	1.2327	0.8534	0.4795	0.072*
C21	1.12771 (15)	0.83567 (16)	0.39363 (10)	0.0572 (5)
H21	1.0706	0.8544	0.4091	0.069*
C22	1.11498 (14)	0.80811 (14)	0.32859 (9)	0.0478 (4)
C23	0.92555 (14)	0.79103 (14)	0.29335 (9)	0.0483 (4)
C24	0.74195 (14)	0.78926 (15)	0.24330 (10)	0.0517 (5)
C25	0.73119 (14)	0.74191 (15)	0.30396 (9)	0.0486 (4)
C26	0.81934 (14)	0.72442 (15)	0.35473 (9)	0.0500 (4)
C27	0.80719 (17)	0.67926 (18)	0.41340 (11)	0.0629 (5)
H27	0.8649	0.6665	0.4477	0.075*
C28	0.71130 (18)	0.65384 (19)	0.42065 (12)	0.0708 (6)
H28	0.7045	0.6235	0.4599	0.085*
C29	0.62350 (16)	0.67269 (18)	0.37022 (12)	0.0667 (6)
H29	0.5585	0.6554	0.3758	0.080*
C30	0.63343 (15)	0.71662 (16)	0.31264 (11)	0.0582 (5)
H30	0.5749	0.7299	0.2789	0.070*
C31	0.85432 (16)	0.88363 (16)	0.18470 (10)	0.0604 (5)
H31A	0.9088	0.9377	0.1997	0.072*
H31B	0.7909	0.9242	0.1668	0.072*
C32	0.87963 (17)	0.81504 (19)	0.13073 (10)	0.0657 (6)
H32A	0.8374	0.7485	0.1238	0.079*
H32B	0.8637	0.8574	0.0903	0.079*
C33	0.5966 (3)	0.8449 (4)	0.02991 (16)	0.1534 (17)
H33A	0.6527	0.8967	0.0355	0.230*
H33B	0.5503	0.8552	-0.0124	0.230*
H33C	0.6232	0.7706	0.0335	0.230*
Cl1	-0.39482 (4)	0.50595 (6)	0.05930 (3)	0.0837 (2)
Cl2	1.43110 (4)	0.81046 (7)	0.46778 (3)	0.0904 (2)
N1	0.02479 (12)	0.49709 (14)	0.23440 (8)	0.0541 (4)
H1A	0.0250 (17)	0.5023 (17)	0.2759 (5)	0.065*
N2	0.12216 (11)	0.56474 (13)	0.16588 (7)	0.0486 (4)

N3	0.20303 (11)	0.49523 (12)	0.27173 (7)	0.0486 (4)	
H3A	0.5847 (15)	0.8370 (17)	0.1097 (7)	0.215*	
N4	0.06204 (17)	0.52954 (17)	0.36770 (9)	0.0716 (5)	
H4A	0.0474 (18)	0.5956 (11)	0.3769 (12)	0.086*	
H4B	0.0261 (17)	0.4888 (18)	0.3870 (11)	0.086*	
N5	1.01963 (12)	0.81229 (14)	0.28200 (8)	0.0553 (4)	
H5A	1.0179 (16)	0.8063 (17)	0.2410 (5)	0.066*	
N6	0.91717 (11)	0.74695 (13)	0.34781 (8)	0.0524 (4)	
N7	0.84205 (12)	0.81837 (12)	0.24149 (7)	0.0490 (4)	
N8	0.98838 (15)	0.78405 (16)	0.14741 (9)	0.0662 (5)	
H8A	1.0023 (18)	0.7195 (11)	0.1344 (11)	0.079*	
H8B	1.0213 (16)	0.8306 (15)	0.1300 (11)	0.079*	
O1	0.37637 (11)	0.50823 (12)	0.31482 (8)	0.0725 (4)	
O2	0.66919 (11)	0.80647 (12)	0.19558 (7)	0.0678 (4)	
O3	0.5479 (2)	0.8615 (2)	0.07449 (11)	0.1432 (10)	
C34	0.5063 (2)	0.4916 (2)	0.50408 (11)	0.131 (2)	0.50
H34A	0.5228	0.5678	0.4981	0.197*	0.50
H34B	0.5664	0.4465	0.5069	0.197*	0.50
H34C	0.4840	0.4844	0.5437	0.197*	0.50
O4	0.4296 (5)	0.4506 (6)	0.4520 (3)	0.168 (2)	0.50
H4C	0.4324	0.4827	0.4183	0.253*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0460 (10)	0.0627 (12)	0.0515 (12)	0.0002 (8)	0.0153 (9)	-0.0065 (9)
C2	0.0538 (11)	0.0646 (12)	0.0466 (11)	-0.0023 (9)	0.0094 (9)	-0.0065 (9)
C3	0.0412 (10)	0.0549 (11)	0.0620 (13)	-0.0048 (8)	0.0086 (9)	0.0037 (9)
C4	0.0465 (10)	0.0491 (10)	0.0682 (13)	-0.0022 (8)	0.0232 (9)	-0.0019 (9)
C5	0.0523 (11)	0.0473 (10)	0.0508 (11)	-0.0065 (8)	0.0197 (9)	-0.0058 (8)
C6	0.0443 (10)	0.0450 (10)	0.0478 (11)	-0.0036 (7)	0.0111 (8)	-0.0012 (8)
C7	0.0424 (10)	0.0478 (10)	0.0461 (11)	0.0011 (7)	0.0066 (8)	-0.0042 (8)
C8	0.0464 (11)	0.0457 (10)	0.0627 (13)	0.0047 (8)	0.0039 (9)	-0.0043 (9)
C9	0.0450 (10)	0.0423 (10)	0.0615 (12)	0.0009 (8)	0.0115 (9)	-0.0078 (8)
C10	0.0462 (10)	0.0442 (10)	0.0550 (11)	0.0008 (8)	0.0149 (8)	-0.0072 (8)
C11	0.0618 (12)	0.0634 (12)	0.0552 (12)	-0.0011 (10)	0.0191 (10)	-0.0021 (10)
C12	0.0767 (15)	0.0650 (13)	0.0743 (15)	-0.0069 (11)	0.0368 (13)	-0.0032 (11)
C13	0.0625 (14)	0.0639 (13)	0.103 (2)	-0.0068 (11)	0.0408 (14)	-0.0064 (13)
C14	0.0447 (11)	0.0554 (12)	0.0912 (17)	-0.0010 (9)	0.0145 (11)	-0.0084 (11)
C15	0.0648 (13)	0.0555 (12)	0.0608 (13)	0.0046 (9)	0.0061 (10)	0.0142 (10)
C16	0.0844 (16)	0.0790 (15)	0.0473 (12)	-0.0093 (12)	0.0059 (11)	0.0110 (11)
C17	0.0589 (12)	0.0484 (10)	0.0532 (11)	-0.0026 (8)	0.0200 (9)	-0.0015 (8)
C18	0.0488 (11)	0.0569 (12)	0.0660 (13)	0.0018 (8)	0.0214 (10)	0.0047 (10)
C19	0.0486 (11)	0.0600 (12)	0.0617 (13)	-0.0071 (9)	0.0111 (9)	0.0072 (10)
C20	0.0562 (12)	0.0727 (13)	0.0518 (12)	-0.0147 (10)	0.0156 (10)	-0.0056 (10)
C21	0.0522 (11)	0.0644 (12)	0.0588 (13)	-0.0074 (9)	0.0208 (10)	-0.0108 (10)
C22	0.0484 (10)	0.0424 (10)	0.0531 (11)	-0.0058 (8)	0.0139 (9)	0.0006 (8)
C23	0.0477 (10)	0.0442 (10)	0.0516 (11)	-0.0012 (8)	0.0097 (9)	-0.0036 (8)

supplementary materials

C24	0.0493 (11)	0.0482 (10)	0.0549 (12)	0.0033 (8)	0.0078 (9)	-0.0086 (9)
C25	0.0479 (10)	0.0437 (10)	0.0534 (11)	-0.0010 (8)	0.0114 (9)	-0.0076 (8)
C26	0.0483 (10)	0.0466 (10)	0.0550 (11)	-0.0017 (8)	0.0128 (9)	-0.0042 (8)
C27	0.0569 (12)	0.0735 (14)	0.0566 (13)	-0.0041 (10)	0.0113 (10)	0.0064 (10)
C28	0.0678 (14)	0.0782 (15)	0.0711 (15)	-0.0055 (11)	0.0261 (12)	0.0073 (12)
C29	0.0541 (12)	0.0679 (14)	0.0820 (16)	-0.0083 (10)	0.0242 (12)	-0.0017 (11)
C30	0.0473 (11)	0.0568 (12)	0.0678 (14)	-0.0025 (9)	0.0092 (10)	-0.0080 (10)
C31	0.0619 (12)	0.0540 (12)	0.0634 (13)	0.0021 (9)	0.0124 (10)	0.0117 (10)
C32	0.0673 (14)	0.0764 (14)	0.0520 (12)	-0.0076 (11)	0.0124 (10)	0.0101 (10)
C33	0.107 (3)	0.280 (6)	0.075 (2)	0.029 (3)	0.025 (2)	0.000 (3)
Cl1	0.0437 (3)	0.1160 (5)	0.0841 (4)	-0.0049 (3)	0.0024 (3)	0.0082 (3)
Cl2	0.0523 (3)	0.1351 (6)	0.0773 (4)	-0.0083 (3)	0.0042 (3)	0.0178 (4)
N1	0.0467 (9)	0.0715 (11)	0.0437 (9)	-0.0048 (7)	0.0104 (8)	-0.0004 (8)
N2	0.0438 (8)	0.0563 (9)	0.0450 (9)	0.0024 (7)	0.0098 (7)	-0.0009 (7)
N3	0.0474 (9)	0.0474 (8)	0.0481 (9)	0.0028 (6)	0.0066 (7)	0.0024 (7)
N4	0.0872 (14)	0.0732 (13)	0.0557 (11)	0.0059 (11)	0.0204 (10)	0.0053 (10)
N5	0.0497 (9)	0.0676 (10)	0.0486 (9)	-0.0061 (7)	0.0122 (8)	-0.0009 (8)
N6	0.0461 (9)	0.0573 (9)	0.0524 (10)	-0.0024 (7)	0.0095 (7)	0.0008 (8)
N7	0.0504 (9)	0.0454 (8)	0.0503 (9)	0.0015 (7)	0.0112 (7)	0.0008 (7)
N8	0.0703 (12)	0.0699 (12)	0.0611 (12)	0.0019 (10)	0.0214 (9)	0.0070 (9)
O1	0.0518 (8)	0.0808 (10)	0.0731 (10)	0.0075 (7)	-0.0065 (8)	0.0072 (8)
O2	0.0562 (8)	0.0828 (10)	0.0571 (9)	0.0043 (7)	0.0004 (7)	0.0008 (7)
O3	0.153 (2)	0.196 (3)	0.0823 (15)	0.088 (2)	0.0324 (15)	0.0356 (16)
C34	0.187 (7)	0.107 (4)	0.098 (4)	-0.014 (4)	0.033 (4)	0.004 (3)
O4	0.170 (6)	0.180 (6)	0.120 (4)	0.016 (4)	-0.028 (4)	0.018 (4)

Geometric parameters (Å, °)

C1—C2	1.377 (3)	C20—H20	0.9300
C1—C6	1.387 (3)	C21—C22	1.381 (3)
C1—H1	0.9300	C21—H21	0.9300
C2—C3	1.374 (3)	C22—N5	1.403 (2)
C2—H2	0.9300	C23—N6	1.296 (2)
C3—C4	1.369 (3)	C23—N5	1.363 (2)
C3—Cl1	1.742 (2)	C23—N7	1.394 (2)
C4—C5	1.373 (3)	C24—O2	1.231 (2)
C4—H4	0.9300	C24—N7	1.394 (2)
C5—C6	1.394 (3)	C24—C25	1.441 (3)
C5—H5	0.9300	C25—C26	1.397 (3)
C6—N1	1.400 (2)	C25—C30	1.399 (3)
C7—N2	1.290 (2)	C26—N6	1.380 (2)
C7—N1	1.367 (2)	C26—C27	1.399 (3)
C7—N3	1.398 (2)	C27—C28	1.364 (3)
C8—O1	1.231 (2)	C27—H27	0.9300
C8—N3	1.390 (2)	C28—C29	1.392 (3)
C8—C9	1.445 (3)	C28—H28	0.9300
C9—C10	1.396 (3)	C29—C30	1.362 (3)
C9—C14	1.406 (3)	C29—H29	0.9300
C10—N2	1.381 (2)	C30—H30	0.9300

C10—C11	1.393 (3)	C31—N7	1.477 (2)
C11—C12	1.369 (3)	C31—C32	1.513 (3)
C11—H11	0.9300	C31—H31A	0.9700
C12—C13	1.387 (3)	C31—H31B	0.9700
C12—H12	0.9300	C32—N8	1.457 (3)
C13—C14	1.358 (3)	C32—H32A	0.9700
C13—H13	0.9300	C32—H32B	0.9700
C14—H14	0.9300	C33—O3	1.287 (4)
C15—N3	1.479 (2)	C33—H33A	0.9600
C15—C16	1.514 (3)	C33—H33B	0.9600
C15—H15A	0.9700	C33—H33C	0.9600
C15—H15B	0.9700	N1—H1A	0.877 (9)
C16—N4	1.451 (3)	N4—H4A	0.853 (10)
C16—H16A	0.9700	N4—H4B	0.858 (10)
C16—H16B	0.9700	N5—H5A	0.863 (9)
C17—C18	1.376 (3)	N8—H8A	0.861 (10)
C17—C22	1.395 (3)	N8—H8B	0.851 (10)
C17—H17	0.9300	O3—H3A	0.835 (10)
C18—C19	1.374 (3)	C34—O4	1.392 (6)
C18—H18	0.9300	C34—H34A	0.9600
C19—C20	1.370 (3)	C34—H34B	0.9600
C19—C12	1.745 (2)	C34—H34C	0.9600
C20—C21	1.376 (3)	O4—H4C	0.8200
C2—C1—C6	120.30 (18)	N6—C23—N5	121.41 (17)
C2—C1—H1	119.8	N6—C23—N7	124.26 (17)
C6—C1—H1	119.8	N5—C23—N7	114.31 (17)
C3—C2—C1	119.97 (18)	O2—C24—N7	120.26 (19)
C3—C2—H2	120.0	O2—C24—C25	124.01 (18)
C1—C2—H2	120.0	N7—C24—C25	115.72 (17)
C4—C3—C2	120.89 (18)	C26—C25—C30	120.28 (18)
C4—C3—C11	120.05 (15)	C26—C25—C24	119.23 (17)
C2—C3—C11	119.06 (16)	C30—C25—C24	120.47 (18)
C3—C4—C5	119.29 (17)	N6—C26—C25	122.04 (18)
C3—C4—H4	120.4	N6—C26—C27	119.58 (18)
C5—C4—H4	120.4	C25—C26—C27	118.36 (17)
C4—C5—C6	121.07 (17)	C28—C27—C26	120.5 (2)
C4—C5—H5	119.5	C28—C27—H27	119.8
C6—C5—H5	119.5	C26—C27—H27	119.8
C1—C6—C5	118.47 (17)	C27—C28—C29	121.0 (2)
C1—C6—N1	123.70 (17)	C27—C28—H28	119.5
C5—C6—N1	117.66 (16)	C29—C28—H28	119.5
N2—C7—N1	121.83 (16)	C30—C29—C28	119.57 (19)
N2—C7—N3	124.32 (16)	C30—C29—H29	120.2
N1—C7—N3	113.83 (16)	C28—C29—H29	120.2
O1—C8—N3	120.28 (19)	C29—C30—C25	120.30 (19)
O1—C8—C9	124.43 (19)	C29—C30—C14	101.64 (14)
N3—C8—C9	115.28 (17)	C25—C30—C14	129.96 (14)
C10—C9—C14	119.71 (19)	C29—C30—H30	119.8
C10—C9—C8	119.60 (16)	C25—C30—H30	119.8

supplementary materials

C14—C9—C8	120.66 (18)	N7—C31—C32	114.31 (16)
N2—C10—C11	119.40 (17)	N7—C31—H31A	108.7
N2—C10—C9	121.80 (17)	C32—C31—H31A	108.7
C11—C10—C9	118.78 (17)	N7—C31—H31B	108.7
C12—C11—C10	120.4 (2)	C32—C31—H31B	108.7
C12—C11—H11	119.8	H31A—C31—H31B	107.6
C10—C11—H11	119.8	N8—C32—C31	110.86 (18)
C11—C12—C13	120.9 (2)	N8—C32—H32A	109.5
C11—C12—H12	119.5	C31—C32—H32A	109.5
C13—C12—H12	119.5	N8—C32—H32B	109.5
C14—C13—C12	119.7 (2)	C31—C32—H32B	109.5
C14—C13—H13	120.1	H32A—C32—H32B	108.1
C12—C13—H13	120.1	O3—C33—H33A	109.5
C13—C14—C9	120.5 (2)	O3—C33—H33B	109.5
C13—C14—C30	103.46 (15)	H33A—C33—H33B	109.5
C9—C14—C30	128.66 (15)	O3—C33—H33C	109.5
C13—C14—H14	119.8	H33A—C33—H33C	109.5
C9—C14—H14	119.8	H33B—C33—H33C	109.5
N3—C15—C16	114.16 (17)	C7—N1—C6	125.38 (16)
N3—C15—H15A	108.7	C7—N1—H1A	114.7 (15)
C16—C15—H15A	108.7	C6—N1—H1A	115.8 (15)
N3—C15—H15B	108.7	C7—N2—C10	118.03 (16)
C16—C15—H15B	108.7	C8—N3—C7	120.67 (16)
H15A—C15—H15B	107.6	C8—N3—C15	117.26 (16)
N4—C16—C15	111.68 (19)	C7—N3—C15	121.90 (15)
N4—C16—H16A	109.3	C16—N4—H4A	119.0 (17)
C15—C16—H16A	109.3	C16—N4—H4B	111.4 (17)
N4—C16—H16B	109.3	H4A—N4—H4B	104 (2)
C15—C16—H16B	109.3	C23—N5—C22	126.00 (17)
H16A—C16—H16B	107.9	C23—N5—H5A	111.5 (15)
C18—C17—C22	120.78 (18)	C22—N5—H5A	119.4 (15)
C18—C17—H17	119.6	C23—N6—C26	117.98 (16)
C22—C17—H17	119.6	C23—N7—C24	120.39 (16)
C19—C18—C17	119.23 (18)	C23—N7—C31	122.22 (15)
C19—C18—H18	120.4	C24—N7—C31	117.31 (16)
C17—C18—H18	120.4	C32—N8—H8A	115.9 (17)
C20—C19—C18	120.71 (19)	C32—N8—H8B	108.7 (16)
C20—C19—C12	119.06 (17)	H8A—N8—H8B	106 (2)
C18—C19—C12	120.23 (16)	C24—O2—O3	164.88 (15)
C19—C20—C21	120.29 (19)	C33—O3—O2	110.7 (2)
C19—C20—H20	119.9	C33—O3—H3A	107.5 (17)
C21—C20—H20	119.9	O4—C34—H34A	112.5
C20—C21—C22	120.17 (18)	O4—C34—H34B	106.7
C20—C21—H21	119.9	H34A—C34—H34B	109.5
C22—C21—H21	119.9	O4—C34—H34C	109.1
C21—C22—C17	118.83 (18)	H34A—C34—H34C	109.5
C21—C22—N5	123.52 (17)	H34B—C34—H34C	109.5
C17—C22—N5	117.52 (17)		
C6—C1—C2—C3	0.2 (3)	C28—C29—C30—C25	0.5 (3)

C1—C2—C3—C4	0.5 (3)	C28—C29—C30—C14	152.10 (18)
C1—C2—C3—C11	-179.32 (15)	C26—C25—C30—C29	-1.3 (3)
C2—C3—C4—C5	-0.7 (3)	C24—C25—C30—C29	-179.48 (18)
C11—C3—C4—C5	179.06 (14)	C26—C25—C30—C14	-143.90 (15)
C3—C4—C5—C6	0.3 (3)	C24—C25—C30—C14	37.9 (2)
C2—C1—C6—C5	-0.5 (3)	C13—C14—C30—C29	178.95 (18)
C2—C1—C6—N1	174.61 (17)	C9—C14—C30—C29	29.7 (2)
C4—C5—C6—C1	0.3 (3)	C13—C14—C30—C25	-33.4 (2)
C4—C5—C6—N1	-175.16 (16)	C9—C14—C30—C25	177.3 (2)
O1—C8—C9—C10	179.95 (17)	N7—C31—C32—N8	-78.3 (2)
N3—C8—C9—C10	1.3 (2)	N2—C7—N1—C6	7.5 (3)
O1—C8—C9—C14	2.1 (3)	N3—C7—N1—C6	-173.77 (16)
N3—C8—C9—C14	-176.57 (16)	C1—C6—N1—C7	35.7 (3)
C14—C9—C10—N2	-179.09 (16)	C5—C6—N1—C7	-149.14 (18)
C8—C9—C10—N2	3.0 (3)	N1—C7—N2—C10	177.40 (16)
C14—C9—C10—C11	-0.8 (3)	N3—C7—N2—C10	-1.2 (3)
C8—C9—C10—C11	-178.69 (17)	C11—C10—N2—C7	178.56 (17)
N2—C10—C11—C12	178.72 (17)	C9—C10—N2—C7	-3.2 (3)
C9—C10—C11—C12	0.4 (3)	O1—C8—N3—C7	175.92 (16)
C10—C11—C12—C13	0.2 (3)	C9—C8—N3—C7	-5.4 (2)
C11—C12—C13—C14	-0.3 (3)	O1—C8—N3—C15	-8.7 (3)
C12—C13—C14—C9	-0.1 (3)	C9—C8—N3—C15	170.03 (16)
C12—C13—C14—C30	-152.49 (18)	N2—C7—N3—C8	5.7 (3)
C10—C9—C14—C13	0.7 (3)	N1—C7—N3—C8	-172.99 (15)
C8—C9—C14—C13	178.53 (18)	N2—C7—N3—C15	-169.51 (18)
C10—C9—C14—C30	145.43 (15)	N1—C7—N3—C15	11.8 (2)
C8—C9—C14—C30	-36.7 (2)	C16—C15—N3—C8	97.6 (2)
N3—C15—C16—N4	75.0 (2)	C16—C15—N3—C7	-87.0 (2)
C22—C17—C18—C19	0.5 (3)	N6—C23—N5—C22	-11.2 (3)
C17—C18—C19—C20	-0.5 (3)	N7—C23—N5—C22	170.10 (16)
C17—C18—C19—C12	179.84 (14)	C21—C22—N5—C23	-33.2 (3)
C18—C19—C20—C21	0.0 (3)	C17—C22—N5—C23	151.00 (17)
C12—C19—C20—C21	179.59 (15)	N5—C23—N6—C26	-177.79 (16)
C19—C20—C21—C22	0.6 (3)	N7—C23—N6—C26	0.8 (3)
C20—C21—C22—C17	-0.6 (3)	C25—C26—N6—C23	3.6 (3)
C20—C21—C22—N5	-176.36 (18)	C27—C26—N6—C23	-178.51 (18)
C18—C17—C22—C21	0.0 (3)	N6—C23—N7—C24	-6.1 (3)
C18—C17—C22—N5	176.04 (16)	N5—C23—N7—C24	172.51 (15)
O2—C24—C25—C26	179.05 (18)	N6—C23—N7—C31	170.51 (17)
N7—C24—C25—C26	-2.5 (2)	N5—C23—N7—C31	-10.8 (2)
O2—C24—C25—C30	-2.7 (3)	O2—C24—N7—C23	-174.86 (16)
N7—C24—C25—C30	175.71 (16)	C25—C24—N7—C23	6.6 (2)
C30—C25—C26—N6	179.15 (16)	O2—C24—N7—C31	8.3 (3)
C24—C25—C26—N6	-2.6 (3)	C25—C24—N7—C31	-170.18 (15)
C30—C25—C26—C27	1.3 (3)	C32—C31—N7—C23	86.7 (2)
C24—C25—C26—C27	179.46 (17)	C32—C31—N7—C24	-96.5 (2)
N6—C26—C27—C28	-178.38 (19)	N7—C24—O2—O3	5.4 (6)
C25—C26—C27—C28	-0.4 (3)	C25—C24—O2—O3	-176.2 (4)
C26—C27—C28—C29	-0.4 (3)	C24—O2—O3—C33	30.1 (7)

supplementary materials

C27—C28—C29—C30

0.4 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N5—H5A \cdots N8	0.863 (9)	1.933 (11)	2.787 (3)	170 (2)
O4—H4C \cdots O1	0.82	2.15	2.889 (6)	151
O3—H3A \cdots O2	0.835 (10)	1.917 (11)	2.744 (3)	170 (2)
N1—H1A \cdots N4	0.877 (9)	1.904 (11)	2.760 (3)	164 (2)
N8—H8B \cdots O4 ⁱ	0.851 (10)	2.467 (14)	3.284 (8)	161 (2)
N8—H8A \cdots N2 ⁱⁱ	0.861 (10)	2.440 (17)	3.160 (2)	142 (2)
N4—H4B \cdots O3 ⁱⁱⁱ	0.858 (10)	2.097 (11)	2.941 (3)	168 (2)
N4—H4A \cdots N6 ^{iv}	0.853 (10)	2.491 (17)	3.223 (3)	144 (2)
C5—H5 \cdots Cg4 ^{iv}	0.93	2.81	3.375 (2)	120
C17—H17 \cdots Cg1 ⁱⁱ	0.93	2.84	3.459 (2)	125
C32—H32A \cdots Cg2 ⁱⁱ	0.97	2.98	3.824 (3)	146

Symmetry codes: (i) $-x+3/2, y+1/2, -z+1/2$; (ii) $x+1, y, z$; (iii) $-x+1/2, y-1/2, -z+1/2$; (iv) $x-1, y, z$.

Fig. 1




Fig. 2

