

1-(3-Chlorophenyl)-3-(2,6-dichlorobenzoyl)thiourea

M. Khawar Rauf,^{a*} Michael Bolte^b and Abdur Rauf^c

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan,

^bInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany, and ^cDepartment of Chemistry, Islamia University of Bahawalpur, Pakistan

Correspondence e-mail: khawar_rauf@hotmail.com


Received 1 December 2008; accepted 20 December 2008

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.094; data-to-parameter ratio = 25.5.

The structure of the title compound, $\text{C}_{14}\text{H}_9\text{Cl}_3\text{N}_2\text{OS}$, is composed of discrete molecules with bond lengths and angles quite typical for thiourea compounds of this class. The plane containing the thiocarbonyl and carbonyl groups subtends dihedral angles of 48.19 (3) and 87.51 (3)° with the planes formed by the 3-chloro and 2,6-dichlorophenyl rings, respectively; the dihedral angle between the two benzene ring planes is 45.32 (3)°. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond stabilizes the molecular conformation and the molecules form intermolecular $\text{N}-\text{H}\cdots\text{S}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, generating a sheet along the a axis.

Related literature

For related structures, see: Khawar Rauf *et al.*, (2006*a,b*; 2007). For a description of the Cambridge structural Database, see: Allen (2002).



Experimental

Crystal data

$\text{C}_{14}\text{H}_9\text{Cl}_3\text{N}_2\text{OS}$

$M_r = 359.64$

Monoclinic, $P2_1/c$
 $a = 10.6589$ (5) Å
 $b = 11.2114$ (5) Å
 $c = 13.2919$ (6) Å
 $\beta = 99.942$ (3)°
 $V = 1564.55$ (12) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.72$ mm⁻¹
 $T = 173$ (2) K
 $0.47 \times 0.47 \times 0.45$ mm

Data collection

Stoe IPDS-II two-circle diffractometer
 Absorption correction: multi-scan (*MULABS*; Spek, 2003; Blessing, 1995)
 $T_{\min} = 0.729$, $T_{\max} = 0.739$

39690 measured reflections
 5066 independent reflections
 4674 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.094$
 $S = 1.06$
 5066 reflections
 199 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.65$ e Å⁻³
 $\Delta\rho_{\min} = -0.66$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2}\cdots\text{O1}$	0.82 (2)	2.07 (2)	2.7190 (13)	136.0 (18)
$\text{N2}-\text{H2}\cdots\text{O1}^{\text{i}}$	0.82 (2)	2.37 (2)	3.0749 (14)	145.5 (18)
$\text{N1}-\text{H1}\cdots\text{S1}^{\text{ii}}$	0.86 (2)	2.47 (2)	3.2974 (10)	163.6 (18)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003) and *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* and *SHELXL97*.

MKR is grateful to the HEC-Pakistan for financial support for the PhD program under scholarship No. [ILC-0363104].

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2127).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Khawar Rauf, M., Badshah, A. & Bolte, M. (2006*a*). *Acta Cryst.* **E62**, o3859–o3861.
 Khawar Rauf, M., Badshah, A. & Bolte, M. (2006*b*). *Acta Cryst.* **E62**, o4296–o4298.
 Khawar Rauf, M., Badshah, A. & Bolte, M. (2007). *Acta Cryst.* **E63**, o2665–o2666.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2009). E65, o234 [doi:10.1107/S1600536808043444]

1-(3-Chlorophenyl)-3-(2,6-dichlorobenzoyl)thiourea

M. Khawar Rauf, M. Bolte and A. Rauf

Comment

The background to this study has been set out in our previous work on the structural chemistry of N,N'-disubstituted thioureas (Khawar Rauf et al., 2006a, 2007). Herein, as a continuation of these studies, the structure of the title compound, (I), is described.

In the structure of the title compound (Fig. 1), bond lengths and bond angles can be regarded as typical for N,N'-disubstituted thiourea compounds as found in the Cambridge Structural Database v5.28 (Allen, 2002) and some related structures (Khawar Rauf et al., 2006b). The molecule exists in the thione form with typical thiourea C—S and C—O bonds, as well as shortened C—N bonds. The thiocarbonyl and carbonyl groups are almost coplanar. The molecule features an intramolecular N—H···O hydrogen bond in the crystal structure. The molecules lying about inversion centers associate via N—H···S intermolecular hydrogen bonds to form dimers on one side and a similar association via N—H···O hydrogen bonding on the other side thus result in a sheet of molecules of (I) along the *a*-axis (Table 1; Fig. 2).

Experimental

Freshly prepared and steam distilled 2,6-dichlorobenzoyl isothiocyanate (2.32 g, 10 mmol) was stirred in acetone (30 ml) for 20 minutes. Neat 3-chloroaniline (1.27 g, 10 mmol) was then added and the resulting mixture was stirred for 1 h. The reaction mixture was then poured into acidified (pH 4) water (approx. 300 ml) and stirred well. The solid product was separated, washed with deionized water and purified by recrystallization from methanol/1,1-dichloromethane (1:10 v/v) to give fine crystals of (I), with an overall yield of 85%.

Refinement

Hydrogen atoms bonded to C were included in calculated positions and refined as riding on their parent C atom with C—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The H atoms bonded to N were freely refined.

Figures




Fig. 1. Molecular structure of (I) showing atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A view of the unit cell of (I) showing hydrogen bonds as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

1-(3-Chlorophenyl)-3-(2,6-dichlorobenzoyl)thiourea

Crystal data

$C_{14}H_9Cl_3N_2OS$	$F_{000} = 728$
$M_r = 359.64$	$D_x = 1.527 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 10.6589 (5) \text{ \AA}$	Cell parameters from 37695 reflections
$b = 11.2114 (5) \text{ \AA}$	$\theta = 3.7\text{--}31.2^\circ$
$c = 13.2919 (6) \text{ \AA}$	$\mu = 0.72 \text{ mm}^{-1}$
$\beta = 99.942 (3)^\circ$	$T = 173 (2) \text{ K}$
$V = 1564.55 (12) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.47 \times 0.47 \times 0.45 \text{ mm}$

Data collection

Stoe IPDS-II two-circle diffractometer	5066 independent reflections
Radiation source: fine-focus sealed tube	4674 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.046$
$T = 173(2) \text{ K}$	$\theta_{\text{max}} = 31.3^\circ$
ω scans	$\theta_{\text{min}} = 3.6^\circ$
Absorption correction: multi-scan (MULABS; Spek, 2003; Blessing, 1995)	$h = -15 \rightarrow 15$
$T_{\text{min}} = 0.729$, $T_{\text{max}} = 0.739$	$k = -16 \rightarrow 16$
39690 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.0404P)^2 + 0.9365P]$
$wR(F^2) = 0.094$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} = 0.001$
5066 reflections	$\Delta\rho_{\text{max}} = 0.65 \text{ e \AA}^{-3}$
199 parameters	$\Delta\rho_{\text{min}} = -0.66 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0111 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.09625 (3)	0.60027 (3)	0.62517 (2)	0.02474 (9)
C11	0.19574 (4)	0.61905 (4)	0.23352 (3)	0.03999 (11)
C12	0.22323 (4)	0.18815 (4)	0.41908 (3)	0.03969 (11)
C13	0.33122 (7)	0.61484 (7)	0.99688 (3)	0.0735 (2)
C1	0.27158 (11)	0.45474 (11)	0.41547 (8)	0.0182 (2)
O1	0.38737 (8)	0.46303 (10)	0.43625 (7)	0.0285 (2)
N1	0.19003 (9)	0.49493 (10)	0.47738 (8)	0.01915 (19)
H1	0.1105 (19)	0.4841 (18)	0.4559 (15)	0.030 (5)*
C2	0.21866 (10)	0.55197 (10)	0.57210 (8)	0.0174 (2)
N2	0.34238 (9)	0.56439 (10)	0.61243 (8)	0.01918 (19)
H2	0.394 (2)	0.5442 (19)	0.5770 (15)	0.032 (5)*
C11	0.20320 (10)	0.39859 (11)	0.31773 (8)	0.0179 (2)
C12	0.16489 (13)	0.46736 (13)	0.23029 (10)	0.0251 (2)
C13	0.09985 (17)	0.41613 (18)	0.14040 (11)	0.0394 (4)
H13	0.0745	0.4639	0.0814	0.047*
C14	0.07264 (17)	0.29526 (19)	0.13789 (12)	0.0421 (4)
H14	0.0275	0.2606	0.0770	0.051*
C15	0.11043 (15)	0.22420 (15)	0.22299 (12)	0.0337 (3)
H15	0.0923	0.1412	0.2208	0.040*
C16	0.17558 (12)	0.27683 (12)	0.31200 (10)	0.0232 (2)
C21	0.39003 (11)	0.62132 (11)	0.70818 (9)	0.0195 (2)
C22	0.34464 (15)	0.58955 (14)	0.79662 (10)	0.0289 (3)
H22	0.2822	0.5288	0.7953	0.035*
C23	0.39279 (16)	0.64880 (16)	0.88702 (10)	0.0340 (3)
C24	0.48645 (15)	0.73570 (15)	0.89168 (11)	0.0332 (3)
H24	0.5180	0.7752	0.9541	0.040*
C25	0.53305 (14)	0.76367 (15)	0.80313 (11)	0.0318 (3)
H25	0.5984	0.8218	0.8053	0.038*
C26	0.48479 (12)	0.70721 (13)	0.71106 (10)	0.0246 (2)
H26	0.5164	0.7273	0.6507	0.030*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.01676 (13)	0.03383 (17)	0.02412 (15)	-0.00159 (11)	0.00495 (10)	-0.01452 (12)
C11	0.0485 (2)	0.02974 (18)	0.0415 (2)	-0.00112 (15)	0.00718 (16)	0.00931 (14)
C12	0.0507 (2)	0.02793 (17)	0.0402 (2)	0.00109 (15)	0.00724 (16)	0.00754 (14)
C13	0.1095 (5)	0.0945 (5)	0.02015 (18)	-0.0585 (4)	0.0216 (2)	-0.0165 (2)
C1	0.0160 (4)	0.0228 (5)	0.0156 (4)	0.0010 (4)	0.0018 (4)	-0.0044 (4)
O1	0.0144 (4)	0.0461 (6)	0.0246 (4)	0.0004 (4)	0.0023 (3)	-0.0134 (4)
N1	0.0133 (4)	0.0275 (5)	0.0163 (4)	-0.0008 (3)	0.0016 (3)	-0.0081 (4)
C2	0.0171 (5)	0.0197 (5)	0.0152 (4)	-0.0009 (4)	0.0018 (4)	-0.0037 (4)
N2	0.0153 (4)	0.0266 (5)	0.0152 (4)	-0.0008 (4)	0.0015 (3)	-0.0061 (4)
C11	0.0160 (4)	0.0231 (5)	0.0147 (4)	-0.0007 (4)	0.0030 (4)	-0.0051 (4)
C12	0.0247 (6)	0.0317 (6)	0.0185 (5)	-0.0009 (5)	0.0023 (4)	0.0000 (4)
C13	0.0408 (8)	0.0571 (10)	0.0173 (6)	-0.0037 (7)	-0.0037 (5)	-0.0014 (6)
C14	0.0391 (8)	0.0612 (11)	0.0243 (6)	-0.0127 (8)	0.0002 (6)	-0.0189 (7)
C15	0.0315 (7)	0.0355 (7)	0.0350 (7)	-0.0098 (6)	0.0080 (5)	-0.0185 (6)
C16	0.0225 (5)	0.0246 (6)	0.0236 (5)	-0.0020 (4)	0.0070 (4)	-0.0059 (4)
C21	0.0185 (5)	0.0235 (5)	0.0151 (4)	-0.0003 (4)	-0.0005 (4)	-0.0049 (4)
C22	0.0358 (7)	0.0330 (7)	0.0168 (5)	-0.0123 (5)	0.0015 (5)	-0.0041 (5)
C23	0.0446 (8)	0.0414 (8)	0.0156 (5)	-0.0123 (7)	0.0041 (5)	-0.0064 (5)
C24	0.0341 (7)	0.0410 (8)	0.0222 (6)	-0.0079 (6)	-0.0016 (5)	-0.0123 (5)
C25	0.0265 (6)	0.0382 (8)	0.0300 (6)	-0.0108 (5)	0.0027 (5)	-0.0129 (6)
C26	0.0206 (5)	0.0305 (6)	0.0229 (5)	-0.0044 (5)	0.0038 (4)	-0.0076 (5)

Geometric parameters (\AA , $^\circ$)

S1—C2	1.6766 (12)	C13—H13	0.9500
C11—C12	1.7313 (15)	C14—C15	1.385 (3)
C12—C16	1.7386 (14)	C14—H14	0.9500
C13—C23	1.7433 (15)	C15—C16	1.3951 (18)
C1—O1	1.2209 (14)	C15—H15	0.9500
C1—N1	1.3721 (14)	C21—C26	1.3912 (18)
C1—C11	1.5128 (15)	C21—C22	1.3925 (18)
N1—C2	1.3980 (14)	C22—C23	1.3912 (18)
N1—H1	0.86 (2)	C22—H22	0.9500
C2—N2	1.3423 (14)	C23—C24	1.389 (2)
N2—C21	1.4360 (14)	C24—C25	1.390 (2)
N2—H2	0.82 (2)	C24—H24	0.9500
C11—C12	1.3957 (17)	C25—C26	1.3947 (17)
C11—C16	1.3957 (17)	C25—H25	0.9500
C12—C13	1.3971 (19)	C26—H26	0.9500
C13—C14	1.385 (3)		
O1—C1—N1	124.03 (10)	C14—C15—C16	118.77 (14)
O1—C1—C11	122.98 (10)	C14—C15—H15	120.6
N1—C1—C11	112.99 (9)	C16—C15—H15	120.6
C1—N1—C2	128.90 (10)	C15—C16—C11	121.96 (13)

C1—N1—H1	116.6 (13)	C15—C16—C12	118.99 (11)
C2—N1—H1	114.5 (13)	C11—C16—C12	119.05 (9)
N2—C2—N1	116.98 (10)	C26—C21—C22	120.68 (11)
N2—C2—S1	125.50 (9)	C26—C21—N2	118.43 (11)
N1—C2—S1	117.52 (8)	C22—C21—N2	120.88 (11)
C2—N2—C21	124.95 (10)	C23—C22—C21	118.48 (13)
C2—N2—H2	117.2 (14)	C23—C22—H22	120.8
C21—N2—H2	117.5 (14)	C21—C22—H22	120.8
C12—C11—C16	117.79 (11)	C24—C23—C22	121.91 (13)
C12—C11—C1	120.90 (11)	C24—C23—C13	119.18 (10)
C16—C11—C1	121.31 (11)	C22—C23—C13	118.89 (12)
C11—C12—C13	121.04 (14)	C23—C24—C25	118.67 (12)
C11—C12—C11	119.67 (10)	C23—C24—H24	120.7
C13—C12—C11	119.28 (12)	C25—C24—H24	120.7
C14—C13—C12	119.59 (15)	C24—C25—C26	120.60 (13)
C14—C13—H13	120.2	C24—C25—H25	119.7
C12—C13—H13	120.2	C26—C25—H25	119.7
C13—C14—C15	120.85 (13)	C21—C26—C25	119.62 (12)
C13—C14—H14	119.6	C21—C26—H26	120.2
C15—C14—H14	119.6	C25—C26—H26	120.2
O1—C1—N1—C2	0.6 (2)	C14—C15—C16—C11	0.2 (2)
C11—C1—N1—C2	-179.35 (12)	C14—C15—C16—C12	179.98 (12)
C1—N1—C2—N2	-4.11 (19)	C12—C11—C16—C15	-0.86 (18)
C1—N1—C2—S1	175.27 (11)	C1—C11—C16—C15	178.74 (12)
N1—C2—N2—C21	179.40 (11)	C12—C11—C16—C12	179.39 (9)
S1—C2—N2—C21	0.07 (18)	C1—C11—C16—C12	-1.01 (16)
O1—C1—C11—C12	-90.46 (16)	C2—N2—C21—C26	-130.55 (14)
N1—C1—C11—C12	89.48 (14)	C2—N2—C21—C22	50.86 (18)
O1—C1—C11—C16	89.95 (16)	C26—C21—C22—C23	2.4 (2)
N1—C1—C11—C16	-90.11 (14)	N2—C21—C22—C23	-179.07 (14)
C16—C11—C12—C13	0.68 (19)	C21—C22—C23—C24	-1.7 (3)
C1—C11—C12—C13	-178.93 (13)	C21—C22—C23—C13	176.96 (13)
C16—C11—C12—C11	179.53 (10)	C22—C23—C24—C25	-0.1 (3)
C1—C11—C12—C11	-0.07 (16)	C13—C23—C24—C25	-178.76 (14)
C11—C12—C13—C14	0.1 (2)	C23—C24—C25—C26	1.3 (3)
C11—C12—C13—C14	-178.73 (14)	C22—C21—C26—C25	-1.2 (2)
C12—C13—C14—C15	-0.8 (3)	N2—C21—C26—C25	-179.82 (13)
C13—C14—C15—C16	0.6 (2)	C24—C25—C26—C21	-0.6 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O1	0.82 (2)	2.07 (2)	2.7190 (13)	136.0 (18)
N2—H2 \cdots O1 ⁱ	0.82 (2)	2.37 (2)	3.0749 (14)	145.5 (18)
N1—H1 \cdots S1 ⁱⁱ	0.86 (2)	2.47 (2)	3.2974 (10)	163.6 (18)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y+1, -z+1$.

Fig. 1




Fig. 2

