

Redetermination of 1,4-dimethoxybenzene

Robbie Iuliucci,^{a*} Cody L. Hoop,^a Atta M. Arif,^b James K. Harper,^b Ronald J. Pugmire^b and David M. Grant^b

^aDepartment of Chemistry, Washington and Jefferson College, 60 South Lincoln Street, Washington, PA 15301, USA, and ^bDepartment of Chemistry, University of Utah, Salt Lake City, UT 84112, USA

Correspondence e-mail: riuliucci@washjeff.edu


Received 9 December 2008; accepted 30 December 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.037; wR factor = 0.101; data-to-parameter ratio = 12.8.

The structure of the centrosymmetric title compound, $\text{C}_8\text{H}_{10}\text{O}_2$, originally determined by Goodwin *et al.* [*Acta Cryst.* (1950), **3**, 279–284], has been redetermined to modern standards of precision to aid in its use as a model compound for ^{13}C chemical-shift tensor measurements in single-crystal NMR studies. In the crystal structure, a $\text{C}-\text{H}\cdots\text{O}$ interaction helps to establish the packing.

Related literature

For previous structural studies of the title compound, see: Goodwin *et al.* (1950); Carter *et al.* (1988).



Experimental

Crystal data

$\text{C}_8\text{H}_{10}\text{O}_2$
 $M_r = 138.16$
 Orthorhombic, *Pbca*

$a = 7.1757$ (3) Å
 $b = 6.2769$ (2) Å
 $c = 16.5573$ (7) Å

$V = 745.76$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.09$ mm⁻¹
 $T = 150$ (1) K
 $0.33 \times 0.30 \times 0.23$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (*DENZO-SMN*; Otwinowski & Minor, 1997)
 $T_{\min} = 0.972$, $T_{\max} = 0.980$

1510 measured reflections
 847 independent reflections
 732 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.101$
 $S = 1.09$
 847 reflections

66 parameters
 All H-atom parameters refined
 $\Delta\rho_{\max} = 0.19$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C4}-\text{H4A}\cdots\text{O1}^i$	1.019 (16)	2.552 (15)	3.4381 (15)	145.1 (10)

Symmetry code: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *WinGX* (Farrugia, 1999) and *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2009).

This work was supported by NSF grant ECC0304433 and NIH grant 5R01GM08521-44. Financial support for CLH was provided by Presidential Discretionary Funds from Washington and Jefferson College.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2878).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Carter, C. M., Facelli, J. C., Alderman, D. W., Grant, D. M., Dalley, N. K. & Wilson, B. E. (1988). *J. Chem. Soc. Faraday Trans. 1*, **84**, 3673–3690.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Goodwin, T. H., Przybylska, M. & Robertson, J. M. (1950). *Acta Cryst.* **3**, 279–284.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o251 [doi:10.1107/S1600536808044231]

Redetermination of 1,4-dimethoxybenzene

R. Iulucci, C. L. Hoop, A. M. Arif, J. K. Harper, R. J. Pugmire and D. M. Grant

Comment

Large single-crystals of organic compounds can be challenging to grow. Substituted methoxybenzenes are one exception and single-crystals on the order of centimeters can be obtained. The ease of crystal growth has enabled substituted methoxybenzenes to be studied by single-crystal NMR experiments. Pioneering work on the development of the two-dimensional single-crystal chemical-shift correlation NMR experiments utilized large crystals of 1,4-dimethoxybenzene (Carter *et al.*, 1988). In 1950 Goodwin *et al.* obtained the first X-ray diffraction structure for 1,4-dimethoxybenzene. This structure (R -factor = 0.12) is shown in Fig. 1 and reported an unusual H–C–C angle of 75.7° , which prompted the acquisition of a second structure (Carter *et al.*, 1988). More typical H–C–C angles were observed with this new refinement and this structure (R -factor = 0.067) was used to assign tensor orientations in the single-crystal NMR analysis. Inadvertently, the second structure was never submitted to the Cambridge Crystallographic database. Here, the acquisition of a third structure is reported to correct this oversight. The new structure (R -factor = 0.038) is shown in Fig. 2. The unit-cell and space group of the previous studies are confirmed.

Acquisition of this third, more accurate, structure is beneficial to NMR studies because the ^{13}C chemical shift tensor data of 1,4-dimethoxybenzene continue to serve as a standard to evaluate new chemical-shift tensor measurement methods as well as to assess electronic structure methods for computing magnetic properties of molecules.

Refinement

The H atoms were located in difference maps and their positions and U_{iso} values were freely refined.

Figures




Fig. 1. The structure of (I) according to Goodwin *et al.* (1950).




Fig. 2. The redetermined structure of (I) from the present study.

1,4-dimethoxybenzene

Crystal data

$C_8H_{10}O_2$

$M_r = 138.16$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 7.1757$ (3) Å

$b = 6.2769$ (2) Å

$c = 16.5573$ (7) Å

$V = 745.76$ (5) Å³

$Z = 4$

$F_{000} = 296$

$D_x = 1.231$ Mg m⁻³

Melting point: 329 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 13221 reflections

$\theta = 1.0$ – 27.5°

$\mu = 0.09$ mm⁻¹

$T = 150$ (1) K

Prism, colorless

$0.33 \times 0.30 \times 0.23$ mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 150$ (1) K

φ and ω scans

Absorption correction: Multi-scan
(DENZO-SMN; Otwinowski & Minor, 1997)

$T_{\min} = 0.972$, $T_{\max} = 0.980$

1510 measured reflections

847 independent reflections

732 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.013$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 3.8^\circ$

$h = -9 \rightarrow 9$

$k = -8 \rightarrow 8$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.101$

$S = 1.09$

Secondary atom site location: difference Fourier map

Hydrogen site location: difference Fourier map

All H-atom parameters refined

$w = 1/[\sigma^2(F_o^2) + (0.0517P)^2 + 0.1487P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

847 reflections $\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
 66 parameters $\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$
 none constraints Extinction correction: none
 Primary atom site location: structure-invariant direct
 methods

Special details

Experimental. The program *DENZO-SMN* (Otwinowski & Minor, 1997) uses a scaling algorithm (Fox & Holmes, 1966) which effectively corrects for absorption effects. High redundancy data were used in the scaling program hence the 'multi-scan' code word was used. No transmission coefficients are available from the program (only scale factors for each frame). The scale factors in the experimental table are calculated from the 'size' command in the *SHELXL97* input file.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.01305 (11)	0.15781 (13)	0.15601 (4)	0.0315 (3)
C2	0.09537 (13)	-0.10269 (16)	0.06126 (6)	0.0257 (3)
C3	-0.09472 (14)	0.18955 (17)	0.01604 (6)	0.0264 (3)
C1	0.00160 (12)	0.08587 (17)	0.07767 (6)	0.0243 (3)
C4	-0.0849 (2)	0.3490 (2)	0.17534 (8)	0.0410 (3)
H2	0.1619 (17)	-0.175 (2)	0.1048 (8)	0.033 (3)*
H3	-0.1604 (18)	0.319 (2)	0.0253 (7)	0.031 (3)*
H4A	-0.061 (2)	0.375 (2)	0.2352 (10)	0.050 (4)*
H4B	-0.221 (3)	0.323 (2)	0.1672 (9)	0.057 (5)*
H4C	-0.036 (2)	0.470 (3)	0.1413 (11)	0.056 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0387 (5)	0.0328 (5)	0.0231 (4)	0.0040 (3)	-0.0018 (3)	-0.0031 (3)
C2	0.0249 (5)	0.0268 (5)	0.0255 (5)	0.0015 (4)	-0.0017 (4)	0.0043 (4)
C3	0.0259 (5)	0.0244 (5)	0.0288 (6)	0.0029 (4)	0.0012 (4)	0.0014 (4)
C1	0.0241 (5)	0.0267 (5)	0.0220 (5)	-0.0028 (4)	0.0012 (3)	0.0007 (4)
C4	0.0534 (8)	0.0377 (7)	0.0321 (6)	0.0092 (6)	-0.0004 (5)	-0.0104 (5)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.3759 (12)	C3—C1	1.3937 (14)
-------	-------------	-------	-------------

supplementary materials

O1—C4	1.4269 (14)	C3—H3	0.953 (13)
C2—C1	1.3883 (15)	C4—H4A	1.020 (16)
C2—C3 ⁱ	1.3912 (15)	C4—H4B	0.998 (18)
C2—H2	0.977 (13)	C4—H4C	1.010 (18)
C3—C2 ⁱ	1.3912 (15)		
C1—O1—C4	117.26 (9)	O1—C1—C3	124.51 (10)
C1—C2—C3 ⁱ	120.83 (9)	C2—C1—C3	119.68 (10)
C1—C2—H2	119.4 (7)	O1—C4—H4A	105.8 (8)
C3 ⁱ —C2—H2	119.8 (7)	O1—C4—H4B	108.4 (9)
C2 ⁱ —C3—C1	119.48 (10)	H4A—C4—H4B	108.8 (12)
C2 ⁱ —C3—H3	118.7 (8)	O1—C4—H4C	109.7 (9)
C1—C3—H3	121.8 (8)	H4A—C4—H4C	111.2 (12)
O1—C1—C2	115.81 (9)	H4B—C4—H4C	112.6 (12)
C4—O1—C1—C2	-178.76 (10)	C3 ⁱ —C2—C1—C3	0.10 (16)
C4—O1—C1—C3	1.69 (15)	C2 ⁱ —C3—C1—O1	179.44 (9)
C3 ⁱ —C2—C1—O1	-179.47 (9)	C2 ⁱ —C3—C1—C2	-0.10 (16)

Symmetry codes: (i) $-x, -y, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4A \cdots O1 ⁱⁱ	1.019 (16)	2.552 (15)	3.4381 (15)	145.1 (10)

Symmetry codes: (ii) $-x, y+1/2, -z+1/2$.

Fig. 1




Fig. 2

