

2-Ethyl-6-methylanilinium 4-methylbenzenesulfonate

Tian-Quan Wu, Lin Xia, Ai-Xi Hu* and Jiao Ye

College of Chemistry and Chemical Engineering, Hunan University, Changsha 410082, People's Republic of China

Correspondence e-mail: axhu0731@yahoo.com.cn


Received 9 January 2009; accepted 18 January 2009

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.038; wR factor = 0.109; data-to-parameter ratio = 17.3.

The title compound, $\text{C}_9\text{H}_{14}\text{N}^+\cdot\text{C}_7\text{H}_7\text{SO}_3^-$, contains a 2-ethyl-6-methylanilinium cation and a 4-methylbenzenesulfonic anion. The cations are anchored between the anions through $\text{N}\cdots\text{O}$ hydrogen bonds. Electrostatic and van der Waals interactions, as well as hydrogen bonds, maintain the structural cohesion.

Related literature

For related structures, see: Benali-Cherif *et al.* (2007); Benslimane *et al.* (2007); Elmali *et al.* (2001); Fábry *et al.* (2001, 2002); Khemiri *et al.* (2008); Muthamizhchelvan *et al.* (2005); Smirani *et al.* (2008); Smirani & Rzaigui (2009).



Experimental

Crystal data

$\text{C}_9\text{H}_{14}\text{N}^+\cdot\text{C}_7\text{H}_7\text{O}_3\text{S}^-$
 $M_r = 307.40$
 Monoclinic, $P2_1/n$
 $a = 15.2514$ (9) Å
 $b = 6.1889$ (4) Å
 $c = 16.9242$ (10) Å
 $\beta = 102.850$ (1)°

$V = 1557.46$ (16) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.22$ mm⁻¹
 $T = 173$ (2) K
 $0.45 \times 0.34 \times 0.27$ mm

Data collection

Bruker SMART 1000 CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)
 $T_{\min} = 0.904$, $T_{\max} = 0.944$

7882 measured reflections
 3354 independent reflections
 2555 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.109$
 $S = 1.08$
 3354 reflections

194 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1C}\cdots\text{O1}^{\text{i}}$	0.91	1.88	2.7727 (18)	168
$\text{N1}-\text{H1B}\cdots\text{O3}^{\text{ii}}$	0.91	1.89	2.7800 (19)	166
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{iii}}$	0.91	1.95	2.7917 (18)	153

Symmetry codes: (i) $x, y, z + 1$; (ii) $-x + 2, -y + 1, -z + 1$; (iii) $x, y + 1, z + 1$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT-Plus (Bruker, 2003); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors express their thanks to the National Key Technology Research and Development Programme (grant No. 2006BAE01A01-4).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2133).

References

- Benali-Cherif, N., Boussekine, H., Boutobba, Z. & Kateb, A. (2007). *Acta Cryst.* **E63**, o3287.
 Benslimane, M., Merazig, H., Bouacida, S., Denbri, S., Beghidja, A. & Ouahab, L. (2007). *Acta Cryst.* **E63**, o3682–o3683.
 Bruker (2001). SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2003). SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
 Elmali, A., Elerman, Y. & Svoboda, I. (2001). *Acta Cryst.* **C57**, 485–486.
 Fábry, J., Krupková, R. & Studnička, V. (2002). *Acta Cryst.* **E58**, o105–o107.
 Fábry, J., Krupková, R. & Vaněk, P. (2001). *Acta Cryst.* **E57**, o1058–o1060.
 Khemiri, H., Akriche, S. & Rzaigui, M. (2008). *Acta Cryst.* **E64**, o526.
 Muthamizhchelvan, C., Saminathan, K., Fraanje, J., Peschar, R. & Sivakumar, K. (2005). *Acta Cryst.* **E61**, o1153–o1155.
 Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Smirani, W., Amri, O. & Rzaigui, M. (2008). *Acta Cryst.* **E64**, o2463.
 Smirani, W. & Rzaigui, M. (2009). *Acta Cryst.* **E65**, o83.

supplementary materials

Acta Cryst. (2009). E65, o368 [doi:10.1107/S160053680900230X]

2-Ethyl-6-methylanilinium 4-methylbenzenesulfonate

T.-Q. Wu, L. Xia, A.-X. Hu and J. Ye

Comment

In the title compound, the proton of the sulfonic group of sulfonic acid has been transferred to the N atom of the 2-ethyl-6-methylaniline molecule, leading to the formation of the molecular complex, (I). In this article we present the crystal structure of the title compound. The removal of the sulfonic acid H atom leads to a shortening of the C10—S1 bond length, showing a partial double-bond character. This behaviour is similar to that observed in many picrate salts (Muthamizhchelvan *et al.* 2005); it is attributed to the loss of the hydroxyl proton, leading to the conversion of the neutral to an anionic state of the molecule.

In the structure (Fig. 1), intermolecular hydrogen bonds are observed, with the N atom of the cation acting as donors. The orientation of the anion and cation facilitates the formation of the expected strong N—H \cdots O hydrogen bonds between amino atom N1 and the sulfonic O atoms; N1 hydrogen bonds to two sulfonic O atoms of adjacent molecules (Fig. 2 and Table 1). These hydrogen bonds are effective in the stabilization of the structure. The crystal structures of several related compounds have been published, e.g. Muthamizhchelvan *et al.* (2008); Benslimane *et al.* (2007); Smirani & Rzaigui (2009); Fábry *et al.* (2001, 2002); Khemiri *et al.* (2008); Benali-Cherif *et al.* (2007); Elmali *et al.* (2001).

Experimental

Added 2-ethyl-6-methylaniline (1.35 g) into a solution of 4-methylbenzenesulfonic acid (1.72 g) and ethanol (10 ml). After 10 min precipitate were formed which were filtered and dried, giving the desired product. Crystals suitable for X-ray structure determination were obtained by slow evaporation of an ethanol solution at room temperature.

Refinement

The methyl H atom were positioned geometrically (C—H = 0.98 Å) and torsion angles refined to fit the electron density [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$]. Other H atoms were placed at calculated positions (methylene C—H = 0.95 Å and aromatic C—H = 0.95 Å) and included in the refinement in a riding mode with [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$].

Figures




Fig. 1. Molecular structure of the title compound showing 30% probability displacement ellipsoids.




Fig. 2. Unit cell packing diagram of (I) showing intermolecular hydrogen bonds; H atoms have been omitted for clarity.

2-Ethyl-6-methylanilinium 4-methylbenzenesulfonate

Crystal data

$C_9H_{14}N^+ \cdot C_7H_7O_3S^-$	$F_{000} = 656$
$M_r = 307.40$	$D_x = 1.311 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 428 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation
$a = 15.2514 (9) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 6.1889 (4) \text{ \AA}$	Cell parameters from 3656 reflections
$c = 16.9242 (10) \text{ \AA}$	$\theta = 2.5\text{--}27.0^\circ$
$\beta = 102.8500 (10)^\circ$	$\mu = 0.22 \text{ mm}^{-1}$
$V = 1557.46 (16) \text{ \AA}^3$	$T = 173 (2) \text{ K}$
$Z = 4$	Block, colourless
	$0.45 \times 0.34 \times 0.27 \text{ mm}$

Data collection

Bruker SMART 1000 CCD diffractometer	3354 independent reflections
Radiation source: fine-focus sealed tube	2555 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
$T = 173(2) \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	$h = -16 \rightarrow 19$
$T_{\text{min}} = 0.904$, $T_{\text{max}} = 0.944$	$k = -7 \rightarrow 7$
7882 measured reflections	$l = -21 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0601P)^2 + 0.2665P]$
$S = 1.08$	where $P = (F_o^2 + 2F_c^2)/3$
3354 reflections	$(\Delta/\sigma)_{\text{max}} = 0.003$
194 parameters	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.34 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.90482 (3)	0.20760 (7)	0.10019 (2)	0.02108 (14)
C1	0.78421 (10)	0.5823 (3)	0.90053 (9)	0.0194 (3)
C2	0.80908 (11)	0.4025 (3)	0.86141 (10)	0.0221 (4)
C3	0.74104 (12)	0.2913 (3)	0.80842 (10)	0.0268 (4)
H3	0.7557	0.1687	0.7802	0.032*
C4	0.65245 (12)	0.3572 (3)	0.79640 (11)	0.0299 (4)
H4	0.6068	0.2791	0.7603	0.036*
C5	0.62989 (12)	0.5364 (3)	0.83675 (11)	0.0275 (4)
H5	0.5687	0.5799	0.8279	0.033*
C6	0.69564 (11)	0.6543 (3)	0.89020 (10)	0.0214 (4)
C7	0.90521 (12)	0.3284 (3)	0.87383 (11)	0.0290 (4)
H7A	0.9409	0.4405	0.8547	0.044*
H7B	0.9077	0.1949	0.8433	0.044*
H7C	0.9297	0.3017	0.9316	0.044*
C8	0.67261 (11)	0.8507 (3)	0.93471 (11)	0.0267 (4)
H8A	0.7064	0.9759	0.9204	0.032*
H8B	0.6937	0.8256	0.9936	0.032*
C9	0.57363 (13)	0.9096 (4)	0.91767 (15)	0.0467 (6)
H9A	0.5525	0.9429	0.8600	0.070*
H9B	0.5653	1.0361	0.9500	0.070*
H9C	0.5392	0.7876	0.9320	0.070*
C10	0.88515 (10)	0.1172 (3)	0.19401 (10)	0.0206 (3)
C11	0.90431 (11)	0.2524 (3)	0.26121 (10)	0.0246 (4)
H11	0.9258	0.3948	0.2564	0.029*
C12	0.89200 (12)	0.1787 (3)	0.33539 (11)	0.0281 (4)
H12	0.9053	0.2716	0.3812	0.034*
C13	0.86041 (11)	-0.0297 (3)	0.34369 (11)	0.0266 (4)
C14	0.84062 (12)	-0.1613 (3)	0.27551 (12)	0.0302 (4)
H14	0.8180	-0.3028	0.2799	0.036*
C15	0.85323 (12)	-0.0899 (3)	0.20085 (11)	0.0283 (4)
H15	0.8401	-0.1825	0.1550	0.034*
C16	0.84628 (14)	-0.1093 (4)	0.42440 (12)	0.0380 (5)
H16A	0.8966	-0.0624	0.4677	0.057*

supplementary materials

H16B	0.8430	-0.2675	0.4238	0.057*
H16C	0.7900	-0.0497	0.4340	0.057*
N1	0.85616 (9)	0.6956 (2)	0.95856 (8)	0.0200 (3)
H1A	0.8349	0.8242	0.9725	0.030*
H1B	0.9036	0.7195	0.9352	0.030*
H1C	0.8743	0.6129	1.0037	0.030*
O1	0.88551 (8)	0.43864 (19)	0.09615 (7)	0.0257 (3)
O2	0.84407 (8)	0.0861 (2)	0.03804 (7)	0.0307 (3)
O3	0.99931 (8)	0.1642 (2)	0.10318 (8)	0.0311 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0223 (2)	0.0194 (2)	0.0217 (2)	0.00109 (16)	0.00531 (16)	-0.00141 (17)
C1	0.0208 (8)	0.0203 (8)	0.0165 (8)	-0.0022 (6)	0.0031 (6)	0.0025 (6)
C2	0.0273 (9)	0.0191 (8)	0.0200 (8)	0.0006 (7)	0.0055 (7)	0.0029 (7)
C3	0.0354 (10)	0.0221 (9)	0.0236 (9)	-0.0033 (7)	0.0078 (7)	-0.0031 (7)
C4	0.0294 (10)	0.0329 (10)	0.0253 (9)	-0.0110 (8)	0.0019 (7)	-0.0032 (8)
C5	0.0217 (9)	0.0322 (10)	0.0276 (9)	-0.0028 (7)	0.0037 (7)	0.0022 (8)
C6	0.0226 (8)	0.0227 (9)	0.0191 (8)	-0.0006 (7)	0.0051 (6)	0.0040 (7)
C7	0.0310 (10)	0.0255 (10)	0.0307 (10)	0.0039 (7)	0.0072 (8)	-0.0050 (8)
C8	0.0208 (9)	0.0275 (9)	0.0313 (10)	0.0002 (7)	0.0047 (7)	-0.0035 (8)
C9	0.0288 (11)	0.0464 (13)	0.0616 (15)	0.0094 (10)	0.0029 (10)	-0.0170 (11)
C10	0.0187 (8)	0.0204 (8)	0.0232 (8)	0.0015 (6)	0.0055 (6)	-0.0007 (7)
C11	0.0241 (9)	0.0231 (9)	0.0267 (9)	-0.0039 (7)	0.0062 (7)	-0.0028 (7)
C12	0.0276 (9)	0.0316 (10)	0.0252 (9)	-0.0024 (8)	0.0060 (7)	-0.0048 (8)
C13	0.0203 (9)	0.0315 (10)	0.0294 (9)	0.0036 (7)	0.0086 (7)	0.0040 (8)
C14	0.0331 (10)	0.0208 (9)	0.0390 (11)	-0.0013 (7)	0.0132 (8)	0.0027 (8)
C15	0.0328 (10)	0.0224 (9)	0.0312 (10)	-0.0029 (7)	0.0102 (8)	-0.0051 (8)
C16	0.0379 (11)	0.0462 (12)	0.0331 (11)	0.0019 (9)	0.0148 (9)	0.0071 (9)
N1	0.0186 (7)	0.0212 (7)	0.0197 (7)	0.0017 (6)	0.0031 (5)	-0.0007 (6)
O1	0.0323 (7)	0.0203 (6)	0.0238 (6)	0.0022 (5)	0.0050 (5)	0.0008 (5)
O2	0.0375 (7)	0.0285 (7)	0.0248 (7)	-0.0043 (6)	0.0044 (5)	-0.0072 (5)
O3	0.0250 (7)	0.0335 (8)	0.0374 (7)	0.0061 (5)	0.0125 (5)	0.0071 (6)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4482 (12)	C8—H8B	0.9900
S1—O3	1.4557 (13)	C9—H9A	0.9800
S1—O1	1.4584 (12)	C9—H9B	0.9800
S1—C10	1.7709 (17)	C9—H9C	0.9800
C1—C2	1.390 (2)	C10—C15	1.385 (2)
C1—C6	1.396 (2)	C10—C11	1.390 (2)
C1—N1	1.477 (2)	C11—C12	1.387 (2)
C2—C3	1.393 (2)	C11—H11	0.9500
C2—C7	1.505 (2)	C12—C13	1.395 (2)
C3—C4	1.382 (3)	C12—H12	0.9500
C3—H3	0.9500	C13—C14	1.389 (3)
C4—C5	1.386 (3)	C13—C16	1.512 (2)

C4—H4	0.9500	C14—C15	1.392 (3)
C5—C6	1.397 (2)	C14—H14	0.9500
C5—H5	0.9500	C15—H15	0.9500
C6—C8	1.512 (2)	C16—H16A	0.9800
C7—H7A	0.9800	C16—H16B	0.9800
C7—H7B	0.9800	C16—H16C	0.9800
C7—H7C	0.9800	N1—H1A	0.9100
C8—C9	1.517 (2)	N1—H1B	0.9100
C8—H8A	0.9900	N1—H1C	0.9100
O2—S1—O3	113.45 (8)	C8—C9—H9A	109.5
O2—S1—O1	112.66 (7)	C8—C9—H9B	109.5
O3—S1—O1	111.75 (7)	H9A—C9—H9B	109.5
O2—S1—C10	106.17 (8)	C8—C9—H9C	109.5
O3—S1—C10	105.92 (7)	H9A—C9—H9C	109.5
O1—S1—C10	106.21 (7)	H9B—C9—H9C	109.5
C2—C1—C6	123.69 (15)	C15—C10—C11	120.17 (16)
C2—C1—N1	117.11 (14)	C15—C10—S1	120.01 (13)
C6—C1—N1	119.15 (14)	C11—C10—S1	119.79 (13)
C1—C2—C3	117.31 (15)	C12—C11—C10	119.79 (16)
C1—C2—C7	122.52 (15)	C12—C11—H11	120.1
C3—C2—C7	120.16 (16)	C10—C11—H11	120.1
C4—C3—C2	120.86 (16)	C11—C12—C13	120.95 (17)
C4—C3—H3	119.6	C11—C12—H12	119.5
C2—C3—H3	119.6	C13—C12—H12	119.5
C3—C4—C5	120.35 (16)	C14—C13—C12	118.31 (16)
C3—C4—H4	119.8	C14—C13—C16	120.74 (17)
C5—C4—H4	119.8	C12—C13—C16	120.94 (17)
C4—C5—C6	121.05 (16)	C13—C14—C15	121.31 (17)
C4—C5—H5	119.5	C13—C14—H14	119.3
C6—C5—H5	119.5	C15—C14—H14	119.3
C1—C6—C5	116.73 (15)	C10—C15—C14	119.45 (17)
C1—C6—C8	121.31 (15)	C10—C15—H15	120.3
C5—C6—C8	121.96 (15)	C14—C15—H15	120.3
C2—C7—H7A	109.5	C13—C16—H16A	109.5
C2—C7—H7B	109.5	C13—C16—H16B	109.5
H7A—C7—H7B	109.5	H16A—C16—H16B	109.5
C2—C7—H7C	109.5	C13—C16—H16C	109.5
H7A—C7—H7C	109.5	H16A—C16—H16C	109.5
H7B—C7—H7C	109.5	H16B—C16—H16C	109.5
C6—C8—C9	115.46 (15)	C1—N1—H1A	109.5
C6—C8—H8A	108.4	C1—N1—H1B	109.5
C9—C8—H8A	108.4	H1A—N1—H1B	109.5
C6—C8—H8B	108.4	C1—N1—H1C	109.5
C9—C8—H8B	108.4	H1A—N1—H1C	109.5
H8A—C8—H8B	107.5	H1B—N1—H1C	109.5
C6—C1—C2—C3	0.8 (2)	O2—S1—C10—C15	27.27 (16)
N1—C1—C2—C3	178.16 (14)	O3—S1—C10—C15	-93.63 (15)
C6—C1—C2—C7	-179.91 (16)	O1—S1—C10—C15	147.40 (13)

supplementary materials

N1—C1—C2—C7	-2.5 (2)	O2—S1—C10—C11	-154.60 (13)
C1—C2—C3—C4	-0.7 (3)	O3—S1—C10—C11	84.50 (14)
C7—C2—C3—C4	179.95 (16)	O1—S1—C10—C11	-34.46 (15)
C2—C3—C4—C5	0.4 (3)	C15—C10—C11—C12	0.5 (3)
C3—C4—C5—C6	0.0 (3)	S1—C10—C11—C12	-177.65 (13)
C2—C1—C6—C5	-0.5 (2)	C10—C11—C12—C13	-0.1 (3)
N1—C1—C6—C5	-177.76 (14)	C11—C12—C13—C14	-0.7 (3)
C2—C1—C6—C8	179.76 (16)	C11—C12—C13—C16	-179.50 (16)
N1—C1—C6—C8	2.5 (2)	C12—C13—C14—C15	1.1 (3)
C4—C5—C6—C1	0.0 (2)	C16—C13—C14—C15	179.94 (17)
C4—C5—C6—C8	179.83 (17)	C11—C10—C15—C14	-0.1 (3)
C1—C6—C8—C9	-179.87 (17)	S1—C10—C15—C14	178.07 (13)
C5—C6—C8—C9	0.4 (3)	C13—C14—C15—C10	-0.8 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1C \cdots O1 ⁱ	0.91	1.88	2.7727 (18)	168
N1—H1B \cdots O3 ⁱⁱ	0.91	1.89	2.7800 (19)	166
N1—H1A \cdots O2 ⁱⁱⁱ	0.91	1.95	2.7917 (18)	153

Symmetry codes: (i) $x, y, z+1$; (ii) $-x+2, -y+1, -z+1$; (iii) $x, y+1, z+1$.

Fig. 1




Fig. 2

