

N-[2-[2-(2,6-Dichloro-3,5-dimethoxyphenyl)ethenyl]phenyl]acetamide

Kartini Ahmad,^a Khalijah Awang,^a Noel F. Thomas,^a Jean-Frederic Faizal Weber^b and Seik Weng Ng^{a*}

^aDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^bFaculty of Pharmacy, Universiti Teknologi MARA, 40450 Shah Alam, Selangor Darul Ehsan, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 24 January 2009; accepted 26 January 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.048; wR factor = 0.117; data-to-parameter ratio = 16.6.

The $\text{C}=\text{C}$ double bond in the title substituted stilbene, $\text{C}_{18}\text{H}_{17}\text{Cl}_2\text{NO}_3$, has a *trans* arrangement of the aryl substituents. The aromatic ring of the 2-acetylaminophenyl substituent is twisted by $39.9(3)^\circ$ with respect to the central $\text{C}-\text{C}=\text{C}-\text{C}$ unit and that of the 2,6-dichloro-3,5-dimethoxyphenyl substituent is twisted by $42.7(3)^\circ$.

Related literature

The compound was synthesized by a ferric chloride-promoted highly atropodiastereoselective cascade reaction, a reaction that illustrates the utility of radical cations in indolostilbene synthesis; see: Ahmad *et al.* (2009).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{17}\text{Cl}_2\text{NO}_3$	$\gamma = 85.969(3)^\circ$
$M_r = 366.23$	$V = 814.65(5)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.5646(3)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.1485(3)\text{ \AA}$	$\mu = 0.42\text{ mm}^{-1}$
$c = 12.2969(5)\text{ \AA}$	$T = 100(2)\text{ K}$
$\alpha = 78.561(2)^\circ$	$0.30 \times 0.03 \times 0.03\text{ mm}$
$\beta = 77.716(2)^\circ$	

Data collection

Bruker SMART APEX	6677 measured reflections
diffractometer	3657 independent reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	2490 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.886$, $T_{\max} = 0.988$	$R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$	220 parameters
$wR(F^2) = 0.117$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
3657 reflections	$\Delta\rho_{\min} = -0.37\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

The authors thank the University of Malaya for supporting this study (Science Fund grant No. 1402034013).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2361).

References

- Ahmad, K., Thomas, N. F., Mukhtar, M. R., Noorbacha, I., Weber, J.-F. F., Nafiah, M. A., Velu, S. S., Takeya, K., Morita, H., Lim, C.-G., Hadi, A. H. A. & Awang, K. (2009). *Tetrahedron*, **65**, 1504–1516.
Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o438 [doi:10.1107/S1600536809003250]

N-{2-[2-(2,6-Dichloro-3,5-dimethoxyphenyl)ethenyl]phenyl}acetamide

K. Ahmad, K. Awang, N. F. Thomas, J.-F. F. Weber and S. W. Ng

Comment

The compound was synthesized by a ferric-chloride promoted, highly atropodiastereoselective cascade reaction, a reaction that illustrates the utility of radical cations in indolostilbene synthesis. The description of the synthesis is given in a recent study (Ahmad *et al.*, 2009).

Experimental

The synthesis is described in an earlier report (Ahmad *et al.*, 2009).

Refinement

Carbon-bound H atoms were placed in calculated positions (C—H 0.95–0.98 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U(\text{C})$. The nitrogen-bound H-atom was similarly treated (N—H 0.88 Å, $U(\text{H}) = 1.2U(\text{N})$).

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{18}\text{H}_{17}\text{Cl}_2\text{NO}_3$ at the 70% probability level. H atoms are drawn as spheres of arbitrary radius.

N-{2-[2-(2,6-Dichloro-3,5-dimethoxyphenyl)ethenyl]phenyl}acetamide

Crystal data

$\text{C}_{18}\text{H}_{17}\text{Cl}_2\text{NO}_3$	$Z = 2$
$M_r = 366.23$	$F_{000} = 380$
Triclinic, $P\bar{1}$	$D_x = 1.493 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.5646 (3) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 9.1485 (3) \text{ \AA}$	Cell parameters from 1123 reflections
$c = 12.2969 (5) \text{ \AA}$	$\theta = 2.6\text{--}26.5^\circ$
$\alpha = 78.561 (2)^\circ$	$\mu = 0.42 \text{ mm}^{-1}$
$\beta = 77.716 (2)^\circ$	$T = 100 (2) \text{ K}$
	Prism, colourless

supplementary materials

$\gamma = 85.969 (3)^\circ$ $0.30 \times 0.03 \times 0.03$ mm
 $V = 814.65 (5) \text{ \AA}^3$

Data collection

Bruker SMART APEX diffractometer	3657 independent reflections
Radiation source: fine-focus sealed tube	2490 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.048$
$T = 100(2)$ K	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.886$, $T_{\text{max}} = 0.988$	$k = -11 \rightarrow 11$
6677 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.117$	$w = 1/[\sigma^2(F_o^2) + (0.046P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.99$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3657 reflections	$\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
220 parameters	$\Delta\rho_{\text{min}} = -0.37 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.21992 (9)	0.40664 (7)	0.21221 (5)	0.02175 (18)
Cl2	0.28411 (9)	0.66314 (7)	0.56854 (5)	0.01944 (18)
O1	0.3311 (3)	0.68767 (19)	0.08280 (14)	0.0248 (5)
O2	0.4014 (2)	0.90427 (18)	0.39962 (15)	0.0203 (4)
O3	0.1911 (3)	0.2547 (2)	0.96831 (17)	0.0353 (5)
N1	0.1666 (3)	0.3365 (2)	0.78526 (18)	0.0198 (5)
H1	0.1734	0.4173	0.7322	0.024*
C1	0.2581 (3)	0.5380 (3)	0.3861 (2)	0.0141 (5)
C2	0.2725 (3)	0.5540 (3)	0.2691 (2)	0.0162 (6)
C3	0.3281 (4)	0.6856 (3)	0.1940 (2)	0.0171 (6)
C4	0.3763 (4)	0.8042 (3)	0.2354 (2)	0.0183 (6)
H4	0.4190	0.8928	0.1845	0.022*
C5	0.3624 (3)	0.7936 (3)	0.3506 (2)	0.0158 (6)
C6	0.3045 (3)	0.6611 (3)	0.4245 (2)	0.0149 (5)
C7	0.3852 (4)	0.8217 (3)	0.0026 (2)	0.0285 (7)

H7A	0.3723	0.8102	-0.0729	0.043*
H7B	0.5119	0.8405	0.0009	0.043*
H7C	0.3085	0.9060	0.0250	0.043*
C8	0.4638 (4)	1.0415 (3)	0.3273 (2)	0.0232 (6)
H8A	0.4852	1.1118	0.3735	0.035*
H8B	0.3721	1.0843	0.2836	0.035*
H8C	0.5768	1.0225	0.2753	0.035*
C9	0.1833 (3)	0.4005 (3)	0.4615 (2)	0.0154 (6)
H9	0.0883	0.3559	0.4413	0.019*
C10	0.2381 (3)	0.3337 (3)	0.5558 (2)	0.0144 (5)
H10	0.3401	0.3734	0.5723	0.017*
C11	0.1548 (3)	0.2039 (3)	0.6366 (2)	0.0159 (6)
C12	0.1058 (4)	0.0794 (3)	0.6023 (2)	0.0193 (6)
H12	0.1223	0.0778	0.5238	0.023*
C13	0.0330 (4)	-0.0426 (3)	0.6812 (2)	0.0207 (6)
H13	-0.0016	-0.1264	0.6565	0.025*
C14	0.0106 (4)	-0.0427 (3)	0.7955 (2)	0.0222 (6)
H14	-0.0367	-0.1275	0.8492	0.027*
C15	0.0568 (4)	0.0805 (3)	0.8323 (2)	0.0203 (6)
H15	0.0404	0.0803	0.9111	0.024*
C16	0.1273 (4)	0.2042 (3)	0.7536 (2)	0.0178 (6)
C17	0.1954 (4)	0.3555 (3)	0.8874 (2)	0.0234 (6)
C18	0.2350 (4)	0.5142 (3)	0.8907 (2)	0.0289 (7)
H18A	0.2924	0.5147	0.9549	0.043*
H18B	0.1216	0.5736	0.8993	0.043*
H18C	0.3165	0.5572	0.8201	0.043*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0352 (4)	0.0150 (3)	0.0164 (3)	-0.0059 (3)	-0.0067 (3)	-0.0029 (3)
Cl2	0.0257 (4)	0.0188 (3)	0.0144 (3)	-0.0049 (3)	-0.0029 (3)	-0.0043 (3)
O1	0.0446 (13)	0.0180 (10)	0.0117 (9)	-0.0104 (9)	-0.0079 (9)	0.0034 (8)
O2	0.0262 (11)	0.0149 (9)	0.0198 (10)	-0.0069 (8)	-0.0029 (8)	-0.0033 (8)
O3	0.0505 (14)	0.0387 (12)	0.0172 (11)	-0.0163 (11)	-0.0106 (10)	0.0027 (10)
N1	0.0279 (13)	0.0169 (11)	0.0133 (11)	-0.0032 (10)	-0.0023 (10)	-0.0004 (9)
C1	0.0100 (12)	0.0137 (12)	0.0179 (13)	0.0009 (9)	-0.0022 (10)	-0.0028 (10)
C2	0.0182 (14)	0.0156 (12)	0.0161 (13)	-0.0017 (10)	-0.0053 (11)	-0.0039 (11)
C3	0.0192 (14)	0.0166 (13)	0.0153 (13)	-0.0008 (11)	-0.0042 (11)	-0.0015 (11)
C4	0.0227 (15)	0.0133 (12)	0.0166 (14)	-0.0038 (11)	-0.0019 (11)	0.0016 (11)
C5	0.0157 (13)	0.0139 (12)	0.0192 (14)	0.0007 (10)	-0.0047 (11)	-0.0056 (11)
C6	0.0151 (13)	0.0183 (13)	0.0103 (12)	0.0009 (10)	-0.0016 (10)	-0.0019 (10)
C7	0.0440 (19)	0.0234 (15)	0.0151 (14)	-0.0083 (14)	-0.0060 (13)	0.0057 (12)
C8	0.0253 (16)	0.0149 (13)	0.0282 (16)	-0.0048 (11)	-0.0065 (13)	0.0012 (12)
C9	0.0195 (14)	0.0121 (12)	0.0149 (13)	-0.0008 (10)	-0.0027 (11)	-0.0037 (10)
C10	0.0153 (13)	0.0119 (12)	0.0158 (13)	-0.0029 (10)	-0.0005 (11)	-0.0038 (10)
C11	0.0165 (14)	0.0148 (13)	0.0150 (13)	0.0008 (10)	-0.0046 (11)	0.0016 (11)
C12	0.0195 (14)	0.0192 (13)	0.0183 (14)	0.0020 (11)	-0.0042 (11)	-0.0020 (11)

supplementary materials

C13	0.0213 (15)	0.0108 (12)	0.0285 (16)	-0.0014 (11)	-0.0034 (12)	-0.0020 (11)
C14	0.0211 (15)	0.0190 (14)	0.0223 (15)	-0.0022 (11)	-0.0023 (12)	0.0046 (12)
C15	0.0213 (15)	0.0217 (14)	0.0147 (13)	0.0006 (11)	-0.0008 (11)	0.0008 (11)
C16	0.0182 (14)	0.0175 (13)	0.0178 (14)	0.0016 (11)	-0.0043 (11)	-0.0040 (11)
C17	0.0220 (15)	0.0321 (16)	0.0155 (14)	-0.0029 (13)	-0.0018 (12)	-0.0046 (13)
C18	0.0397 (19)	0.0290 (16)	0.0211 (15)	-0.0074 (14)	-0.0058 (14)	-0.0105 (13)

Geometric parameters (\AA , $^\circ$)

C11—C2	1.742 (2)	C8—H8A	0.9800
Cl2—C6	1.748 (2)	C8—H8B	0.9800
O1—C3	1.359 (3)	C8—H8C	0.9800
O1—C7	1.437 (3)	C9—C10	1.333 (4)
O2—C5	1.355 (3)	C9—H9	0.9500
O2—C8	1.433 (3)	C10—C11	1.471 (3)
O3—C17	1.211 (3)	C10—H10	0.9500
N1—C17	1.365 (3)	C11—C12	1.389 (3)
N1—C16	1.411 (3)	C11—C16	1.410 (3)
N1—H1	0.8800	C12—C13	1.386 (4)
C1—C6	1.397 (3)	C12—H12	0.9500
C1—C2	1.399 (3)	C13—C14	1.379 (4)
C1—C9	1.476 (3)	C13—H13	0.9500
C2—C3	1.395 (3)	C14—C15	1.386 (4)
C3—C4	1.387 (3)	C14—H14	0.9500
C4—C5	1.382 (3)	C15—C16	1.391 (4)
C4—H4	0.9500	C15—H15	0.9500
C5—C6	1.398 (3)	C17—C18	1.513 (4)
C7—H7A	0.9800	C18—H18A	0.9800
C7—H7B	0.9800	C18—H18B	0.9800
C7—H7C	0.9800	C18—H18C	0.9800
C3—O1—C7	118.2 (2)	H8B—C8—H8C	109.5
C5—O2—C8	118.00 (19)	C10—C9—C1	125.3 (2)
C17—N1—C16	128.5 (2)	C10—C9—H9	117.3
C17—N1—H1	115.7	C1—C9—H9	117.3
C16—N1—H1	115.7	C9—C10—C11	125.8 (2)
C6—C1—C2	116.4 (2)	C9—C10—H10	117.1
C6—C1—C9	124.0 (2)	C11—C10—H10	117.1
C2—C1—C9	119.5 (2)	C12—C11—C16	118.6 (2)
C3—C2—C1	122.4 (2)	C12—C11—C10	122.6 (2)
C3—C2—Cl1	117.50 (19)	C16—C11—C10	118.8 (2)
C1—C2—Cl1	120.11 (19)	C13—C12—C11	120.9 (2)
O1—C3—C4	124.3 (2)	C13—C12—H12	119.6
O1—C3—C2	116.4 (2)	C11—C12—H12	119.6
C4—C3—C2	119.4 (2)	C14—C13—C12	120.1 (2)
C5—C4—C3	120.1 (2)	C14—C13—H13	119.9
C5—C4—H4	120.0	C12—C13—H13	119.9
C3—C4—H4	120.0	C13—C14—C15	120.3 (2)
O2—C5—C4	124.6 (2)	C13—C14—H14	119.8
O2—C5—C6	115.8 (2)	C15—C14—H14	119.8

C4—C5—C6	119.6 (2)	C14—C15—C16	119.8 (2)
C5—C6—C1	122.2 (2)	C14—C15—H15	120.1
C5—C6—Cl2	115.48 (18)	C16—C15—H15	120.1
C1—C6—Cl2	122.29 (19)	C15—C16—N1	122.6 (2)
O1—C7—H7A	109.5	C15—C16—C11	120.3 (2)
O1—C7—H7B	109.5	N1—C16—C11	117.0 (2)
H7A—C7—H7B	109.5	O3—C17—N1	123.4 (3)
O1—C7—H7C	109.5	O3—C17—C18	122.7 (3)
H7A—C7—H7C	109.5	N1—C17—C18	114.0 (2)
H7B—C7—H7C	109.5	C17—C18—H18A	109.5
O2—C8—H8A	109.5	C17—C18—H18B	109.5
O2—C8—H8B	109.5	H18A—C18—H18B	109.5
H8A—C8—H8B	109.5	C17—C18—H18C	109.5
O2—C8—H8C	109.5	H18A—C18—H18C	109.5
H8A—C8—H8C	109.5	H18B—C18—H18C	109.5
C6—C1—C2—C3	-0.7 (4)	C2—C1—C6—Cl2	176.94 (19)
C9—C1—C2—C3	174.7 (2)	C9—C1—C6—Cl2	1.8 (4)
C6—C1—C2—Cl1	179.43 (19)	C6—C1—C9—C10	-40.8 (4)
C9—C1—C2—Cl1	-5.2 (3)	C2—C1—C9—C10	144.2 (3)
C7—O1—C3—C4	-0.9 (4)	C1—C9—C10—C11	174.6 (2)
C7—O1—C3—C2	179.2 (2)	C9—C10—C11—C12	45.5 (4)
C1—C2—C3—O1	-178.1 (2)	C9—C10—C11—C16	-136.0 (3)
Cl1—C2—C3—O1	1.8 (3)	C16—C11—C12—C13	-0.6 (4)
C1—C2—C3—C4	2.0 (4)	C10—C11—C12—C13	177.9 (2)
Cl1—C2—C3—C4	-178.1 (2)	C11—C12—C13—C14	-0.9 (4)
O1—C3—C4—C5	177.6 (2)	C12—C13—C14—C15	1.5 (4)
C2—C3—C4—C5	-2.5 (4)	C13—C14—C15—C16	-0.5 (4)
C8—O2—C5—C4	-0.8 (4)	C14—C15—C16—N1	176.0 (2)
C8—O2—C5—C6	179.1 (2)	C14—C15—C16—C11	-1.1 (4)
C3—C4—C5—O2	-178.2 (2)	C17—N1—C16—C15	20.7 (4)
C3—C4—C5—C6	1.9 (4)	C17—N1—C16—C11	-162.1 (3)
O2—C5—C6—C1	179.5 (2)	C12—C11—C16—C15	1.6 (4)
C4—C5—C6—C1	-0.6 (4)	C10—C11—C16—C15	-177.0 (2)
O2—C5—C6—Cl2	2.3 (3)	C12—C11—C16—N1	-175.6 (2)
C4—C5—C6—Cl2	-177.7 (2)	C10—C11—C16—N1	5.8 (4)
C2—C1—C6—C5	0.0 (4)	C16—N1—C17—O3	0.3 (5)
C9—C1—C6—C5	-175.1 (2)	C16—N1—C17—C18	179.8 (2)

supplementary materials

Fig. 1

