

[1,3-Bis(diphenylphosphino)pentane- κ^2P,P']tetracarbonylchromium(0)

Omar bin Shawkataly,^{a*}‡ Daniel T. Thangadurai,^a Mohd. Aslam A. Pankhi,^a S. M. Shahinoor Dusal Islam^a and Hoong-Kun Fun^b

^aChemical Sciences Programme, Centre for Distance Education, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bX-Ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia
Correspondence e-mail: omarsa@usm.my


Received 7 October 2008; accepted 10 January 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.031; wR factor = 0.072; data-to-parameter ratio = 20.4.

In the title compound, $[\text{Cr}(\text{C}_{29}\text{H}_{30}\text{P}_2)(\text{CO})_4]$, the Cr atom is octahedrally coordinated by four carbonyl ligands and one bidentate phosphine ligand, which is bounded as a chelate in a *cis* position. The average Cr–P and Cr–C bond lengths are 2.377 and 1.865 Å, respectively.

Related literature

For chromium–carbonyl complexes see: Shawkataly *et al.* (1996, 1997, 2006); for Cr–C bond lengths see: Bennett *et al.* (1971); Ueng & Shih (1992). For Cr–C and C–O distances see Whitaker & Jeffery (1967); Jost *et al.* (1975). For a description of the Cambridge Structural Database, see: Allen (2002).



‡ On secondment to: Multimedia University, Melaka Campus, Jalan Ayer Keroh Lama, 74750 Melaka, Malaysia.

Experimental

Crystal data

$[\text{Cr}(\text{C}_{29}\text{H}_{30}\text{P}_2)(\text{CO})_4]$	$V = 2966.55$ (8) Å ³
$M_r = 604.51$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 13.3013$ (2) Å	$\mu = 0.53$ mm ⁻¹
$b = 14.2333$ (2) Å	$T = 293$ (2) K
$c = 15.6694$ (3) Å	$0.48 \times 0.42 \times 0.28$ mm

Data collection

Siemens SMART CCD diffractometer	24593 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2001)	7364 independent reflections
$(SADABS$; Sheldrick, 2001)	6364 reflections with $I > 2\sigma(I)$
$R_{\text{int}} = 0.049$	$R_{\text{int}} = 0.049$
$T_{\min} = 0.785$, $T_{\max} = 0.866$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	$\Delta\rho_{\max} = 0.19$ e Å ⁻³
$wR(F^2) = 0.072$	$\Delta\rho_{\min} = -0.30$ e Å ⁻³
$S = 1.03$	Absolute structure: Flack (1983), 3256 Friedel pairs
7364 reflections	Flack parameter: -0.001 (13)
361 parameters	H-atom parameters constrained

Table 1
Selected geometric parameters (Å, °).

Cr1–C1	1.851 (2)	Cr1–C4	1.901 (2)
Cr1–C2	1.8650 (19)	Cr1–P2	2.3736 (5)
Cr1–C3	1.872 (2)	Cr1–P3	2.3847 (5)
P2–Cr1–P3	91.389 (18)		

Data collection: *SMART* (Siemens, 1994); cell refinement: *SAINT* (Siemens, 1994); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS86* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We thank the Malaysian Government and Universiti Sains Malaysia for support (IRPA grant Nos. 09–02–05–0008 and 190–9609–2801).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KJ2106).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Bennett, M. J., Cotton, F. A. & LaPrade, M. D. (1971). *Acta Cryst.* **B27**, 1899–1904.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Jost, A., Rees, B. & Yelon, W. B. (1975). *Acta Cryst.* **B31**, 2649–2658.
- Shawkataly, O. bin, Islam, S. D., Fun, H.-K. & Didierjean, C. (2006). *Acta Cryst.* **E62**, m1086–m1087.
- Shawkataly, O. bin, Saminathan, T., Fun, H.-K. & Sivakumar, K. (1996). *Acta Cryst.* **C52**, 1352–1355.
- Shawkataly, O. bin, Umathavan, A., Ramalingam, K., Fun, H.-K. & Ibrahim, A. R. (1997). *Acta Cryst.* **C53**, 1543–1545.

- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1994). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Ueng, C.-H. & Shih, G.-Y. (1992). *Acta Cryst. C* **48**, 988–991.
- Whitaker, A. & Jeffery, J. W. (1967). *Acta Cryst. B* **23**, 977–984.

supplementary materials

Acta Cryst. (2009). E65, m250-m251 [doi:10.1107/S1600536809001202]

[1,3-Bis(diphenylphosphino)pentane- κ^2P,P']tetracarbonylchromium(0)

O. bin Shawkataly, D. T. Thangadurai, M. A. A. Pankhi, S. M. Shahinoor Dusal Islam and H.-K. Fun

Comment

It is generally believed that the metal (M) to carbon monoxide bond involves both $OC—M\sigma$ -bonding and $M—CO\pi$ -bonding. In view of this phenomenon, the bonding characteristics of metal carbonyls with a phosphine ligand in phosphine-substituted metal carbonyls are of interest. A search of the Cambridge Structural Database (Version 5.29; Allen, 2002) revealed only 88 complexes of group VI metal carbonyls with a 3-carbon backbone bidentate phosphine. However, there are only a few examples of chromium carbonyl complexes (Shawkataly *et al.*, 2006). Previously, we reported several crystal structures of phosphine-substituted group VI metal carbonyls (Shawkataly *et al.*, 1996, 1997). We present here the crystal structure of the title compound.

The title compound has an expected octahedral geometry (Fig. 1). The $Cr—C$ bond lengths of the *cis* carbonyl ligands (with respect to the P atom) are slightly longer than those for the *trans* carbonyl group (Table 1). This trend was also observed in $Cr[Ph_2P(CH_2)_2PPh_2](CO)_4$ (Bennett *et al.*, 1971) and $Cr[Ph_2P(CH_2)_4PPh_2](CO)_4$ (Ueng & Shih, 1992). The bidentate phosphine bite angle [91.389 (18) $^\circ$] is intermediate between that observed in $Cr[Ph_2P(CH_2)_2PPh_2](CO)_4$ (83.41 (8) $^\circ$) and that in $Cr[Ph_2P(CH_2)_4PPh_2](CO)_4$ (93.29 (5) $^\circ$). Comparison of the mean $Cr—C$ and $C—O$ distances in the title compound [1.872 (2) and 1.145 (6) \AA , respectively] with those in $Cr(CO)_6$ [1.909 (3) and 1.137 (4) \AA , respectively (Whitaker & Jeffery, 1967); and 1.918 (2) and 1.141 (2) \AA , respectively (Jost *et al.*, 1975)], indicates stronger bonding owing to the back-bonding abilities of the bidentate phosphine. The $Cr—P$ bond lengths, with an average values of 2.3792 (5) \AA , are relatively short inspite of the presence of the bulky phosphine ligand.

Experimental

A mixture of $Cr(CO)_6$ (1.064 mmol) and $Ph_2P(CH_3)CH(CH_2)CH(CH_3)PPh_2$ (1.065 mmol) was refluxed in a purified mixture of petroleum ether (60–80 $^\circ\text{C}$, 25 ml) and butanol (20 ml) for *ca* 12 h under nitrogen atmosphere. The solvent was evaporated and the crude product was dissolved in acetone (5 ml) and filtered. Yellow crystals (75% yield) were obtained by slow evaporation of the acetone solution at room temperature. Analysis calculated for $C_{33}H_{30}CrO_4P_2$: C 65.55, H 5.01%; found C 65.54, H 5.00%.

Refinement

All H atoms were placed at calculated positions and refined using a riding model, with $C—H = 0.93\text{--}0.98\text{\AA}$, $C—H = 0.97\text{\AA}$ (methylene) and $C—H = 0.96\text{\AA}$ (methyl) and $U_{iso}(H) = 1.2U_{eq}(C, \text{aromatic, methylene})$ and $U_{iso}(H) = 1.5U_{equ}(C \text{ methyl})$. A rotating group model was used for the methyl group. The number of Friedel pairs are 3260.

supplementary materials

Figures




Fig. 1. View of the title compound (50% probability displacement ellipsoids).

[2,4-Bis(diphenylphosphino)pentane- κ^2P,P']tetracarbonylchromium(0)

Crystal data

[Cr(C ₂₉ H ₃₀ P ₂)(CO) ₄]	$F_{000} = 1256$
$M_r = 604.51$	$D_x = 1.354 \text{ Mg m}^{-3}$
Orthorhombic, $P_{2_1}2_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 13.3013 (2) \text{ \AA}$	Cell parameters from 427 reflections
$b = 14.2333 (2) \text{ \AA}$	$\theta = 2\text{--}27.5^\circ$
$c = 15.6694 (3) \text{ \AA}$	$\mu = 0.53 \text{ mm}^{-1}$
$V = 2966.55 (8) \text{ \AA}^3$	$T = 293 \text{ K}$
$Z = 4$	Prism, yellow
	$0.48 \times 0.42 \times 0.28 \text{ mm}$

Data collection

Siemens SMART CCD diffractometer	7364 independent reflections
Radiation source: fine-focus sealed tube	6364 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.049$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	$h = -17 \rightarrow 12$
$T_{\text{min}} = 0.785, T_{\text{max}} = 0.866$	$k = -18 \rightarrow 18$
24593 measured reflections	$l = -18 \rightarrow 20$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.031$	$w = 1/[\sigma^2(F_o^2) + (0.0349P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$

$wR(F^2) = 0.072$	$(\Delta/\sigma)_{\max} = 0.001$
$S = 1.03$	$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
7364 reflections	$\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$
361 parameters	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 3256 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: -0.001 (13)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cr1	0.21688 (2)	0.11992 (2)	0.931300 (16)	0.03170 (7)
P2	0.33880 (3)	0.10714 (3)	0.82130 (3)	0.03095 (10)
P3	0.10424 (3)	0.01926 (3)	0.85607 (3)	0.03203 (10)
O1	0.35830 (13)	0.24334 (14)	1.02996 (12)	0.0745 (5)
O2	0.07737 (12)	0.14004 (13)	1.08126 (9)	0.0648 (5)
O3	0.29780 (14)	-0.05494 (12)	1.01576 (10)	0.0644 (4)
O4	0.11867 (14)	0.29523 (11)	0.85872 (12)	0.0696 (5)
C1	0.30535 (15)	0.19547 (15)	0.99132 (13)	0.0448 (5)
C2	0.12769 (14)	0.13194 (15)	1.02249 (12)	0.0434 (5)
C3	0.26815 (15)	0.01112 (15)	0.98236 (11)	0.0417 (4)
C4	0.15612 (16)	0.22871 (15)	0.88349 (12)	0.0436 (5)
C5	0.01810 (14)	0.06112 (14)	0.77239 (11)	0.0382 (4)
C6	0.05328 (15)	0.12712 (14)	0.71520 (12)	0.0425 (4)
H6	0.1137	0.1577	0.7262	0.051*
C7	0.00026 (19)	0.14868 (16)	0.64175 (14)	0.0573 (6)
H7	0.0263	0.1915	0.6028	0.069*
C8	-0.09146 (19)	0.10630 (19)	0.62657 (14)	0.0642 (7)
H8	-0.1276	0.1210	0.5776	0.077*
C9	-0.12896 (18)	0.0428 (2)	0.68371 (15)	0.0625 (7)
H9	-0.1911	0.0150	0.6736	0.075*
C10	-0.07513 (16)	0.01932 (17)	0.75678 (14)	0.0513 (5)
H10	-0.1012	-0.0241	0.7952	0.062*
C11	0.01881 (13)	-0.03718 (14)	0.93242 (12)	0.0390 (4)
C12	-0.06244 (15)	0.01504 (18)	0.96266 (13)	0.0523 (5)
H12	-0.0752	0.0743	0.9401	0.063*

supplementary materials

C13	-0.12452 (17)	-0.0206 (2)	1.02619 (15)	0.0691 (8)
H13	-0.1788	0.0144	1.0459	0.083*
C14	-0.10527 (18)	-0.1081 (2)	1.05985 (14)	0.0720 (8)
H14	-0.1470	-0.1322	1.1022	0.086*
C15	-0.0251 (2)	-0.1601 (2)	1.03155 (14)	0.0639 (7)
H15	-0.0126	-0.2190	1.0548	0.077*
C16	0.03757 (16)	-0.12478 (16)	0.96805 (12)	0.0475 (5)
H16	0.0922	-0.1600	0.9494	0.057*
C17	0.32879 (14)	0.20310 (13)	0.74299 (11)	0.0356 (4)
C18	0.29810 (16)	0.19096 (15)	0.65920 (13)	0.0486 (5)
H18	0.2886	0.1307	0.6378	0.058*
C19	0.2816 (2)	0.26759 (18)	0.60726 (16)	0.0643 (6)
H19	0.2594	0.2584	0.5516	0.077*
C20	0.29715 (19)	0.35629 (19)	0.63617 (18)	0.0676 (7)
H20	0.2856	0.4074	0.6006	0.081*
C21	0.33051 (18)	0.37031 (16)	0.71937 (18)	0.0620 (6)
H21	0.3426	0.4308	0.7392	0.074*
C22	0.34562 (16)	0.29421 (14)	0.77231 (14)	0.0477 (5)
H22	0.3672	0.3038	0.8281	0.057*
C23	0.47292 (13)	0.11750 (14)	0.84941 (12)	0.0374 (4)
C24	0.54190 (15)	0.14519 (15)	0.78829 (14)	0.0488 (5)
H24	0.5195	0.1622	0.7342	0.059*
C25	0.64357 (16)	0.14802 (18)	0.80620 (17)	0.0592 (6)
H25	0.6889	0.1676	0.7647	0.071*
C26	0.67731 (16)	0.12194 (17)	0.88523 (16)	0.0590 (6)
H26	0.7458	0.1230	0.8971	0.071*
C27	0.61055 (17)	0.09426 (16)	0.94707 (16)	0.0577 (6)
H27	0.6339	0.0768	1.0007	0.069*
C28	0.50785 (15)	0.09223 (15)	0.92967 (14)	0.0477 (5)
H28	0.4627	0.0739	0.9718	0.057*
C29	0.16603 (14)	-0.07883 (12)	0.79726 (11)	0.0348 (4)
H29	0.2055	-0.1151	0.8385	0.042*
C30	0.09214 (18)	-0.14639 (15)	0.75374 (14)	0.0520 (5)
H30A	0.0463	-0.1710	0.7954	0.078*
H30B	0.1287	-0.1972	0.7280	0.078*
H30C	0.0552	-0.1133	0.7106	0.078*
C31	0.23919 (13)	-0.04033 (13)	0.72887 (11)	0.0355 (4)
H31A	0.2505	-0.0899	0.6874	0.043*
H31B	0.2054	0.0105	0.6993	0.043*
C32	0.34278 (14)	-0.00379 (13)	0.75768 (11)	0.0356 (4)
H32	0.3806	0.0104	0.7056	0.043*
C33	0.40092 (16)	-0.08103 (14)	0.80401 (15)	0.0497 (5)
H33A	0.4652	-0.0572	0.8216	0.074*
H33B	0.4104	-0.1334	0.7663	0.074*
H33C	0.3637	-0.1009	0.8533	0.074*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cr1	0.02906 (13)	0.03611 (14)	0.02992 (12)	0.00083 (12)	-0.00075 (11)	-0.00429 (12)
P2	0.0271 (2)	0.0337 (2)	0.0320 (2)	0.00001 (19)	-0.00125 (17)	-0.00244 (18)
P3	0.0275 (2)	0.0372 (2)	0.0314 (2)	-0.00126 (18)	-0.00068 (18)	-0.00108 (19)
O1	0.0532 (10)	0.0809 (12)	0.0895 (13)	-0.0080 (9)	-0.0075 (9)	-0.0435 (11)
O2	0.0598 (9)	0.0915 (13)	0.0432 (8)	0.0009 (9)	0.0148 (7)	-0.0130 (8)
O3	0.0705 (11)	0.0641 (10)	0.0585 (9)	0.0125 (9)	0.0022 (8)	0.0236 (8)
O4	0.0799 (12)	0.0527 (10)	0.0763 (11)	0.0254 (9)	-0.0090 (10)	0.0008 (9)
C1	0.0381 (10)	0.0484 (11)	0.0480 (11)	0.0004 (9)	0.0006 (9)	-0.0132 (9)
C2	0.0410 (10)	0.0521 (12)	0.0370 (10)	0.0002 (9)	-0.0035 (8)	-0.0068 (9)
C3	0.0394 (10)	0.0519 (11)	0.0337 (9)	-0.0011 (9)	0.0015 (8)	0.0012 (8)
C4	0.0415 (11)	0.0472 (11)	0.0422 (10)	0.0042 (10)	-0.0019 (9)	-0.0086 (9)
C5	0.0335 (9)	0.0448 (10)	0.0362 (9)	0.0057 (8)	-0.0038 (8)	-0.0019 (8)
C6	0.0423 (10)	0.0425 (11)	0.0428 (10)	0.0046 (9)	-0.0039 (8)	0.0008 (9)
C7	0.0696 (15)	0.0531 (13)	0.0493 (12)	0.0118 (12)	-0.0021 (12)	0.0099 (11)
C8	0.0666 (15)	0.0781 (17)	0.0478 (12)	0.0153 (14)	-0.0230 (11)	-0.0005 (12)
C9	0.0440 (12)	0.0866 (18)	0.0569 (14)	-0.0002 (12)	-0.0165 (11)	-0.0009 (13)
C10	0.0398 (11)	0.0650 (14)	0.0491 (12)	-0.0049 (10)	-0.0078 (9)	0.0047 (11)
C11	0.0296 (8)	0.0541 (11)	0.0333 (9)	-0.0074 (8)	-0.0019 (8)	-0.0020 (9)
C12	0.0381 (10)	0.0736 (15)	0.0451 (11)	0.0030 (11)	0.0015 (9)	0.0009 (11)
C13	0.0354 (11)	0.124 (2)	0.0482 (12)	0.0020 (14)	0.0072 (10)	0.0005 (15)
C14	0.0499 (13)	0.125 (2)	0.0411 (11)	-0.0204 (16)	0.0055 (10)	0.0173 (15)
C15	0.0670 (16)	0.0786 (17)	0.0461 (12)	-0.0227 (14)	-0.0002 (12)	0.0155 (12)
C16	0.0479 (11)	0.0541 (12)	0.0406 (10)	-0.0079 (11)	0.0025 (9)	0.0027 (10)
C17	0.0290 (9)	0.0384 (10)	0.0394 (9)	-0.0028 (8)	0.0018 (8)	0.0020 (8)
C18	0.0485 (12)	0.0506 (11)	0.0465 (11)	-0.0083 (10)	-0.0059 (9)	0.0091 (9)
C19	0.0620 (14)	0.0724 (16)	0.0585 (13)	-0.0095 (14)	-0.0087 (12)	0.0269 (12)
C20	0.0587 (14)	0.0624 (15)	0.0818 (17)	0.0047 (12)	0.0057 (14)	0.0332 (14)
C21	0.0523 (13)	0.0380 (11)	0.0958 (19)	-0.0006 (10)	0.0202 (13)	0.0070 (12)
C22	0.0428 (11)	0.0426 (11)	0.0577 (12)	-0.0026 (9)	0.0070 (10)	-0.0020 (9)
C23	0.0286 (8)	0.0377 (9)	0.0458 (9)	0.0011 (8)	-0.0039 (7)	-0.0074 (9)
C24	0.0366 (10)	0.0571 (13)	0.0528 (12)	0.0000 (9)	-0.0012 (9)	-0.0013 (10)
C25	0.0312 (10)	0.0642 (15)	0.0822 (16)	-0.0029 (10)	0.0048 (11)	-0.0068 (13)
C26	0.0320 (10)	0.0591 (14)	0.0857 (16)	0.0036 (10)	-0.0140 (11)	-0.0225 (13)
C27	0.0479 (12)	0.0627 (14)	0.0624 (14)	0.0135 (11)	-0.0213 (11)	-0.0126 (11)
C28	0.0408 (10)	0.0550 (12)	0.0472 (11)	0.0045 (9)	-0.0063 (9)	-0.0056 (10)
C29	0.0337 (9)	0.0345 (9)	0.0362 (9)	-0.0024 (8)	0.0010 (8)	-0.0023 (7)
C30	0.0532 (12)	0.0475 (12)	0.0553 (12)	-0.0159 (10)	0.0072 (10)	-0.0128 (10)
C31	0.0350 (9)	0.0388 (10)	0.0327 (8)	-0.0024 (7)	0.0014 (7)	-0.0058 (7)
C32	0.0302 (8)	0.0393 (9)	0.0373 (9)	0.0003 (8)	0.0057 (7)	-0.0054 (8)
C33	0.0403 (11)	0.0421 (11)	0.0665 (14)	0.0091 (9)	-0.0008 (10)	-0.0081 (10)

Geometric parameters (\AA , $^\circ$)

Cr1—C1	1.851 (2)	C16—H16	0.9300
Cr1—C2	1.8650 (19)	C17—C18	1.386 (3)

supplementary materials

Cr1—C3	1.872 (2)	C17—C22	1.394 (3)
Cr1—C4	1.901 (2)	C18—C19	1.379 (3)
Cr1—P2	2.3736 (5)	C18—H18	0.9300
Cr1—P3	2.3847 (5)	C19—C20	1.357 (4)
P2—C17	1.8409 (19)	C19—H19	0.9300
P2—C23	1.8434 (17)	C20—C21	1.391 (4)
P2—C32	1.8680 (18)	C20—H20	0.9300
P3—C11	1.8352 (19)	C21—C22	1.379 (3)
P3—C5	1.8405 (18)	C21—H21	0.9300
P3—C29	1.8637 (18)	C22—H22	0.9300
O1—C1	1.152 (2)	C23—C24	1.384 (3)
O2—C2	1.144 (2)	C23—C28	1.388 (3)
O3—C3	1.146 (2)	C24—C25	1.382 (3)
O4—C4	1.138 (2)	C24—H24	0.9300
C5—C6	1.380 (3)	C25—C26	1.368 (3)
C5—C10	1.397 (3)	C25—H25	0.9300
C6—C7	1.384 (3)	C26—C27	1.372 (3)
C6—H6	0.9300	C26—H26	0.9300
C7—C8	1.382 (3)	C27—C28	1.393 (3)
C7—H7	0.9300	C27—H27	0.9300
C8—C9	1.367 (3)	C28—H28	0.9300
C8—H8	0.9300	C29—C30	1.535 (3)
C9—C10	1.391 (3)	C29—C31	1.548 (2)
C9—H9	0.9300	C29—H29	0.9800
C10—H10	0.9300	C30—H30A	0.9600
C11—C16	1.389 (3)	C30—H30B	0.9600
C11—C12	1.395 (3)	C30—H30C	0.9600
C12—C13	1.389 (3)	C31—C32	1.540 (2)
C12—H12	0.9300	C31—H31A	0.9700
C13—C14	1.376 (4)	C31—H31B	0.9700
C13—H13	0.9300	C32—C33	1.528 (3)
C14—C15	1.371 (4)	C32—H32	0.9800
C14—H14	0.9300	C33—H33A	0.9600
C15—C16	1.392 (3)	C33—H33B	0.9600
C15—H15	0.9300	C33—H33C	0.9600
C1—Cr1—C2	87.81 (8)	C18—C17—C22	118.39 (18)
C1—Cr1—C3	91.82 (9)	C18—C17—P2	124.08 (15)
C2—Cr1—C3	88.86 (9)	C22—C17—P2	117.33 (15)
C1—Cr1—C4	89.84 (9)	C19—C18—C17	120.5 (2)
C2—Cr1—C4	87.52 (9)	C19—C18—H18	119.7
C3—Cr1—C4	175.96 (9)	C17—C18—H18	119.7
C1—Cr1—P2	88.80 (6)	C20—C19—C18	121.0 (2)
C2—Cr1—P2	176.35 (6)	C20—C19—H19	119.5
C3—Cr1—P2	89.89 (6)	C18—C19—H19	119.5
C4—Cr1—P2	93.83 (6)	C19—C20—C21	119.7 (2)
C1—Cr1—P3	178.55 (7)	C19—C20—H20	120.2
C2—Cr1—P3	91.96 (6)	C21—C20—H20	120.2
C3—Cr1—P3	86.74 (6)	C22—C21—C20	119.8 (2)
C4—Cr1—P3	91.58 (6)	C22—C21—H21	120.1

P2—Cr1—P3	91.389 (18)	C20—C21—H21	120.1
C17—P2—C23	99.78 (9)	C21—C22—C17	120.6 (2)
C17—P2—C32	105.87 (8)	C21—C22—H22	119.7
C23—P2—C32	99.66 (8)	C17—C22—H22	119.7
C17—P2—Cr1	112.20 (6)	C24—C23—C28	118.61 (17)
C23—P2—Cr1	118.79 (6)	C24—C23—P2	119.95 (14)
C32—P2—Cr1	118.14 (6)	C28—C23—P2	121.31 (15)
C11—P3—C5	102.74 (9)	C25—C24—C23	121.1 (2)
C11—P3—C29	105.52 (9)	C25—C24—H24	119.4
C5—P3—C29	99.48 (9)	C23—C24—H24	119.4
C11—P3—Cr1	109.25 (6)	C26—C25—C24	119.8 (2)
C5—P3—Cr1	123.28 (7)	C26—C25—H25	120.1
C29—P3—Cr1	114.67 (6)	C24—C25—H25	120.1
O1—C1—Cr1	178.18 (18)	C25—C26—C27	120.3 (2)
O2—C2—Cr1	176.29 (17)	C25—C26—H26	119.8
O3—C3—Cr1	177.93 (18)	C27—C26—H26	119.8
O4—C4—Cr1	176.72 (18)	C26—C27—C28	120.1 (2)
C6—C5—C10	118.50 (18)	C26—C27—H27	119.9
C6—C5—P3	118.16 (14)	C28—C27—H27	119.9
C10—C5—P3	122.65 (16)	C23—C28—C27	120.0 (2)
C5—C6—C7	121.2 (2)	C23—C28—H28	120.0
C5—C6—H6	119.4	C27—C28—H28	120.0
C7—C6—H6	119.4	C30—C29—C31	108.47 (15)
C8—C7—C6	119.8 (2)	C30—C29—P3	113.99 (14)
C8—C7—H7	120.1	C31—C29—P3	110.76 (13)
C6—C7—H7	120.1	C30—C29—H29	107.8
C9—C8—C7	119.9 (2)	C31—C29—H29	107.8
C9—C8—H8	120.1	P3—C29—H29	107.8
C7—C8—H8	120.1	C29—C30—H30A	109.5
C8—C9—C10	120.7 (2)	C29—C30—H30B	109.5
C8—C9—H9	119.7	H30A—C30—H30B	109.5
C10—C9—H9	119.7	C29—C30—H30C	109.5
C9—C10—C5	119.9 (2)	H30A—C30—H30C	109.5
C9—C10—H10	120.0	H30B—C30—H30C	109.5
C5—C10—H10	120.0	C32—C31—C29	118.62 (15)
C16—C11—C12	118.76 (19)	C32—C31—H31A	107.7
C16—C11—P3	122.94 (15)	C29—C31—H31A	107.7
C12—C11—P3	117.91 (16)	C32—C31—H31B	107.7
C13—C12—C11	120.6 (2)	C29—C31—H31B	107.7
C13—C12—H12	119.7	H31A—C31—H31B	107.1
C11—C12—H12	119.7	C33—C32—C31	110.44 (16)
C14—C13—C12	119.6 (2)	C33—C32—P2	111.66 (13)
C14—C13—H13	120.2	C31—C32—P2	114.61 (12)
C12—C13—H13	120.2	C33—C32—H32	106.5
C15—C14—C13	120.6 (2)	C31—C32—H32	106.5
C15—C14—H14	119.7	P2—C32—H32	106.5
C13—C14—H14	119.7	C32—C33—H33A	109.5
C14—C15—C16	120.2 (3)	C32—C33—H33B	109.5
C14—C15—H15	119.9	H33A—C33—H33B	109.5

supplementary materials

C16—C15—H15	119.9	C32—C33—H33C	109.5
C11—C16—C15	120.2 (2)	H33A—C33—H33C	109.5
C11—C16—H16	119.9	H33B—C33—H33C	109.5
C15—C16—H16	119.9		
C1—Cr1—P2—C17	−87.62 (9)	C13—C14—C15—C16	−0.2 (4)
C3—Cr1—P2—C17	−179.44 (9)	C12—C11—C16—C15	1.3 (3)
C4—Cr1—P2—C17	2.15 (9)	P3—C11—C16—C15	173.94 (17)
P3—Cr1—P2—C17	93.82 (7)	C14—C15—C16—C11	−0.6 (3)
C1—Cr1—P2—C23	28.07 (10)	C23—P2—C17—C18	121.48 (17)
C3—Cr1—P2—C23	−63.75 (10)	C32—P2—C17—C18	18.42 (19)
C4—Cr1—P2—C23	117.84 (10)	Cr1—P2—C17—C18	−111.79 (16)
P3—Cr1—P2—C23	−150.49 (8)	C23—P2—C17—C22	−63.80 (17)
C1—Cr1—P2—C32	148.80 (10)	C32—P2—C17—C22	−166.86 (15)
C3—Cr1—P2—C32	56.98 (9)	Cr1—P2—C17—C22	62.93 (16)
C4—Cr1—P2—C32	−121.44 (9)	C22—C17—C18—C19	−2.1 (3)
P3—Cr1—P2—C32	−29.76 (7)	P2—C17—C18—C19	172.58 (18)
C2—Cr1—P3—C11	−23.62 (10)	C17—C18—C19—C20	1.5 (4)
C3—Cr1—P3—C11	65.13 (9)	C18—C19—C20—C21	0.2 (4)
C4—Cr1—P3—C11	−111.19 (9)	C19—C20—C21—C22	−1.3 (4)
P2—Cr1—P3—C11	154.94 (7)	C20—C21—C22—C17	0.7 (3)
C2—Cr1—P3—C5	96.96 (10)	C18—C17—C22—C21	1.0 (3)
C3—Cr1—P3—C5	−174.29 (9)	P2—C17—C22—C21	−174.05 (17)
C4—Cr1—P3—C5	9.39 (10)	C17—P2—C23—C24	−33.81 (18)
P2—Cr1—P3—C5	−84.48 (8)	C32—P2—C23—C24	74.30 (18)
C2—Cr1—P3—C29	−141.81 (9)	Cr1—P2—C23—C24	−155.96 (14)
C3—Cr1—P3—C29	−53.06 (9)	C17—P2—C23—C28	150.40 (17)
C4—Cr1—P3—C29	130.62 (9)	C32—P2—C23—C28	−101.49 (17)
P2—Cr1—P3—C29	36.75 (7)	Cr1—P2—C23—C28	28.25 (19)
C11—P3—C5—C6	164.12 (15)	C28—C23—C24—C25	−0.1 (3)
C29—P3—C5—C6	−87.46 (16)	P2—C23—C24—C25	−176.02 (18)
Cr1—P3—C5—C6	40.56 (18)	C23—C24—C25—C26	0.9 (4)
C11—P3—C5—C10	−25.5 (2)	C24—C25—C26—C27	−0.9 (4)
C29—P3—C5—C10	82.87 (18)	C25—C26—C27—C28	0.2 (4)
Cr1—P3—C5—C10	−149.11 (15)	C24—C23—C28—C27	−0.6 (3)
C10—C5—C6—C7	−3.0 (3)	P2—C23—C28—C27	175.27 (16)
P3—C5—C6—C7	167.71 (16)	C26—C27—C28—C23	0.5 (3)
C5—C6—C7—C8	2.5 (3)	C11—P3—C29—C30	56.18 (16)
C6—C7—C8—C9	−0.6 (4)	C5—P3—C29—C30	−49.99 (16)
C7—C8—C9—C10	−0.8 (4)	Cr1—P3—C29—C30	176.46 (12)
C8—C9—C10—C5	0.2 (4)	C11—P3—C29—C31	178.84 (12)
C6—C5—C10—C9	1.7 (3)	C5—P3—C29—C31	72.67 (14)
P3—C5—C10—C9	−168.63 (18)	Cr1—P3—C29—C31	−60.89 (13)
C5—P3—C11—C16	134.04 (16)	C30—C29—C31—C32	−156.37 (17)
C29—P3—C11—C16	30.26 (18)	P3—C29—C31—C32	77.82 (19)
Cr1—P3—C11—C16	−93.51 (16)	C29—C31—C32—C33	58.3 (2)
C5—P3—C11—C12	−53.21 (17)	C29—C31—C32—P2	−68.9 (2)
C29—P3—C11—C12	−156.99 (15)	C17—P2—C32—C33	151.59 (14)
Cr1—P3—C11—C12	79.24 (15)	C23—P2—C32—C33	48.44 (15)
C16—C11—C12—C13	−1.1 (3)	Cr1—P2—C32—C33	−81.72 (14)

supplementary materials

P3—C11—C12—C13	-174.20 (17)	C17—P2—C32—C31	-81.88 (14)
C11—C12—C13—C14	0.3 (4)	C23—P2—C32—C31	174.97 (14)
C12—C13—C14—C15	0.4 (4)	Cr1—P2—C32—C31	44.80 (15)

supplementary materials

Fig. 1

