

trans,trans,trans-Diaquabis(nicotinamide- κN)bis(2-nitrobenzoato- κO)cadmium(II) dihydrate

Kou-Lin Zhang,^a Bo Yang,^a Jian-Guo Lin^a and Seik Weng Ng^{b*}

^aCollege of Chemistry and Chemical Engineering, Yangzhou University, Yangzhou 225002, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603, Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 21 January 2009; accepted 16 February 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.034; wR factor = 0.093; data-to-parameter ratio = 10.8.

The cadmium atom in the title compound, $[\text{Cd}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$, lies on a center of inversion in an all-*trans* octahedral environment. In the crystal, the complex interacts with the uncoordinated water molecules through $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a layered network.

Related literature

There are several examples of diaquadi(arylcarboxylato)di(nicotinamide)metal(II) compounds. For recent examples, see: Hökelek & Necefoğlu (2007a,b); Hökelek *et al.* (2007); Koksharova *et al.* (2006); Şahin *et al.* (2007a,b); Stachova *et al.* (2006); Çaylak *et al.* (2007).



Experimental

Crystal data

$[\text{Cd}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$
 $M_r = 760.94$
Monoclinic, $P2_1/n$
 $a = 7.9365 (8)\text{ \AA}$

$b = 19.589 (2)\text{ \AA}$
 $c = 10.059 (1)\text{ \AA}$
 $\beta = 103.178 (2)^\circ$
 $V = 1522.6 (3)\text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.80\text{ mm}^{-1}$

$T = 293\text{ K}$
 $0.50 \times 0.18 \times 0.18\text{ mm}$

Data collection

Bruker SMART area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.619$, $T_{\max} = 0.866$

4427 measured reflections
2651 independent reflections
2396 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.093$
 $S = 1.13$
2651 reflections
246 parameters
9 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.42\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.04\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1w-H11 \cdots O2w	0.85 (4)	1.92 (4)	2.764 (4)	174 (4)
O1w-H12 \cdots O2 ⁱ	0.85 (4)	1.95 (5)	2.718 (4)	150 (4)
O2w-H21 \cdots O5 ⁱ	0.85 (3)	2.08 (3)	2.910 (4)	166 (4)
O2w-H22 \cdots O1 ⁱⁱ	0.85 (3)	2.00 (1)	2.846 (3)	177 (5)
N3-H31 \cdots O2 ⁱⁱⁱ	0.85 (3)	2.22 (2)	3.038 (4)	165 (4)
N3-H32 \cdots O5 ^{iv}	0.85 (3)	2.05 (3)	2.873 (4)	164 (4)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$; (iii) $x, y, z + 1$; (iv) $-x, -y + 1, -z + 2$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Foundation of Jiangsu Provincial Key Program of Physical Chemistry in Yangzhou University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2136).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.
- Bruker (2000). *SAINT* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Çaylak, N., Hökelek, T. & Necefoğlu, H. (2007). *Acta Cryst. E63*, m1341–m1343.
- Hökelek, T., Çaylak, N. & Necefoğlu, H. (2007). *Acta Cryst. E63*, m1873–m1874.
- Hökelek, T. & Necefoğlu, H. (2007a). *Acta Cryst. E63*, m1078–m1080.
- Hökelek, T. & Necefoğlu, H. (2007b). *Acta Cryst. E63*, m1279–m1281.
- Koksharova, T. V., Sadikov, G. G., Antsyshkina, A. S., Gritsenko, I. S., Sergienko, V. S. & Egorova, O. A. (2006). *Russ. J. Inorg. Chem.* **51**, 895–900.
- Şahin, O., Büyükgüngör, O., Köse, D. A. & Necefoğlu, H. (2007a). *Acta Cryst. C63*, m510–m512.
- Şahin, O., Büyükgüngör, O., Köse, D. A., Ozturkkan, E. F. & Necefoğlu, H. (2007b). *Acta Cryst. C63*, m243–m245.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stachova, P., Melnik, M., Korobik, M., Mrozinski, M., Koman, M., Glowiacz, T. & Valigura, D. (2006). *Inorg. Chim. Acta*, **360**, 1517–1522.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m292 [doi:10.1107/S1600536809005479]

trans,trans,trans-Diaquabis(nicotinamide- κN)bis(2-nitrobenzoato- κO)cadmium(II) dihydrate

K.-L. Zhang, B. Yang, J.-G. Lin and S. W. Ng

Experimental

A water/methanol (1:1 v/v) solution (3 ml) of cadmium nitrate trihydrate (0.082 g, 0.3 mmol) was added to a water/methanol (1:1 v/v) solution (3 ml) of 2-nitrobenzoic acid (0.100 g, 0.6 mmol), sodium hydroxide (0.024 g, 0.6 mmol) and nicotinamide (0.073 g, 0.6 mmol). A white powder was obtained after several days; this was recrystallized from DMF/methanol (3:1 v/v) to give colorless crystals in 50% yield. CH&N elemental analysis. Calculated for $C_{26}H_{28}CdN_6O_{14}$: C 41.04 H 3.68 N 11.04%; found: C 40.08, H 3.87, N 10.93%.

Refinement

Carbon-bound H atoms were placed in calculated positions and were allowed to ride on the parent atoms. N and O-bound H atoms were located in a difference Fourier map, and were refined with distance restraints $N-H = O-H = 0.85 \pm 0.01 \text{ \AA}$; for the water molecules, an additional $H \cdots H = 1.39 \pm 0.01 \text{ \AA}$ restraint was used. Their temperature factors were freely refined.

The measurements are 100% at the 2θ limit of 50 °.

Figures




Fig. 1. Thermal ellipsoid plot of $Cd(H_2O)_2(C_7H_4NO_4)_2(C_6H_6N_2O)_2 \cdot 2H_2O$; displacement ellipsoids are drawn at the 50% probability level, and H atoms as spheres of arbitrary radii.

trans,trans,trans-Diaquabis(nicotinamide- κN)bis(2-nitrobenzoato- κO)cadmium(II) dihydrate

Crystal data

$[Cd(C_7H_4NO_4)_2(C_6H_6N_2O)_2(H_2O)_2] \cdot 2H_2O$	$F_{000} = 772$
$M_r = 760.94$	$D_x = 1.660 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 7.9365 (8) \text{ \AA}$	Cell parameters from 3417 reflections
$b = 19.589 (2) \text{ \AA}$	$\theta = 2.1\text{--}25.1^\circ$
$c = 10.059 (1) \text{ \AA}$	$\mu = 0.80 \text{ mm}^{-1}$
$\beta = 103.178 (2)^\circ$	$T = 293 \text{ K}$
$V = 1522.6 (3) \text{ \AA}^3$	Rod, colorless
$Z = 2$	$0.50 \times 0.18 \times 0.18 \text{ mm}$

supplementary materials

Data collection

Bruker SMART area-detector diffractometer	2651 independent reflections
Radiation source: medium-focus sealed tube	2396 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.016$
$T = 295 \text{ K}$	$\theta_{\text{max}} = 25.1^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 3$
$T_{\text{min}} = 0.619, T_{\text{max}} = 0.866$	$k = -20 \rightarrow 23$
4427 measured reflections	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.0433P)^2 + 2.3534P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.13$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2651 reflections	$\Delta\rho_{\text{max}} = 0.42 \text{ e \AA}^{-3}$
246 parameters	$\Delta\rho_{\text{min}} = -1.04 \text{ e \AA}^{-3}$
9 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.5000	0.5000	0.5000	0.02349 (13)
O1	0.6570 (3)	0.40982 (12)	0.4385 (2)	0.0312 (5)
O2	0.4256 (3)	0.36571 (14)	0.2970 (3)	0.0428 (7)
O3	0.3595 (4)	0.23243 (19)	0.1217 (4)	0.0693 (10)
O4	0.4552 (5)	0.2124 (2)	0.3355 (4)	0.0835 (12)
O5	0.0789 (4)	0.48477 (17)	0.8538 (3)	0.0505 (8)
O1W	0.7685 (3)	0.55083 (13)	0.5819 (3)	0.0362 (6)
H11	0.842 (5)	0.561 (2)	0.536 (4)	0.055 (14)*
H12	0.737 (6)	0.5868 (14)	0.616 (5)	0.082 (19)*
O2W	0.9923 (3)	0.57871 (14)	0.4149 (3)	0.0375 (6)
H21	0.987 (5)	0.556 (2)	0.342 (3)	0.079 (18)*
H22	1.097 (2)	0.581 (2)	0.460 (3)	0.050 (13)*
N1	0.4713 (4)	0.23414 (16)	0.2260 (4)	0.0438 (8)
N2	0.5129 (3)	0.44979 (15)	0.7119 (3)	0.0291 (6)
N3	0.1940 (4)	0.44256 (19)	1.0627 (3)	0.0405 (8)

H31	0.274 (4)	0.424 (2)	1.122 (3)	0.055 (13)*
H32	0.114 (4)	0.457 (2)	1.098 (3)	0.049 (12)*
C1	0.6994 (4)	0.32348 (16)	0.2832 (3)	0.0237 (6)
C2	0.6421 (4)	0.26190 (17)	0.2198 (3)	0.0282 (7)
C3	0.7386 (5)	0.22316 (19)	0.1503 (4)	0.0386 (9)
H3	0.6942	0.1829	0.1070	0.047 (12)*
C4	0.9033 (5)	0.2454 (2)	0.1459 (4)	0.0407 (9)
H4	0.9714	0.2196	0.1008	0.047 (12)*
C5	0.9655 (5)	0.3057 (2)	0.2087 (4)	0.0430 (9)
H5	1.0761	0.3205	0.2063	0.057 (13)*
C6	0.8643 (4)	0.34455 (19)	0.2755 (4)	0.0337 (8)
H6	0.9075	0.3855	0.3161	0.046 (12)*
C7	0.5842 (4)	0.36885 (16)	0.3458 (3)	0.0262 (7)
C8	0.3792 (4)	0.46351 (17)	0.7674 (3)	0.0256 (7)
H8	0.2975	0.4951	0.7238	0.044 (12)*
C9	0.3553 (4)	0.43356 (17)	0.8857 (3)	0.0273 (7)
C10	0.4767 (5)	0.3864 (2)	0.9499 (4)	0.0399 (9)
H10	0.4645	0.3648	1.0295	0.044 (11)*
C11	0.6165 (5)	0.3721 (2)	0.8941 (4)	0.0461 (10)
H11A	0.7004	0.3410	0.9362	0.055 (13)*
C12	0.6301 (5)	0.40434 (19)	0.7751 (4)	0.0352 (8)
H12A	0.7239	0.3942	0.7375	0.052 (12)*
C13	0.1972 (4)	0.45472 (18)	0.9337 (3)	0.0312 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.02157 (19)	0.0277 (2)	0.0224 (2)	-0.00033 (12)	0.00743 (13)	0.00046 (12)
O1	0.0292 (12)	0.0324 (13)	0.0303 (12)	0.0020 (10)	0.0034 (10)	-0.0075 (10)
O2	0.0268 (13)	0.0550 (17)	0.0443 (15)	0.0049 (12)	0.0033 (11)	-0.0207 (13)
O3	0.0398 (17)	0.078 (2)	0.085 (3)	-0.0141 (16)	0.0042 (17)	-0.027 (2)
O4	0.087 (3)	0.098 (3)	0.078 (3)	-0.041 (2)	0.044 (2)	0.003 (2)
O5	0.0388 (16)	0.084 (2)	0.0318 (15)	0.0297 (15)	0.0135 (12)	0.0122 (14)
O1W	0.0273 (13)	0.0406 (14)	0.0416 (15)	-0.0066 (11)	0.0094 (11)	-0.0085 (12)
O2W	0.0276 (13)	0.0452 (15)	0.0390 (15)	0.0019 (11)	0.0061 (11)	-0.0039 (12)
N1	0.0417 (19)	0.0363 (18)	0.057 (2)	-0.0114 (14)	0.0190 (17)	-0.0155 (16)
N2	0.0251 (14)	0.0348 (15)	0.0274 (15)	0.0037 (12)	0.0059 (11)	0.0027 (12)
N3	0.0366 (17)	0.066 (2)	0.0223 (15)	0.0158 (16)	0.0128 (13)	0.0093 (15)
C1	0.0281 (16)	0.0240 (16)	0.0173 (14)	0.0011 (13)	0.0015 (12)	-0.0017 (12)
C2	0.0282 (17)	0.0258 (17)	0.0313 (17)	-0.0024 (13)	0.0083 (14)	-0.0016 (14)
C3	0.051 (2)	0.0263 (18)	0.041 (2)	-0.0017 (16)	0.0154 (17)	-0.0088 (16)
C4	0.042 (2)	0.040 (2)	0.045 (2)	0.0102 (17)	0.0185 (17)	-0.0024 (17)
C5	0.0305 (19)	0.047 (2)	0.056 (3)	-0.0013 (17)	0.0203 (18)	-0.0033 (19)
C6	0.0303 (18)	0.0349 (19)	0.0359 (19)	-0.0044 (15)	0.0075 (15)	-0.0083 (15)
C7	0.0254 (16)	0.0243 (16)	0.0297 (17)	0.0019 (13)	0.0078 (13)	0.0013 (13)
C8	0.0246 (16)	0.0309 (18)	0.0201 (15)	0.0033 (13)	0.0025 (12)	0.0029 (13)
C9	0.0282 (17)	0.0336 (18)	0.0199 (15)	0.0028 (14)	0.0051 (13)	-0.0012 (13)
C10	0.042 (2)	0.053 (2)	0.0267 (18)	0.0173 (18)	0.0117 (15)	0.0150 (17)

supplementary materials

C11	0.041 (2)	0.062 (3)	0.035 (2)	0.027 (2)	0.0095 (17)	0.0190 (19)
C12	0.0303 (18)	0.046 (2)	0.0317 (19)	0.0088 (16)	0.0117 (15)	0.0022 (16)
C13	0.0304 (18)	0.0375 (19)	0.0274 (17)	0.0051 (15)	0.0098 (14)	0.0027 (14)

Geometric parameters (\AA , $^\circ$)

Cd1—O1 ⁱ	2.325 (2)	C1—C2	1.391 (4)
Cd1—O1	2.325 (2)	C1—C6	1.391 (5)
Cd1—O1W ⁱ	2.326 (2)	C1—C7	1.512 (4)
Cd1—O1W	2.326 (2)	C2—C3	1.377 (5)
Cd1—N2	2.329 (3)	C3—C4	1.388 (5)
Cd1—N2 ⁱ	2.329 (3)	C3—H3	0.9300
O1—C7	1.266 (4)	C4—C5	1.378 (6)
O2—C7	1.244 (4)	C4—H4	0.9300
O3—N1	1.211 (5)	C5—C6	1.386 (5)
O4—N1	1.214 (5)	C5—H5	0.9300
O5—C13	1.237 (4)	C6—H6	0.9300
O1W—H11	0.85 (4)	C8—C9	1.378 (5)
O1W—H12	0.85 (4)	C8—H8	0.9300
O2W—H21	0.85 (3)	C9—C10	1.383 (5)
O2W—H22	0.85 (3)	C9—C13	1.502 (5)
N1—C2	1.476 (4)	C10—C11	1.382 (5)
N2—C8	1.334 (4)	C10—H10	0.9300
N2—C12	1.339 (4)	C11—C12	1.379 (5)
N3—C13	1.325 (4)	C11—H11A	0.9300
N3—H31	0.84 (3)	C12—H12A	0.9300
N3—H32	0.85 (3)		
O1 ⁱ —Cd1—O1	180.000 (1)	C1—C2—N1	120.5 (3)
O1 ⁱ —Cd1—O1W ⁱ	85.20 (9)	C2—C3—C4	118.7 (3)
O1—Cd1—O1W ⁱ	94.80 (9)	C2—C3—H3	120.7
O1 ⁱ —Cd1—O1W	94.80 (9)	C4—C3—H3	120.7
O1—Cd1—O1W	85.20 (9)	C5—C4—C3	119.8 (3)
O1W ⁱ —Cd1—O1W	180.00 (12)	C5—C4—H4	120.1
O1 ⁱ —Cd1—N2	89.52 (9)	C3—C4—H4	120.1
O1—Cd1—N2	90.48 (9)	C4—C5—C6	120.4 (3)
O1W ⁱ —Cd1—N2	89.36 (10)	C4—C5—H5	119.8
O1W—Cd1—N2	90.64 (10)	C6—C5—H5	119.8
O1 ⁱ —Cd1—N2 ⁱ	90.48 (9)	C5—C6—C1	121.4 (3)
O1—Cd1—N2 ⁱ	89.52 (9)	C5—C6—H6	119.3
O1W ⁱ —Cd1—N2 ⁱ	90.64 (10)	C1—C6—H6	119.3
O1W—Cd1—N2 ⁱ	89.36 (10)	O2—C7—O1	125.0 (3)
N2—Cd1—N2 ⁱ	180.0	O2—C7—C1	117.4 (3)
C7—O1—Cd1	119.7 (2)	O1—C7—C1	117.5 (3)
Cd1—O1W—H11	126 (3)	N2—C8—C9	123.7 (3)
Cd1—O1W—H12	100 (3)	N2—C8—H8	118.2
H11—O1W—H12	109 (4)	C9—C8—H8	118.2

H21—O2W—H22	109.4 (17)	C8—C9—C10	118.0 (3)
O3—N1—O4	124.7 (4)	C8—C9—C13	116.7 (3)
O3—N1—C2	118.2 (4)	C10—C9—C13	125.3 (3)
O4—N1—C2	117.1 (4)	C11—C10—C9	118.9 (3)
C8—N2—C12	117.9 (3)	C11—C10—H10	120.5
C8—N2—Cd1	115.1 (2)	C9—C10—H10	120.5
C12—N2—Cd1	126.6 (2)	C12—C11—C10	119.3 (3)
C13—N3—H31	126 (3)	C12—C11—H11A	120.4
C13—N3—H32	122 (2)	C10—C11—H11A	120.4
H31—N3—H32	111.4 (18)	N2—C12—C11	122.2 (3)
C2—C1—C6	116.4 (3)	N2—C12—H12A	118.9
C2—C1—C7	122.4 (3)	C11—C12—H12A	118.9
C6—C1—C7	121.0 (3)	O5—C13—N3	122.8 (3)
C3—C2—C1	123.3 (3)	O5—C13—C9	119.2 (3)
C3—C2—N1	116.2 (3)	N3—C13—C9	118.0 (3)
O1W ⁱ —Cd1—O1—C7	22.2 (2)	C4—C5—C6—C1	1.0 (6)
O1W—Cd1—O1—C7	-157.8 (2)	C2—C1—C6—C5	-0.3 (5)
N2—Cd1—O1—C7	111.6 (2)	C7—C1—C6—C5	-174.8 (3)
N2 ⁱ —Cd1—O1—C7	-68.4 (2)	Cd1—O1—C7—O2	-11.8 (5)
O1 ⁱ —Cd1—N2—C8	29.7 (2)	Cd1—O1—C7—C1	163.9 (2)
O1—Cd1—N2—C8	-150.3 (2)	C2—C1—C7—O2	-27.5 (5)
O1W ⁱ —Cd1—N2—C8	-55.5 (2)	C6—C1—C7—O2	146.7 (3)
O1W—Cd1—N2—C8	124.5 (2)	C2—C1—C7—O1	156.4 (3)
O1 ⁱ —Cd1—N2—C12	-157.2 (3)	C6—C1—C7—O1	-29.4 (4)
O1—Cd1—N2—C12	22.8 (3)	C12—N2—C8—C9	0.0 (5)
O1W ⁱ —Cd1—N2—C12	117.6 (3)	Cd1—N2—C8—C9	173.7 (3)
O1W—Cd1—N2—C12	-62.4 (3)	N2—C8—C9—C10	-0.3 (5)
C6—C1—C2—C3	-1.2 (5)	N2—C8—C9—C13	-179.8 (3)
C7—C1—C2—C3	173.2 (3)	C8—C9—C10—C11	0.7 (6)
C6—C1—C2—N1	178.0 (3)	C13—C9—C10—C11	-179.9 (4)
C7—C1—C2—N1	-7.5 (5)	C9—C10—C11—C12	-0.8 (7)
O3—N1—C2—C3	-69.1 (5)	C8—N2—C12—C11	-0.1 (6)
O4—N1—C2—C3	108.4 (4)	Cd1—N2—C12—C11	-173.0 (3)
O3—N1—C2—C1	111.6 (4)	C10—C11—C12—N2	0.6 (7)
O4—N1—C2—C1	-70.9 (5)	C8—C9—C13—O5	15.8 (5)
C1—C2—C3—C4	1.9 (6)	C10—C9—C13—O5	-163.7 (4)
N1—C2—C3—C4	-177.4 (3)	C8—C9—C13—N3	-161.5 (3)
C2—C3—C4—C5	-1.1 (6)	C10—C9—C13—N3	19.0 (6)
C3—C4—C5—C6	-0.3 (6)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1w—H11···O2w	0.85 (4)	1.92 (4)	2.764 (4)	174 (4)
O1w—H12···O2 ⁱ	0.85 (4)	1.95 (5)	2.718 (4)	150 (4)
O2w—H21···O5 ⁱ	0.85 (3)	2.08 (3)	2.910 (4)	166 (4)

supplementary materials

O2w—H22···O1 ⁱⁱ	0.85 (3)	2.00 (1)	2.846 (3)	177 (5)
N3—H31···O2 ⁱⁱⁱ	0.85 (3)	2.22 (2)	3.038 (4)	165 (4)
N3—H32···O5 ^{iv}	0.85 (3)	2.05 (3)	2.873 (4)	164 (4)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+1, -z+1$; (iii) $x, y, z+1$; (iv) $-x, -y+1, -z+2$.

Fig. 1

